

Vol. 39 No. 2 2026

Print ISSN: 0971-8184

Online ISSN: 0976-1926

Indian Journal of Plant Genetic Resources

*An International Journal for
Conservation and Use of Plant Diversity*



Indian Society of Plant Genetic Resources
New Delhi, India

INDIAN SOCIETY OF PLANT GENETIC RESOURCES

(Registration No. 5/18336)

The Society was founded in 1987 with the following objectives:

- To serve and promote the scientific cause and to advance academic interest in the field of plant genetic resources.
- To disseminate knowledge relating to various aspects on plant genetic resources.
- To provide a forum for organizing symposia/conferences with a view to develop close relationship among the scientists engaged and interested in plant genetic resources activities.

Indian Journal of Plant Genetic Resources, the official publication of the Society, is published thrice in a year. The contribution to the journal, except for invited papers, is open to the members of the society only.

Membership to the society is open to all the individuals/institutions interested in various aspects of plant genetic resources. The membership fee is as follows:

Membership*	Inland	Foreign
Annual Member	Rs. 1,500.00	US\$ 120.00
Life Member	Rs. 6,000.00	US\$ 1,800.00
Institutional (Annual)	Rs. 30,000.00	US\$ 1,500.00
Institutional (5 years)	Rs. 2,00,000.00	US\$ 5,000.00
Institutional (10 years)	Rs. 4,00,000.00	US\$ 10,000.00

Issues of the journal published earlier are also available.

Limited space is available for advertisement of interest to botanists/geneticists/plant breeders and all those concerned with plant genetic resources.

CORRESPONDENCE relating to membership, advertisement and other related matters should be addressed to the General Secretary, Indian Society of Plant Genetic Resources, Office Block A, 2nd Floor, National Agricultural Science Complex, Dev Prakash Shastri Marg, Opposite Dasghara, Pusa Campus, New Delhi - 110012, India (E-mail: ispgr2015@gmail.com).

COMMUNICATIONS regarding the publication of research papers should be addressed to Editor-in-Chief, Indian Journal of Plant Genetic Resources, Indian Society of Plant Genetic Resources, Office Block A, 2nd Floor, National Agricultural Science Complex, Dev Prakash Shastri Marg, Opposite Dasghara, Pusa Campus, New Delhi - 110012, India. E-mail: ispgr2015@gmail.com

Print ISSN: 0971-8184
Online ISSN: 0976-1926


Indian Journal of Plant Genetic Resources

Vol. 39 No. 2 2026

Editor-in-Chief R K Tyagi

Editors (Foreign) Ehsan Dulloo, Michael Halewood, Shivali Sharma

Editors (Indian) V Celia Chalam, Sudip K Dutta, Sherry Rachel Jacob, Vikender Kaur,
Rajiv Kumar, Mohal Lal, K Pradheep, Aditya Pratap, S Rajkumar,
Somnath Roy, RK Salgotra, Rakesh Singh, Amit Siingh,
Sheikh M Sultan, Kuldeep Tripathi, Manjusha Verma



INDIAN SOCIETY OF PLANT GENETIC RESOURCES

Office Block A, 2nd Floor, National Agricultural Science Complex, Dev Prakash Shastri Marg,
Opposite Dasghara, Pusa Campus, New Delhi - 110012, India
Website: www.ispgr.nbpgri.ernet.in

INDIAN SOCIETY OF PLANT GENETIC RESOURCES

Patrons	MS Swaminathan, RS Paroda, Mangala Rai
Honorary Fellows	MS Swaminathan, GS Khush, S Rajaram, RS Paroda, Mangala Rai, Emile Frison, RB Singh, SK Datta, HY Mohan Ram, KS Gill, SK Vasal, Kamal Bawa, PL Gautam, BS Dhillon, Sanjay Kumar

EXECUTIVE COUNCIL (2025-27)

President	RS Paroda (New Delhi)
Vice Presidents	RC Agrawal (New Delhi); RK Tyagi (New Delhi)
General Secretary	Anuradha Agrawal (New Delhi)
Joint Secretary	Sherry Rachel Jacob (New Delhi)
Treasurer	Kuldeep Tripathi (New Delhi)
Ex Officio Member	DK Yadava, Director, ICAR-NBPGR, New Delhi
Councillors	North Zone: Pavan Kumar Malav, Sandeep Kumar (New Delhi) East Zone: Amit Kumar (Umiam), Somnath Roy (Hazaribag) West Zone: Kartar Singh (Ajmer) South Zone: N Sivaraj (Hyderabad), Rohini MR (Bengaluru) Central Zone: Vaidurya Pratap Sahi (Prayagraj), Vidya Sagar (Varanasi)

EDITORIAL BOARD

Editor-in-Chief

RK TYAGI

Indian Society of Plant Genetic Resources (ISPGR)
editor.pgr@gmail.com

Foreign Editors

Michael HALEWOOD

Head of Policy Unit
Bioversity International Rome, Italy
m.halewood@cgiar.org

Ehsan DULLOO

Principal Scientist
Team Leader, Integrated Conservation Strategies
Bioversity International Mauritius
e.dulloo@cgiar.org

Shivali Sharma

Senior Expert Specialist- Plant Genetic Resources
& Pre-breeding (Consultant)
Global Crop Diversity Trust (GCDT) & Adjunct Associate
Professor, Murdoch University, Australia
Shivali.sharma@murdoch.edu.au

Indian Editors

V Celia CHALAM

Principal Scientist
Division of Plant Quarantine
National Bureau of Plant Genetic Resources
Pusa Campus, New Delhi-110 012
celia.chalam@icar.org.in; mailcelia@gmail.com

Sherry Rachel JACOB

Senior Scientist
Division of Germplasm Conservation
ICAR-National Bureau of Plant Genetic Resources
Pusa Campus, New Delhi-110 012
Sherry.Jacob@icar.org.in; sherryrachel@yahoo.co.in

Rajiv KUMAR

Principal Scientist (Horticulture)
Division of Flower and Medicinal Crop ICAR-Indian
Institute of Horticultural Research Hessaraghatta lake
post, Bengaluru-560089, Karnataka
florirajiv@gmail.com

K PRADHEEP

Principal Scientist
ICAR-NBPGR Regional Station - Thrissur, Vellanikkara, KAU
P.O. Thrissur-680656, Kerala
k.pradheep@icar.org.in; hortpradheep@gmail.com

Sudip K DUTTA

Senior Scientist (Horticulture)
ICAR Research Complex for NEH Region Sikkim Centre
Tadong, Gangtok-737102, Sikkim
Sudip.Dutta@icar.org.in

Vikender KAUR

Senior Scientist
Division of Germplasm Evaluation
ICAR-National Bureau of Plant Genetic Resources
Pusa, New Delhi-110012, India
vikender.kaur@icar.org.in;
kaurvikender@gmail.com

Mohan LAL

Principal Scientist (Plant Breeding & Genetics)
Agrotechnology Technology & Rural Development Division,
CSIR-North-East Institute of Science & Technology
(CSIR-NEIST)
P.O.- RRL, Jorhat 785 006, Assam
mohan@neist.res.in; drmohanlal80@gmail.com

Aditya PRATAP

Principal Scientist
ICAR-Indian Institute of Pulses Research
Kanpur-208 024, Uttar Pradesh
adityapratapgarg@gmail.com

S RAJKUMAR

Senior Scientist

Division of Genomic Resources
ICAR-National Bureau of Plant Genetic Resources Pusa
Campus, New Delhi-110012
s.rajkumar@icar.org.in; rajnbpgr@gmail.com

RK SALGOTRA

Professor & Coordinator

School of Biotechnology

Sher-e-Kashmir University of Agricultural Sciences &
Technology of Jammu-180 009
rks_2959@rediffmail.com

Amit SINGH

Senior Scientist

Division of Genomic Resources

ICAR-National Bureau of Plant Genetic Resources
Pusa Campus, New Delhi-110012
Amit.Singh5@icar.org.in; amit_singh79@yahoo.com

Sheikh M SULTAN

Principal Scientist & Officer-In-Charge

ICAR-NBPGR, Regional Station, Srinagar

ICAR-CITH Campus, KD Farm, Old Airfield, Srinagar-190
005, Jammu & Kashmir mohmmad.sheikh@icar.org.in;
Sheikhmsultan@gmail.com

Somnath ROY

Scientist

Central Rainfed Upland Rice Research Station
ICAR-National Rice Research Institute
Hazaribag-825 301, Jharkhand
sroypbr@gmail.com

Rakesh SINGH

Principal Scientist & Head

Division of Genomic Resources

National Bureau of Plant Genetic Resources
Pusa Campus, New Delhi-110 012
rakesh.singh2@icar.org.in; singhnbpr@yahoo.com

Kuldeep TRIPATHI

Senior Scientist

Division of Germplasm Evaluation

ICAR-National Bureau of Plant Genetic Resources
Pusa Campus, New Delhi-110 012
kuldeep.tripathi@icar.org.in; kdttripathi89@gmail.com

Manjusha VERMA

Principal Scientist

Division of Genomic Resources

ICAR-National Bureau of Plant Genetic Resources
Pusa Campus, New Delhi-110 012
Manjusha.Verma@icar.org.in; manjusha_v@yahoo.com

CONTENTS

Review Article

- 1 Sustainability Benefits Following the Adoption of Genetically Modified Crops in Canada** **1**
Stuart J Smyth

Research Articles

- 2 Deciphering Genetic Variability and Stability for Yield Improvement in Mungbean (*Vigna radiata*)** **9**
Chandra Mohan Singh, Mukul Kumar, Hitesh Kumar, Priyanka Kaundal, Chirag, Pradyum Maurya, Ruchi, Anurag and Aditya Pratap
- 3 Aroma and Agro-morphological Diversity in Traditional Rice (*Oryza sativa*) Germplasm from Eastern India** **20**
Pooja Belwal, Amrita Banerjee, Anima Mahato, Priyamedha, Sougata Bhattacharya, Nabaneeta Basak, Nimai Prasad Mandal and Somnath Roy
- 4 Assessment of Molecular Diversity and Phylogenetic Analysis Among Chilli Genotypes Using Start Codon Targeted (SCoT) Markers** **29**
Hanuman Ram, DK Samadia, Chet Ram and AK Verma
- 5 Assessment of Antioxidant Activity and Quality Attributes in Indigenous Pran (Kashmiri Shallot) Genotypes** **38**
Gazala Nazir, Suraya Ahsan, Mudasir Magray, Khursheed Hussain, Saima Tabasum, UH Masoodi, Syed Berges, Asima Amin, Abida Jabeen and AA Lone
- 6 Morpho-biochemical Characterization and Multivariate Assessment of Genetic Diversity in Rajmah (*Phaseolus vulgaris* L.) Genotypes from Northeast India** **43**
Bedashruti Saikia, Rupam Borgohain Gayatree Goswami, Jutika Das and Anubrat Borah
- 7 Fatty Acid Composition of Seeds of Three Native *Garcinia* Species from Andaman and Nicobar Islands, India** **53**
Pooja Bohra and Ajit Arun Waman
- 8 Evaluation of Barley (*Hordeum vulgare* L.) Germplasm for Heat Tolerance Using Germination Indices and Seedling Traits** **58**
Mohmad Amin Bhagat, Navjyot Kaur, Prinka Goyal and Simarjit Kaur
- 9 Effect of Various Seed Priming Agents on Germination, Seedling Vigour and Grain Yield of Dry Direct-Seeded Rice in the Northwestern Indo Gangetic Plains of India** **67**
Sukham Madaan, Mangat Ram, Lalita, Sumit Saini, Ashish Jain, Rakesh Kumar, Mahaveer Bochalya, Amit Kumar, Ashwani Kumar, Vishal Gandhi, Charan Singh, Parvesh Kumar, Sarita Devi and Anita Kumari
- 10 Morphological Diversity Analysis in Brinjal (*Solanum melongena*) Germplasm** **74**
TH Singh, Sushil Pandey, Chitra D Pandey and Suresh R Yerasu
- 11 Preliminary Characterization of Wild *Actinidia Callosa* Var. *Strigillosa* as Potential Genetic Resources from North Sikkim, India** **83**
Abhilash Padhan, Dinesh Singh Thakur, Vishal Singh Rana, RK Dogra, Narender Negi, Akriti Chauhan and Pooja Sharma
- 12 Present status of *Pubistylus andamanensis* Thoth. in Andaman Islands and its IUCN Red List Assessment** **100**
CS Purohit, Vivek CP, BC Dey and LJ Singh
- 13 Correlation and Path Analysis for Seed Yield and Components Traits in Mothbean [*Vigna aconitifolia* (Jacq) Marechal] under Arid Environment** **105**
Savita Meena, NK Sharma, Pinky Yadav, Rukoo Chawla and Ajit Kumar Meena
- 14 Genetic Variation and Elite Clone Selection in *Gmelina arborea* for Seed Source Development in Northeast India** **114**
Manish Kumar Singh
- 15 Exploring Maize (*Zea mays* L.) Landraces: Characterization for Yield and Quality Traits to Foster Genetic Improvement** **125**
Heena Sharma, Surinder Sandhu, Gagandeep Kaur and RK Bhavyasree

Book Review

- 16 A Review of Code Paroda: A Life in Service of Agricultural Science** **135**
A Amarender Reddy
- 17 Plant Germplasm Registration Notice** **143**
- 18 Authors Guidelines** **169**

REVIEW ARTICLE

Sustainability Benefits Following the Adoption of Genetically Modified Crops in Canada

Stuart J Smyth^{1*}

Abstract

The adoption of genetically modified canola in 1995 in Western Canada was done to improve the control of weeds during crop production. Thirty years later, Western Canadian agriculture has witnessed a complete transformation, whereby the use of summerfallow as the leading form of weed control has virtually disappeared, to be replaced by continuous cropping, zero or minimum tillage land management practices. Of course there are trade-offs. The removal of tillage practices has increased soil moisture conservation and reduced soil erosion, but it has also resulted in an increase in herbicide applications. However, the environmental impact from the current suite of herbicides applied are considerably lower than previous ones, which often required soil incorporation instead of foliar application. All food production practices have impacts on the environment and biodiversity, however, the key is to identify best practices that contribute to increased production coupled with a lower environmental footprint. This article reviews the transition that Western Canadian agriculture has undergone over the past 30 years, highlighting the benefits for the environment and biodiversity.

Keywords: Biodiversity, Genetically modified crops, Innovation, Regulation, Tillage

¹Department of Agricultural and Resource Economics, University of Saskatchewan, 51 Campus Drive Saskatoon, Saskatchewan, S7N 5A8, Canada.

***Author for correspondence:** stuart.smyth@usask.ca

Received: 12/02/2026

Revised: 23/05/2026

Accepted: 26/05/2026

How to cite this article: Smyth SJ (2026) Sustainability Benefits Following the Adoption of Genetically Modified Crops in Canada. *Indian J. Plant Genet. Resour.* 39(2):1-8.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.176026>

Introduction

Effective weed control is a leading factor in crop yield maximization. Inefficient weed control enables weeds to use scarce soil moisture and nutrients, rather than these being available for the seeded crop. The high rate of seed production by weeds results in a compounding of the problem from one year to the next, if improperly controlled. For example, striga is an extremely noxious weed in Africa and if not properly controlled can result in yield losses of 100% (Benjamin *et al.*, 2024). The lost yield has significant economic impacts, with one estimate indicating that parasitic weeds in Africa account for economic losses of US\$200 million per year, increasing at a rate of US\$30 million per year (Rodenburg *et al.*, 2016). Yield losses from ineffective weed control exist across all crop kinds, as research from India found yield losses of up to 76% for soybeans, 71% for groundnut, 66% for rice, and 65% for corn (Gharde *et al.*, 2018).

The inability to efficiently control weeds in dryland Canadian Prairie agriculture can reduce wheat yields by up to 20% (Flessner *et al.*, 2021) and canola yields by up to 40% (Harker, 2001). Greater yield variability resulting from higher weed populations can reduce yields to the point that the crop is no longer profitable. Weeds propagate at far higher rates than domesticated crops, as a high-yielding wheat variety is capable

of producing 40-50 seeds per plant, while palmer amaranth, for example, can produce at least 100,000 seeds per plant (Grain Farmers of Ontario 2025). Kochia is a noxious weed commonly found in the Prairies, which is capable of producing up to 25,000 seeds per plant per year (Government of Manitoba, no date). Given the dramatically higher seed production, inefficient weed control not only has adverse economic impacts during the growing season, but also spills over and further impacts subsequent crops. Early germinating weeds are capable of growing rapidly and providing a canopy that shields seeded crops from sunlight, resulting in minimal or even no crop production within weedy areas of crop land.

Tillage has historically been used to control weeds in virtually all crop-producing regions. Adverse impacts of tillage use are increased soil erosion and reduced moisture conservation. The decade of the 1930s witnessed millions of tons of topsoil erosion due to the prolonged drought and consistent tillage practices across the major crop-producing regions of both Canada and the United States. The 1980s were a similar decade of below average precipitation, resulting in further soil erosion in these regions. During this decade, farmers and industry increasingly realized practices needed to change to ensure the future sustainability of crop production. Equipment innovations enabled greater adoption of direct seeding practices, reducing the need for tillage as farmers began to transition tillage out of their land management practices.

A parallel innovation that complemented the removal of tillage from land management practices was the commercialization and adoption of genetically modified (GM) and mutagenic herbicide-tolerant (HT) canola varieties, beginning in the mid-1990s. In-crop herbicides before this time provided limited weed control, especially for broad-leaf weeds and thistles. The initial GM canola varieties that were glyphosate and glufosinate-tolerant provided consistent year-over-year effective weed control that farmers required to completely remove tillage as the leading form of weed control. As a result, the adoption of GMHT canola became the most rapidly adopted innovation in the history of Canadian Prairie agricultural production.

This article provides a review of crop production practices in the Canadian Prairies before the commercialization of HT crops in 1995 and in the subsequent 30 years. It highlights how improving the efficiency of weed control has driven the improved sustainability of agriculture across the crop-producing regions of Western Canada.

Crop Production Practices Prior to GM Crop Adoption

Following the settlement of the three Prairie provinces of Alberta, Saskatchewan, and Manitoba in the latter half of the 19th century, weed control quickly became of key importance. This issue quickly arose as many of the weeds that had affected crop production in Europe were transported to this region via seed shipments that settlers brought with them. The result was that within several years of land being broken for crop production, weeds began to establish and negatively affect yields. Weed control was achieved through the use of summerfallow, which is a process whereby a field is left fallow for an entire growing season and routinely tilled to prevent weed growth. The number of required tillage passes was dependent on the amount of rainfall in the region, as in some years, only 2-3 tillage passes would be required, while in others, as many as 10 may have been needed. Fig. 1 illustrates the total summerfallow area for the three Prairie provinces, which peaked in 1970 at 14.9 million hectares (36.8 million acres) and has consistently declined to 600,000 ha in 2023 (1.5 million acres).




Fig. 1: Canadian Prairie summer fallow acreage, 1913-2022.

Source: Statistics Canada, 2023.

Given the importance of summerfallow for efficient weed control in the 20th century, many parts of the Canadian Prairies had two-year crop rotations that consisted of wheat one year and summerfallow the next. In areas with greater livestock densities, three-year rotations were common, where wheat would be grown for cash sales on summerfallow, followed by a crop used for forage purposes, such as oats, barley, or clover. The third year would be summerfallow. Crop diversity was limited, as there were few legume varieties developed for Prairie production, other than forage varieties as part of livestock nutritional needs. Many farms only produced wheat, as it was the most highly priced cereal

crop, and if a farm was strictly involved in grain production, other cereal crops had lower economic value. With the introduction of rapeseed varieties in the 1950s and the initial pea varieties in the 1970s, crop diversity slowly began to increase.

Near the end of the 20th century, a lengthy period of drought impacted this region, reducing yields for most of the 1980s. During this drought period, it became increasingly apparent for the need to change land management practices as topsoil was frequently being blown away from summerfallow fields. During this period, on windy days, the sky would be filled with dirt and ditches were often filled with blown dirt.

Innovative farmers and short-line equipment manufacturers began to experiment with new seeding technologies, such as those that no longer turned the soil over, such as with diskers, but rather used a chisel to create a furrow for seed placement, leaving the soil untilled. While some experimenting initiated in the late 1960s, this expanded in the 1970s and by the 1980s, manufacturing of direct seeding implements was occurring. The increased adoption of zero tillage and minimum tillage equipment throughout this period were contributing factor to the reduction in summerfallow acreage in Fig. 1.

There have been some periodic increases in the use of tillage, especially in years with higher snowfall and spring precipitation, as farmers will use tillage as part of the efforts to dry soils down to the point that they can be seeded. Additionally, the province of Manitoba typically receives higher precipitation than the other two provinces, which contributes to explaining the higher rate of summerfallow variability, as it is not uncommon that land is too wet to seed in the spring, forcing farmers to summerfallow unseeded fields to control weeds.

The high utilization of tillage reduced the potential for improved biodiversity, given that fields were routinely tilled to ensure weeds were not growing, thereby depriving many populations of sufficient feed sources, not to mention protection from ground cover. The consistent tillage passes over summerfallow fields additionally contributed to reduced biodiversity in crop fields, especially during summerfallow years.

Coming out of the 1980s, Prairie agriculture had been adversely affected by the decade-long drought. Drought, combined with high winds, once again created significant soil erosion problems. Many farmers realized that changes would be required to ensure the long-term sustainability of their land. This realization improved farmers' willingness to approach innovative technologies with a more positive approach than may previously have been the case, as without new products and technology, things would not improve. Land

management changes were occurring across the Prairies during the 1980s, which would only increase in the subsequent decades.

Crop Production Practices Following GM Crop Adoption

The final handful of years of the 20th century witnessed one of the most significant innovations in the history of crop agriculture, the commercialization of genetically modified herbicide-tolerant (GMHT) crop, namely canola, corn, cotton and soy, although in the 1990s, only GMHT canola was grown in the Prairies. The potential to efficiently control weeds in crops now existed, resulting in a transition in land management practices as the use of summerfallow to control weeds consistently declined following the commercialization of GMHT canola in 1995. Over the ensuing decade, summerfallow acreages decreased as crop production became a continuous process, frequently involving the use of zero tillage or minimum tillage land management systems.

As identified above, efficient weed control is a key factor towards ensuring that yields are as high as can be possible. Given the lack of efficient in-crop herbicides to control weeds, the historically leading method for efficient weed control was through the practice of summerfallow. The commercialization of GMHT crops, but GM canola in particular, in 1995, provided a new option for efficient weed control. The first two GM canola varieties that were commercialized were Monsanto's (now Bayer) glyphosate-tolerant canola and Agrevo's (now Bayer) glufosinate-tolerant canola. Pioneer Hi-Bred also had a variety of imidazolinone-tolerant canola commercialized in 1995. Following the risk assessment of HT canola varieties by the Canadian Food Inspection Agency (CFIA), these varieties entered a two-year seed multiplication program, consisting of 30,000 acres in 1995 and 230,000 acres in 1996 (Smyth and Phillips, 2001).

Initial farm level adoption occurred in 1997, with virtually full adoption within 15 years (Fig. 2). This is considerably faster than the average length of time for peak adoption, which is estimated to be 22-23 years in length (Alston and Pardey, 2021). By the third planting season in 1999, over 50% of seeded acres were done with HT varieties. By 2004, adoption reached 75% and by 2009 was estimated to be 91%. The adoption rate of HT canola varieties represents the most rapid adoption of any agricultural technology in the history of Canadian agriculture. This rapid rate of adoption indicates that farmers experienced efficient and economic weed control by adopting HT canola varieties. There is virtually no non-HT canola currently produced.




Fig. 2: Canadian HT canola adoption rates, 1997-2013 (seeded acres percentage).

Source: Canola Council of Canada (no date).

With the 2026 crop year representing the 30th year of commercial HT canola production in the Canadian Prairies, significant additional benefits have been realized. With the rapid adoption of HT canola in the decade from 1997-2007, this corresponds to the significant decline in crop summerfallow acres in Fig. 1. In Saskatchewan alone, summerfallow dropped from 4 million hectares to less than 400,000, a decrease of over 90%. The effective weed control provided by HT crops enabled farmers to remove summerfallow from their crop rotations. This has resulted in many fields across the Canadian Prairies being in zero tillage or minimum tillage practices for 20 years or even longer with some of the earliest adopters. The contrast in the use of tillage between crop-producing regions that adopted HT crop technology compared to those that failed to do so is starkly evident in Fig. 3. Europe rejected GM crop technologies, with only Spain and Portugal producing small acre ages of GM corn. The remainder of Europe's crop-producing regions rely on older, more environmentally impactful pesticides as well as tillage for weed control options.

The results of a farmer survey following a decade of HT crop production identified substantial environmental benefits, with 86% of respondents indicating they had experienced reduced soil erosion and 83% reporting higher levels of soil moisture conservation (Smyth *et al.*, 2011a). Further benefits were identified through the reduction in the environmental




Fig. 3: Comparison of tillage rates between Canada and Europe.

Source: EuroStat (2016); Sutherland *et al.*, (2021).

impacts of canola in-crop herbicides applied. Before the commercialization of HT canola varieties, there were very limited options for in-crop herbicide controls, resulting in many herbicides having to be soil incorporated before, or in conjunction with, seeding. The transition to effective in-crop foliar-applied herbicides reduced the environmental impact of herbicides, with the environmental impact of the top five herbicides applied to canola production dropping by 53% between 1995 and 2006 (Smyth *et al.*, 2011b). A key reason for the decline in the environmental impact of canola herbicide use was the drop in the volume of herbicide active ingredient needing to be applied, which dropped from 648 grams per hectare to 401 grams per hectare, a 38% reduction. Between 1995 and 2006, this represented a reduction in the volume of herbicide active ingredient being applied to canola fields by an estimated 1.5 million kilograms.

The increase in canola acres is represented by the reduction in summerfallow acres. In 1995, the three Prairie provinces of Alberta, Saskatchewan and Manitoba had combined summerfallow acres of 13 million (Fig. 1). Between 1995 and 2025, seeded canola acres increased by 8.5 million acres (Fig. 4). The increased canola production did not displace other crop production; it replaced annual summerfallow acres.




Fig. 4: Canadian Prairie seeded canola acres, 1986-2025 (millions of acres).

Source: Canola Council of Canada 2025.

A complement to the adoption of HT canola varieties was increased investment into the development of new pea and lentil varieties across the Prairies. This allowed farmers to increase the diversity of their crop rotations, which expanded to include alterations between cereals, pulses, and oilseeds. From 2000-2008, lentil production ranged from 1.3 to 2 million acres. This rose significantly over the subsequent decade, such that from 2020 to 2024, production increased to a range from 3.7 to 4.3 million acres (Sask Pulse, 2025). This increased crop type rotation diversity reflects a significant change from earlier decades, where rotations consisted of wheat and summerfallow.

Table 1: Change in average annual Saskatchewan fertilizer use, 1991-1994 to 2016-2019

Period	Crop yield (bu/ac)	Fertilizer applied (lbs/ac)	Fertilizer rate (lbs/bu)	Crop acres (millions)
1991-94	40	231	6	33.3
2016-19	51	467	9	40.5
Percent increase	28%	102%	50%	22%

Rotation diversity ensures that different herbicide modes of action are applied to fields, thereby contributing to ensuring that the development of herbicide resistance in weed populations is minimized, as this follows good agronomic practices.

The diversity in the number of crops in a rotation has impacted fertilizer use, with greater volumes being applied. This increase in applications has contributed to a corresponding yield increase. While fertilizer volumes have increased, the correct assessment is to determine how much crop is produced per pound of fertilizer applied, resulting in a fertilizer use efficiency calculation. Between the periods of 1991-1994 and 2016-2019, Saskatchewan farmers increased their average annual total fertilizer application by 102% (Table 1). This suggests that for every 1% increase in fertilizer application, there is a corresponding 0.87% increase in crop yields. From 1991-1994 to 2016-2019, Saskatchewan farmers converted 7.2 million acres of summerfallow to continuous crop production, thereby increasing the overall fertilizer use, as summerfallow production does not require the use of fertilizers. This conversion accounts for 44% of the increase in fertilizer usage. Additionally, there was a shift in fertilizer application timing, particularly due to advances in equipment and technology that allow for in-crop fertilizer applications. In the 1991-1994 period, no in-crop fertilizer applications were made, while during the 2016-2019 period, 13% of total fertilizer applications were made in-crop (Lika *et al.*, 2024a). This advancement resulted in less post-harvest fertilizer application, which can be susceptible to spring melting and run-off, whereby nutrients are unintentionally transferred to watersheds.

The increased crop rotation diversity and the resulting alternating herbicide modes of action have made a significant contribution to lowering the development of herbicide resistance (HR) in new weeds. In the period from 1988-1995, an average of 2.5 new HR weeds were reported each year. The adoption of HT crops has contributed to a reduction in the reporting of new HR weeds, as the rate of reporting across the Canadian Prairies has continually dropped. As farmers learned how to best integrate HT crops into their rotations, the rate of new HR weed reporting dropped

precipitously. Between 2016 and 2021, only three new HR weeds were reported in all of the Canadian Prairies (Sutherland *et al.*, 2025). This drop in the development of new HR weeds has been driven by increased crop rotation diversity as the number of farmers practising at least three crops in their rotation increased from 59% from 1991-1995 to 80% from 2016-2019, while the percentage of farmers practicing two-year rotations decreased from 39 to 2% over the same period (Sutherland *et al.*, 2025).

The transition in land management practices away from rotations that relied on tillage as the leading form of weed control to continuous cropping involving HT crops in the rotation has transitioned crop land from being a carbon emitter to a sequesterer. Summerfallow practices required frequent passes during the growing season to ensure efficient weed control, with the total number of passes varying by region and precipitation levels. Each tillage pass released carbon into the atmosphere, resulting in higher amounts of carbon released during the summerfallow year than was sequestered during the crop year. It is estimated that the reliance on tillage by farmers in Saskatchewan from 1991 to 1994 resulted in a 1,000-hectare farm releasing 33.4 mg of soil organic carbon (SOC) annually, while storing 16.7 Mg of SOC from crop production. By the 2016 to 2019 period, the same 1,000-hectare farm would have sequestered 119 Mg of SOC annually, in addition to a further reduction of 421 Mg of SOC due to the reduction in summerfallow practices (Sutherland *et al.*, 2021). If the change in emissions was compared to the average emission of an automobile, a 1,000-hectare farm from 1991-95 would have released 15 times more carbon than an average automobile per year, and by the 2016-2019 period, it would have sequestered the equivalent emissions from 95 cars.

With crop land being in minimum or zero tillage land management practices for 20 years or even longer, farmers are experiencing increases in soil health. While many farmer observations are challenging to quantify, such as reduced soil erosion following substantial rainfall, it is possible to quantify changes in soil organic matter. Soil testing is a common practice in Prairie crop land, as it measures the level of soil nutrients available in a specific field each fall, providing farmers with greater

knowledge about crop nutrient requirements for the subsequent crop year. A standard part of each soil test is to report the level of soil organic matter. Research on changes in soil organic matter levels in Saskatchewan identified that soil organic matter levels increased 41% between the periods of 1990-1999 and 2015-2024 (Sutherland *et al.*, forthcoming). Increases in soil organic matter were reported across all of the major soil zone classifications in the province.

The above confirmation of changes in land management and weed management practices dating back to the early 1990s confirms the substantial impacts of agricultural innovations. The combination of improved crop genetics, such as HT crops and nitrogen-fixing pulse crops, the adoption of more benign crop pesticides—especially herbicides, and equipment advances enabling minimum and zero tillage land management practices all combine to create a system of sustainable crop production. While those critical of modern crop production practices suggest that continuous cropping and the resulting increase in pesticide applications are a detriment, the evidence from 30 years of HT crop production as part of a crop rotation soundly rejects this argument.

Benefits for Biodiversity

One of the leading debates regarding the transition away from utilizing tillage as the leading form of weed control to diverse, but constant, crop production, which relies on pesticides to control production threats, such as plant diseases, insects, and weeds. The total volume of pesticide active ingredient applied has increased due to the millions of additional crop acres that are presently part of crop rotations, as compared to crop rotations that relied on summerfallow for weed control. Even though more crop acres are planted each year, and more pesticides are applied, the environmental impact of those being applied is frequently lower. Comparing the in-crop herbicides applied to Saskatchewan crops between the periods of 1991-1994 and 2016-2019 identified that the environmental impact of those applied was 65% lower, while 45% less herbicide active ingredient was applied per acre (Lika *et al.*, 2024b). The decline in the environmental impacts of the herbicides being applied contributes to mitigating impacts on non-target organisms.

One of the most significant changes for biodiversity has been the transition to continuous plant matter in fields. When rotations were predominantly wheat followed by summerfallow, the field was barren soil every other year, limiting biodiversity as there was no ground or crop cover for many species. Summerfallow

fields consisted of black dirt, which was highly susceptible to wind and water erosion. Constant crop cover provides increased habitats for beneficial insects, including pollinators, as well as ground-nesting birds.

Assessing pollinator populations, especially wild populations, is extremely challenging, such that few studies exist. One metric that is used to assess pollinator populations is the number of honey bee hives. Even though these hives are managed, they are subject to the full array of population stressors, such as diseases, mites, land use change, and pesticides. While total crop acres have increased over the past 35 years, resulting in larger volumes of pesticide active ingredient being applied, the environmental impacts of the increased applications are lower, resulting in increased honey bee hives across Canada (Fig. 5). The number of honey bee hives has increased by 14% over the 12 years from 2012-2024.




Fig. 5: Honey bee colonies in Canada, 2012-2024.

Source: Government of Canada 2025.

There are additional uncorroborated reports from farmers at public events that they believe there has been an improvement in biodiversity following the removal of tillage as the leading form of weed control. As the increase in soil organic matter has been quantified, farmers indicate there has also been a corresponding reduction in soil compaction due to increases in earthworm populations. While this has not been confirmed through peer-reviewed analysis, it is a form of citizen science engagement, whereby farmers provide valued insights into field-level changes occurring on their farms over time.

Agriculture and the production of food will always have impacts on biodiversity. The only way for agriculture and food production not to have an impact on biodiversity would be to stop food production entirely. This is not a realistic option, so the challenge becomes how to reduce the biodiversity impacts of agriculture and food production as much as is reasonably possible, without adversely impacting the overall amount of food produced. Research indicates that the adoption of GM crops contributes to reducing the impacts of crop agriculture on biodiversity (Carpenter, 2011). Farmers across the Prairies have

substantially adopted HT canola, the majority of which is GM; however, all HT canola production provides biodiversity benefits. The ability of farmers to adopt production practices that include continuous cropping and zero tillage or minimum tillage ensures that the environmental impacts of these practices have been substantially reduced, thereby providing biodiversity benefits.

Summary

Over the past 30 years, agriculture in Canada has consistently demonstrated advances in sustainable production practices, as well as in mitigating impacts on biodiversity. This achievement has been possible due to significant and consistent investments into research and development (R&D) of new products and technologies that have driven these advances. Increases in sustainable agricultural practices and reduced biodiversity impacts are only possible from sustained R&D investments; they are not possible by returning to historic crop production, as these varieties are more susceptible to insect, weed, and disease pest pressures, which results in lower yields, as well as resulting in greater environmental and biodiversity impacts due to the application of older, more toxic pesticides.

Crop production in the Canadian Prairies has aptly demonstrated the value of investing in new agricultural products and technologies. Research into the initial application of genetically modified technology began in the mid to late 1980s. A decade of this research resulted in the initial commercialization of GM crops in the mid-1990s. This region is now globally recognized as among the most sustainable producers of crops (Bamber *et al.*, 2025). As the sustainability of crop production increases, the biodiversity benefits are correspondingly enhanced through innovations such as more efficient fertilizer applications, application of pesticides with lower environmental impacts, and the removal of summerfallow practices from land management systems. However, this would not have been achieved were it not for the 40 years of innovation investments and 30 years of commercial production.

A significant challenge for agriculture, especially regarding contributing to meeting international climate change commitments, is how to contribute to meeting these objectives in the face of increased pressure from environmental organizations focused on preventing the integration of innovative strategies into domestic policy and funding frameworks. The cornerstone of international sustainability principles is to produce more food per acre of land, while reducing the level of inputs required, as well as reducing the land area for food

production. This triple challenge will be impossible to meet without dramatic advancements in innovative products and technologies.

As has been empirically established in the Canadian Prairie regions, the adoption of GM crops, especially those that are herbicide-tolerant, is the driving force of sustainability advances and a mitigation of the factors that can be stressors to biodiversity. This has been confirmed with 91% of Saskatchewan farmers indicating that the use of glyphosate is the number one factor for their ability to increase the utilization of sustainable land management practices in their crop rotations (Sutherland *et al.*, 2021). The evidence clearly supports the tremendous advantages of the removal of tillage as a land management practice and how this is correlated to crop-producing regions that have a lengthy history of HT crop adoption and production. Improving soil health is an important factor for all farmers, and 30 years of HT crop production in the Canadian Prairies has demonstrated increased levels of soil organic matter, a key component of healthy soil definitions. This improvement would not have been possible were it not for the efficient weed control provided by HT crops, enabling the permanent removal of tillage practices.

As international climate agreements and bioeconomy frameworks become further integrated into domestic policies for agriculture and environment, evidence is going to be required to provide the basis for policy. While many activist environmental groups have invested massive fiscal resources in the creation and dissemination of false information regarding the safety and benefits of HT crops and biotechnology (Ryan *et al.*, 2020), the evidence from adopting farms and countries overwhelmingly refutes these activist claims. While there is a small but increasing component of research confirming the benefits of GM crops towards mitigating the effects of climate change, more research is required (Smyth *et al.*, 2024). The quantification of increased sustainable agriculture following the adoption of biotechnology and HT crops is in its infancy, and research into the impacts on biodiversity is very scant. Research in these fields is important to inform policymakers in non-GM crop adopting countries that the adoption of these sustainable crops provides a wide range of benefits in terms of increased crop and food production, as well as in reducing agriculture's environmental footprint. It is research of this type that is more important than ever, as if countries are going to be able to make domestic sustainability improvements, which can potentially reduce the impacts of agriculture and food production on biodiversity, greater adoption rates of GM crop and gene editing technologies will need to happen.

References

- Alston JM and PG Pardey (2021) The economics of agricultural innovation. In: CB Barrett and DR Just (eds.) *Handbook of Agricultural Economics*. Elsevier, Amsterdam, pp 3895-3980.
- Bamber N, I Turner and N Pelletier (2025) Rapeseed, wheat and peas grown in Canada have considerably lower carbon footprints than those from major international competitors. *Nature Food* 6: 757-761. <https://doi.org/10.1038/s43016-025-01212-0>
- Benjamin J, I Oluwadamilola, OK Babalola, EV Oziegbe, DO Oyedokun, AM Akinyemi and A Adebayo (2024) Cereal production in Africa: the threat of certain pests and weeds in a changing climate; a review. *Agric. Food Sec.* 13(18). <https://doi.org/10.1186/s40066-024-00470-8>
- Canola Council of Canada (2025) Canadian canola production statistics. Available at: <https://www.canolacouncil.org/markets-stats/production/>
- Canola Council of Canada (No date) Estimated acreage, production and percentage of herbicide-tolerant canola in Canada. Available at: <https://www.canolacouncil.org/markets-stats/production/>
- Carpenter J (2011) Impact of GM crops on biodiversity. *GM Crops and Food* 2(1): 7-23. doi:10.4161/gmcr.2.1.15086
- EuroStat (2016) Share of tillage practices in the EU27 2016. Available at: https://ec.europa.eu/eurostat/statistics-explained/index.php?title=File:Figure1_Share_of_tillage_practices_in_the_EU27_2016.png
- Flessner ML, IC Burke, JA Dille, WJ Everman, MJ VanGessel, B Tidemann, MR Manuchehri, N Soltani and PH Sikkema (2021) Potential wheat yield loss due to weeds in the United States and Canada. *Weed Technol.* 35(6): 916-23. <https://doi.org/10.1017/wet.2021.78>
- Gharde Y, PK Singh, RP Dubey and PK Gupta (2018) Assessment of yield and economic losses in agriculture due to weeds in India. *Crop Protect.* 107: 12-18. <https://doi.org/10.1016/j.cropro.2018.01.007>
- Government of Canada (2025) Statistical overview of the Canadian honey and bee industry, 2024. Available at: <https://agriculture.canada.ca/en/sector/horticulture/reports/statistical-overview-canadian-honey-and-bee-industry-2024>
- Government of Manitoba (no date) Managing kochia. Available at: <https://www.gov.mb.ca/agriculture/crops/weeds/print,managing-kochia.html#:~:text=Kochia%20seed%20production%20ranges%20from,way%20of%20spreading%20its%20seeds> (accessed 18 December 2025).
- Grain Farmers of Ontario (2025) Fact sheet: Palmer amaranth. Available at: https://gfo.ca/wp-content/uploads/2025/10/Fact_Sheet_Alert-Palmer.pdf
- Harker N (2001) Survey of yield losses due to weeds in central Alberta. *Can. J. of Plant Sci.* 81(2): 339-42. <https://cdnsiencepub.com/doi/pdf/10.4141/P00-102>
- Lika E, C Sutherland, S Gleim and SJ Smyth (2024a) Analyzing shifts in fertilizer use with crop rotation practices in Saskatchewan across distinct periods from 1991-94 to 2016-19. *Front. Sustain. Food Syst.* 8(1412443). doi: 10.3389/fsufs.2024.1412443
- Lika E, C Sutherland, S Gleim and SJ Smyth (2024b) Quantifying changes in the environmental impact of in-crop herbicide use in Saskatchewan, Canada. *Weed Tech.* 38(e28): 1-8. doi: 10.1017/wet.2024.15
- Rodenburg J, M Demont, SJ Zwart and L Bastiaans (2016) Parasitic weed incidence and related economic losses in rice in Africa. *Agric. Ecosyst. Environ.* 235(1): 306-317. <https://doi.org/10.1016/j.agee.2016.10.020>
- Ryan CD, AJ Schaul, R Butner and JT Swarthout (2020) Monetizing disinformation in the attention economy; The case of genetically modified organisms (GMOs). *Euro. Man. Jour.* 38(1): 7-18. <https://doi.org/10.1016/j.emj.2019.11.002>
- Saskatchewan Pulse (2025) 2024 Seeded Pulse Acreage. Available at: <https://saskpulse.com/resources/2024-seeded-pulse-acreage/>
- Smyth SJ, PWB Phillips and D Castle (2024) An assessment of the linkages between GM crop biotechnology and climate change mitigation. *GM Crops and Food* 15(1): 150-169. <https://doi.org/10.1080/21645698.2024.2335701>
- Smyth SJ, M Gusta, K Belcher, PWB Phillips and D Castle (2011a) Environmental impacts from herbicide tolerant canola production in Western Canada. *Agric. Syst.* 104(5): 403-410.
- Smyth SJ, M Gusta, K Belcher, PWB Phillips and D Castle (2011b) Changes in herbicide use following the adoption of HR canola in Western Canada. *Weed Tech.* 25(3): 492-500. <https://doi.org/10.1016/j.agry.2011.01.004>
- Smyth SJ and PWB Phillips (2001) Competitors co-operating: Establishing a supply chain to manage genetically modified canola. *Int. Food Agribus. Manage. Rev.* 4(1): 51-66.
- Statistics Canada (2023) Table 32-10-0359-01 - Estimated areas, yield, production, average farm price and total farm value of principal field crops, in metric and imperial units. <https://doi.org/10.25318/3210035901-eng>
- Sutherland C, S Gleim and SJ Smyth (Forthcoming) Impacts of reduced tillage on soil health in Saskatchewan.
- Sutherland C, S Gliem, S Lubieniechi and SJ Smyth (2025) Rate of herbicide resistant weed development: A Canadian Prairie case study. *GM Crops and Food* 16(1): 252-262. <https://doi.org/10.1080/21645698.2025.2477231>
- Sutherland C, S Gleim and SJ Smyth (2021) Correlating genetically modified crops, glyphosate use and increased carbon sequestration. *Sustainability* 13: 11679. <https://doi.org/10.3390/su132111679>

RESEARCH ARTICLE

Deciphering Genetic Variability and Stability for Yield Improvement in Mungbean (*Vigna radiata*)

Chandra Mohan Singh^{1*}, Mukul Kumar¹, Hitesh Kumar¹, Priyanka Kaundal¹, Chirag¹, Pradyum Maurya¹, Ruchi¹, Anurag¹ and Aditya Pratap²

Abstract

Mungbean [*Vigna radiata* (L.) R. Wilczek] is an important short-duration and widely adaptable pulse crop known for its nutritional and soil-ameliorative properties. However, its productivity still lags due to photo- and thermo-period sensitivity, the seasonal specificity of the available cultivars, and susceptibility to a wide range of biotic and abiotic factors. Therefore, breeding for developing stable elite cultivars is urgently required to improve productivity. A total of 55 genotypes of mungbean, including released varieties and elite lines, were tested over a period of three years to identify the stable elite lines for high yield. A wide range of variability was recorded for seed yield per plant (SYP) and its key component traits. Based on correlation coefficient analysis, PL and NSP were found to be key components. The significant interaction of genotype $\times$ year indicated the crucial role of the environment on SYP. The correlation coefficient analysis revealed that pod length (PL) and number of seeds per pod (NSP) were key yield contributing traits. GGE biplot analysis demonstrated that five genotypes, i.e. G42 (BMS 19-3), G17 (Shikha), G49 (ML 134), G18 (TRAM-18), and G27 (BMS 18-4), were found most stable and high-yielding genotypes over the year. Further, the molecular characterisation of these selected genotypes with SSR and functional markers opens the scope for their effective utilization in the mungbean breeding programme.

Keywords: Adaptability, DNA fingerprinting, Functional markers, Greengram, G $\times$ E interaction, Promising genotypes, Stability, SSR

¹Department of Genetics and Plant Breeding, Banda University of Agriculture and Technology, Banda, 210001, India.

²Project coordination Unit, All Indian Coordinated Project on Kharif Pulses, ICAR- Indian Institute of Pulses Research, Kanpur, 208024, India.

Received: 11/02/2026

Revised: 06/06/2026

Accepted: 08/06/2026

How to cite this article: Singh CM, M Kumar, H Kumar, P Kaundal, Chirag, P Maurya, Ruchi, Anurag and A Pratap (2026) Deciphering Genetic Variability and Stability for Yield Improvement in Mungbean (*Vigna radiata*). *Indian J. Plant Genet. Resour.* 39(2): 9-19.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.176011>

Introduction

Mungbean [*Vigna radiata* (L.) R. Wilczek] is an important short-duration grain legume that is cultivated over a wide range of environments (Singh *et al.*, 2022). It is rich in protein (25-28%) and several micronutrients (Pratap *et al.*, 2022; Zafar *et al.*, 2023). It improves the soil health by fixing the free atmospheric nitrogen into the soil and also improves the carbon content of the soil, which leads to yield improvement of the subsequent crop (Diatla *et al.*, 2020; Arya *et al.*, 2024). However, the productivity of mungbean is quite low due to several factors such as non-availability of quality seeds of improved varieties, abiotic and biotic stresses, poor crop management practices, low harvest index and post-harvest losses (Nair *et al.*, 2019; Pratap *et al.*, 2019).

Yield is a complex trait that always depends on its component traits and is determined by its genotype, environment and their interactions (Kumar *et al.*, 2023a; Kumar *et al.*, 2023b). Hence, consideration of environmental effect is necessary as many factors overshadow the genetic potential of the genotype and lead to their low performance (Parihar *et al.*, 2022; Bomma *et al.*, 2024; Sadaiah *et al.*, 2026; Irfan *et al.*, 2026). The significant gap between the observed yield and the potential yield of any cultivar occurs due to its variable

performance in different environments and the major reason behind this is the lack of performance stability.

Genotype by environment interaction (GEI) provides opportunities to select genotypes that interact positively with a specific environment (specific adaptation) or record outstanding performance in a wider range of environments (general adaptation). Further, it helps the breeders for a robust evaluation and appropriate selection of genotypes for yield stability over the environments or adaptability in the target environment (Abou-Khater *et al.*, 2022; Rao *et al.*, 2023; Wodebo *et al.*, 2023; Shobeiri *et al.*, 2024; Jafarullakhan *et al.*, 2024; Mohammadi *et al.*, 2024; Ortiz *et al.*, 2024; Kona *et al.*, 2024; Torres-Ordoñez *et al.*, 2024; Ma *et al.*, 2024). The characterization of identified genotypes for breeding programmes also needs attention for their effective utilization in further breeding programmes. Molecular markers have evolved as a unique tool for assessing genetic variability under the influence of the environment. Nowadays, various marker systems are available (Kyu *et al.*, 2024; Kohli *et al.*, 2024; Sao *et al.*, 2025; Ghallab *et al.*, 2025). Previous workers also gave emphasis on various types of omics approaches for characterizing the *Vigna* species for crop improvement (Raizada *et al.*, 2019; Altaf *et al.*, 2026; Pookhamsak *et al.*, 2026). Chugh *et al.*, (2022) also focused on molecular tools for improving the oilseeds and legumes. Keeping the above in view, the present study was undertaken with 55 genotypes of mungbean to evaluate the influence of $G \times E$ interaction on seed yield and its component traits; and identify the high yielding stable mungbean genotype for cultivation.

Materials and Methods

Plant materials, field experimentations and data recording

The present experiment was conducted with 55 genotypes of mungbean, including released varieties, breeding lines and checks (Supplementary Table 1). The experiments were conducted at the Research Farm of Banda University of Agriculture and Technology, Banda (25.5°N, 80.3°E and 113 MSL), India. The experiments were laid out in randomized block design (RBD) with three replications during *kharif* 2020-22. Each entry was grown in four rows of 2m length with 45 cm $\times$ 10 cm plant geometry. All the cultural operations were carried out as per the standard package and practices recommended for raising a good crop in the region. Harvesting of the crop was done twice, first through pod picking when >75% pods were mature and final

harvesting was done after 10 days of the first picking. The observations were recorded on six important yield traits and their contributing traits, such as days to maturity (DM), number of pods per plant (NPP), pod length (PL), number of seeds per pod (NSP), seed index (SI) and seed yield per plant (SYP). All the traits except DM were recorded on five random plants and averaged in each entry in each replication, whereas DM was calculated on a plot basis.

Statistical analysis

The data analysis for two-way ANOVA, genetic parameters and correlation coefficient were done by using R-Statistical software version 4.0 (package "Metan"). The GGE biplot analysis for SYP was done by estimating each element of the matrix using GEA-R software.

DNA extraction and fingerprinting of selected promising genotypes

Six mungbean genotypes, including two elite lines such as G42 (BMS 19-3), G27 (BMS 18-4); three released varieties, i.e. G17 (Shikha), G49 (ML 134), G18 (TRAM-18), and one check variety (IPM 02-3), were subjected to DNA fingerprinting. The 200 mg tiny leaves of selected genotypes were frozen in liquid nitrogen and homogenized using tissuelyser-II (Qiagen). The homogenized samples were further subjected to DNA extraction using Qiagen's DNeasy[®] Plant Mini Kit as per the manufacturer's protocol. After extraction, the quality of extracted DNA was examined on 0.8% agarose gel and quantified using Nano Drop (Qiagen QIAxpert). Further, the set of 12 markers (10 functional and 2 SSR) were used for the fingerprinting of genotypes through polymerase chain reaction (Supplementary Table 2). The 20 μ L PCR reaction mix, 10 μ L of PCR master mix (Thermo Scientific), 1 μ L of each forward and reverse primer, and 6 μ L nuclease-free water were mixed with 2 μ L genomic DNA. The polymerase chain reaction (PCR) amplification was carried out in a thermo-cycler (Veriti, Applied Biosystems) with the programming: initial denaturation at 95°C for 3 min, followed by 35 cycles of denaturation at 95°C for 30 sec, annealing (52-58°C) for 30 sec, extension at 72°C for 30 sec, and the final extension for 7 min at 72°C. Thereafter, the amplified PCR product was resolved through electrophoresis in 3% (w/v) agarose gel containing 0.5 μ g/mL ethidium bromide. A 1kb DNA ladder (Thermo Scientific) was utilized in gel electrophoresis to determine the size of the amplicons. The gels were visualized under a gel documentation unit (Vilber E-box).

Results

Genetic variability assessment

The two-way ANOVA showed significant differences ($P < 0.05$) among the yield and yield components for all traits under study (Table 1). The effect of year was also found to be significant for all the traits, except PL. In contrast, the genotype $\times$ year interaction was non-significant for most of the traits, except for SYP.

The boxplots (Fig. 1) illustrate the distribution of six traits under study, i.e. DM, NPP, PL, NSP, SI and SYP, across

three years. Two traits, such as DM and PL, did not differ significantly across the years and were found to be relatively stable. In NPP, Year 1 showed the highest median value, Year 3 the lowest median value, while Year 2 was found intermediate. For NSP, Years 1 and 2 were statistically similar and superior to Year 3. The SI exhibited a significant decline from Year 1 and Year 2 to Year 3, showing that environmental conditions in Year 3 negatively affected this trait. The most pronounced difference was observed for SYP, where Year 1 recorded the highest yield, followed by Year 2 and then Year 3, with

Table 1: Pooled analysis of variance for different traits in mungbean during *kharif* 2020-22

	DF	DM	NPP	NSP	PL	SI	SYP
Genotype	54	280.07**	443.67**	4.531**	7.95**	3.40**	21.47**
Year	2	57.69**	569.71**	27.45**	0.16	12.61**	417.91**
Gen $\times$ Year	108	2.89	9.96	0.093	0.06	0.07	1.004**
Residuals	324	3.30	8.32	0.14	0.081	0.06	0.500
Mean	-	63.68	20.92	10.34	7.72	3.62	8.43
CV (%)	-	2.71	9.55	13.67	18.31	27.60	11.85

** $P < 0.01$ level of significance.




Fig. 1: Box plot analysis of 55 mungbean genotypes across the three years.

all years differing significantly from each other. Overall, the results indicate that environmental variation had little influence on DM and PL, whereas NPP, NSP, SI, and especially SYP were substantially affected by year-to-year changes, resulting in reduced performance under the conditions prevailing in Year 3.

Trait association analysis

Correlation analysis showed a strong, significant association between SYP and PL ($r = 0.74$, $P < 0.001$) (Fig. 2). NSP exhibited a positive and significant correlation with PL ($r = 0.61$, $P < 0.001$) and SYP ($r = 0.61$, $P < 0.001$), indicating that these qualities contribute considerably to yield enhancement. SYP showed strong positive associations with DM ($r = 0.29$, $P < 0.05$) and PL ($r = 0.35$, $P < 0.01$), but non-significant with NSP ($r = 0.23$). The SI had a substantial positive correlation with DM ($r = 0.40$, $P < 0.01$).




Fig. 2: Pearson's correlation matrix depicting relationships among yield and its component traits. DM, Days to maturity; NPP, Number of pods per plant; PL, Pod length; NSP, Number of seeds per pod; SI, Seed index; SYP, Seed yield per plant; ns, non-significant.

Genetic parameters

The result revealed that the PCV was slightly higher than the corresponding GCV for all the traits (Table 2). Out of six traits studied, only NPP exhibited high GCV and PCV estimates, whereas three traits, such as PL, SI and SYP, exhibited moderate GCV and PCV estimates. The traits, i.e. DM and NSP, showed low GCV and PCV estimates. In terms of heritability, all the traits showed high heritability. The high GCV, PCV, and heritability coupled with high GAM were recorded for NPP, PL, SI and SYP, whereas the traits such as DM and NSP showed high heritability coupled with moderate GAM, but the extent of GCV and PCV was low to moderate.

GGE biplot analysis

The GGE biplot is an effective approach to identify the ideal genotypes for the target trait under the specific or macro environments correctly. The ideal genotype should have superiority for the target trait in terms of mean performance, in addition to the minimum interactions with the environment. The biplot view ranked the genotypes based on the average yield of test entries over the three years. PC 1 explained about 55.36% variability, whereas PC 2 explained 42.93% of the total variations of the year, which indicated the cumulative variation of 98.29% (Fig. 3A). Each genotype positioned along the AEC axis illustrated its average performance across the environments, while the length of the perpendicular projection from the AEC axis denoted the stability of genotypes. In this study, G27, G48, G25, G47, G5, G50, G22 and G26 had superior mean performance and the most stable genotypes across the environments. The proximity of these genotypes to the ideal genotype position further validated their superior adaptability and reliable performance across the environments. These genotypes thus signify interesting prospects for extensive trials and prospective application in future breeding projects focused on enhancing adaptation and performance stability. The

Table 2: Analysis of genetic parameters of different traits in promising genotypes of mungbean

	DM	NPP	PL	NSP	SI	SYP
Maximum	87.33	38.43	10.35	12.03	5.21	13.17
Minimum	54.67	5.47	5.88	8.11	2.16	4.49
Mean	63.68	20.92	7.72	10.34	3.62	8.43
CD 5%	1.58	2.93	0.25	0.30	0.24	1.00
GCV (%)	8.56	32.60	11.90	6.68	16.51	17.38
PCV (%)	8.70	33.72	12.07	6.91	17.03	18.86
Heritability (bs) (%)	97.00	93.00	97.00	93.00	94.00	85.00

environment plays an important role in the projection of the phenotype in the case of quantitative traits. Therefore, screening of genotypes under multiple environments would be more effective to tag the potential genotypes. Discrimination and representativeness are based on the ability to discriminate the test genotypes, the ability to represent the mega environment and their joint response. In the GGE biplot, the lengths the vectors demonstrated the discrimination ability, while the angle between the environmental vectors and the AEC abscissa shows the representativeness of the testing location. Among the three test environments, E3 had the longest environmental vectors, indicating the most "discriminating year" followed by E2 and E1 (Fig. 3B). However, E2 followed by E1 showed a minimum angle with the average environment, which indicated that these were the most ideal testing years, accounting for both discrimination ability and representativeness. The

results demonstrated the polygon view with six genotypes, namely G27, G42, G2, G32, G44 and G6 at the vertices (Fig. 3C). All three test years were scattered in two sectors within the biplot, which indicated that these environments could be divided into two mega environments (MEs). The ME-I was represented by E1 and E2, whereas ME-II was represented by E3. All three years within the MEs exhibited $<90^\circ$ angle with each other. Most of the mungbean genotypes were placed near the origin, indicating their consistency in terms of yield performance. The genotypes, viz. G42, G17, G49, G18, G27 were found to be the most stable and high yielding over the environments (Fig. 3D).

DNA fingerprinting of selected promising genotypes

DNA fingerprinting of six selected genotypes of mungbean, i.e. G18 (TRAM-18), G42 (BMS 19-3), G49 (ML 134), IPM 02-3, G27 (BMS 18-4), and G17 (Shikha) was done by employing twelve markers. Out of these 12




Fig. 3: GGE biplot of 55 mungbean genotypes for seed yield over the years. 2020 (E1); 2021 (E2), 2022 (E3).

(A) Mean vs stability biplot; (B) Discriminateness vs representativeness biplot; (C) Which-Won-Where biplot; (D) Genotype ranking biplot.

markers, only six markers showed polymorphism (Fig. 4). *VrSnRK2.6c* exhibited a range of 460–475 bp in the test genotypes, the amplicons remained within the size of 180–200 bp and 140–160 bp for *VrSnRK2.1* and *VrHSfA6a*, respectively. Likewise, low polymorphism was exhibited by *VrHSfA6b*, *Mchr3-76* and *HAAS-VR-1437*.




Fig. 4: DNA fingerprinting of promising genotypes of mungbean using markers.
L. Ladder (1kb); 1–6, mungbean genotypes (1. G18 (TRAM-18); 2. G42 (BMS 19-3); 3. G49 (ML 134); 4. IPM 02-3; 5. G27 (BMS 18-4); 6. G17 (Shikha).

Discussion

Evaluation and characterization of the breeding materials followed by assessment of their genetic variations, are the most important activities for crop improvement (Mahajan *et al.*, 2023; Singh & Collins, 2024; Ikram *et al.*, 2024). These help in the effective utilization of trait-specific promising genotypes in breeding programmes. In most of the crops, the genetic variation has been narrowed down significantly due to the repeated and frequent utilization of the common parents in breeding programmes (Singh *et al.*, 2023a; Singh *et al.*, 2023b). Similarly, in mungbean also, many of the modern varieties have been developed by deploying only a narrow spectrum of the existing variability due to frequent use of common parents in the breeding programme (Schreinemachers *et al.*, 2019). Therefore, identification of trait-specific diverse parental lines, which can help in broadening the genetic base of the cultivated lines, should be our prime objective in developing new varieties (Tripathi *et al.*, 2023a; Ogiso-Tanaka *et al.*, 2023; Mahalingam *et al.*, 2023). The present study showed a considerable amount of genetic variability for key agro-morphological traits in the panel of promising mungbean genotypes. The extent and magnitude of genetic variation, along with its usefulness, can be understood by the genetic parameters. In general, estimates of PCV were higher than the corresponding GCV in the present study, as the PCV includes the variability explained by the environment in addition to the GCV. However, the traits

like DM, NPP, PL, NSP, and SI had small differences in their GCV and PCV estimates, which indicated that these traits are mostly governed by genetic factors, and the environment has a little role to play. Similarly, high h^2 bs coupled with high GAM also indicated the preponderance of additive gene action in most of the traits under study. Earlier, breeding methods based on heritability and GAM were also advocated by some of the workers; however, considering the high estimates of GCV, PCV, heritability and GAM, strong association could be exhibited for these traits; therefore, indirect selection by simple plant selection methods would be more effective (Singh *et al.*, 2015). On the other hand, promising accessions for traits showing low or moderate heritability and genetic advance could be improved through hybridization programme.

The results also indicated that the traits like PL, NSP, and SI were strongly associated with SYP, suggesting their strength as selection criteria for improving the SYP of mungbean. Therefore, these traits may be chosen as selection criteria for improving SYP. Similar findings on the strong association of component traits with SYP have also been reported by (Tran and Truong 2023; Bhardwaj *et al.*, 2024; Chauhan *et al.*, 2024).

The performance of the genotypes is always influenced by the environmental conditions where they are grown. SYP is a dependent variable and is an outcome of the cumulative effect of its component traits. Therefore, tagging the key traits with stable performance is more important than the single-environment evaluation. GGE biplot serves a major role in selecting these traits under multiple environments. Therefore, the present investigation was undertaken to identify high-yielding and stable mungbean promising varieties or genotypes over the three years using GGE biplot analysis. Employing the GGE biplot approach to understand GEI is considered to be a systematic approach for isolating the best performing genotypes in accordance with the environment through ranking of phenotypic performance of genotypes and test environments (Yan & Tinker 2006; Arshadi *et al.*, 2018; Oral 2018; Kendal *et al.*, 2019). The discrimination ability of an environment is revealed by the length of environmental vectors, where E3 indicated the most “discriminating year”, whereas E2 followed by E1, showed the minimum angle with the average environment, indicating that these were the most ideal testing environments. The mean vs stability analysis shortlisted the 39 varieties/ promising genotypes over the years and also exhibited superior performance than the check variety IPM 02-3. ‘Which-Won-Where’ analysis is the most efficient way of delineating the GEI of

genotypes through plotting the multi-year or multi-environment data in a polygon view of a GGE biplot (Yan and Kang, 2002). The genotypes, i.e. G42 (BMS 19-3), G17 (Shikha), G49 (ML 134), G18 (TRAM-18), G27 (BMS 18-4), were found to be most stable and high yielding over the years. Singh et al., (2020) also employed a similar approach to select the potential mungbean genotypes. Likewise, Iqbal et al., (2021) also adopted a similar approach of GGE biplot analysis to identify the stable and potential heat-tolerant mungbean lines.

Molecular breeding is one of the tools for characterization, assessing genetic variations without any environmental interruptions. Molecular breeding is now sounder after the decoding of whole genome sequence of mungbean (Kang et al., 2014). Many of the *Vigna* germplasm had been characterized through different types of molecular markers (Pratap et al., 2021; Singh et al., 2021; Sahu et al., 2023; Tripathi et al., 2023b; Kumar et al., 2024). These molecular markers are also used in gene discovery (Purwar et al., 2023; Singh et al., 2023; Sen Gupta et al., 2025), QTL mapping (Subhashree et al., 2024; Reddappa et al., 2025) and marker-assisted breeding (Chugh et al., 2022; Singh et al., 2025). Functional markers are one of them, which can be effectively utilized in assessing genetic variations, tagging genomic regions and also for gene introgression (Salgotra and Stewart 2022; Xue et al., 2024). In the present study, SSRs and functional markers have been used for DNA fingerprinting, and results indicated the significant variations among the promising mungbean genotypes for selected markers. The comprehensive fingerprinting with a large set of markers will help in accelerating the effective utilization of these genotypes in the breeding programme for yield improvement.

Conclusion

The study revealed that PL, NPP and SI were observed to be the key traits over the years, hence may be chosen as the selection criteria for improving the yield. Based on GGE biplot analysis, five genotypes, i.e. G42 (BMS 19-3), G17 (Shikha), G49 (ML 134), G18 (TRAM-18), and G27 (BMS 18-4) were found to be the most stable and high-yielding ones over the years. These genotypes should be used as the potential parents for mungbean improvement. Further, BMS 18-4 and BMS 19-3 are new promising entries and could be evaluated further in varietal yield trials for their potential as new varieties.

Acknowledgement

Authors are grateful for the financial support from the Ministry of Agriculture, Agriculture Education and

Research, Govt. of Uttar Pradesh, under the project "Center of Excellence in Dryland Agriculture (CEDA/36-2018)".

References

- Abou-Khater LF Maalouf, A Jighly, D Rubiales and S Kumar (2022) Adaptability and stability of faba bean (*Vicia faba* L.) accessions under diverse environments and herbicide treatments. *Plants* 11(3): 251.
- Altat MT (2026) Assessing mungbean genetic diversity with start codon targeted (SCoT) markers; a step towards climate-tolerant varieties for global food security. *BMC Plant Bio.* 26: 531. <https://doi.org/10.1186/s12870-026-08357-5>
- Arshadi A, E Karami, A Sartip, M Zare and P Rezabakhsh (2018) Genotypes performance in relation to drought tolerance in barley using multi-environment trials. *Agron. Res.* 16: 521. <http://hdl.handle.net/10492/3785>
- Arya M, SB Mishra, KM Maring and R Kant (2024) Stability analysis for biological nitrogen fixation and seed yield in mungbean [*Vigna radiata* (L.) Wilczek] genotypes. *Legume Res.* 47(2): 201-205. Doi: 10.18805/LR-5256
- Bhardwaj R, Gayacharan, BH Gawade, P Pathania, A Talukdar, P Kumar and GP Singh (2024) Identification of heat-tolerant mungbean genotypes through morpho-physiological evaluation and key gene expression analysis. *Front. Genet.* 15: 1482956.
- Bomma N, HB Shruthi, CD Soregaon, A Gaddameedi, K Suma, J Pranati and PI Gangashetty (2024) Multi-environment testing for G × E interactions and identification of high-yielding, stable, medium-duration pigeonpea genotypes employing AMMI, GGE biplot, and YREM analyses. *Front. PlantSci.* 15: 1396826.
- Chauhan DS, SK Verma, RK Panwar, A Arora, K Pragati, A Bhatt and A Mehra (2024) Multivariate analysis of genetic diversity in mung bean [*Vigna radiata* (L.) Wilczek] using Mahalanobis statistic. *Int J Plant Soil Sci.* 36(8): 1138-1145.
- Chugh V, D Kaur, S Purwar, P Kaushik, V Sharma, H Kumar and RB Dubey (2022) Applications of molecular markers for developing abiotic-stress-resilient oilseed crops. *Life* 13(1): 88.
- Diatta AA, WE Thomason, O Abaye, TL Thompson, ML Battaglia, LJ Vaughan and JF Filho (2020) Assessment of nitrogen fixation by mungbean genotypes in different soil textures using 15N natural abundance method. *J. Soil Sci. Plant Nutr.* 20(4): 2230-2240.
- Ghallab DS and EH Zaghloul (2025) Metabolomic insights into phytochemical diversity among the different parts of *Mangrove Avicennia marina* and their prebiotic effects on marine probiotics growth. *Food Biosci.* 71: 107229.
- Ikram S, S Bhattarai and KB Walsh (2024) Screening new mungbean varieties for terminal drought tolerance. *Agriculture* 14(8): 1328.
- Iqbal J, G Shabbir, KN Shah and A Qayyum (2021) Deciphering of genotype × environment interaction to identify stable heat-tolerant mung bean genotypes by GGE biplot analysis. *J. Soil Sci. Plant Nutr.* 21(3): 2551-2561.
- Irfan M, MA Bhat, U Rashid, KM Alwutayd, FA Safhi, NM Al Aboud and MA Wani (2026) Multi-environment stability and genotype × environment interaction analysis for grain yield and micronutrient (Fe and Zn) contents in mungbean (*Vigna*

- radiata* L.). *BMC Plant Biol.* 26(1): 292.
- Jafarullakhan SS, V Vijayakumar, KV Singh, NA Saravanan, VA Giridhari, S Rathinavelu and S Rajaprakasam (2024) Assessing the yield and nutrient potential of horse gram mutants (*Macrotyloma uniflorum* Lam. Verdc.), an underutilized legume through a multi-environment-based experiment. *Sci. Rep.* 14(1): 16305.
- Kang YJ, SK Kim, MY Kim, P Lestari, KH Kim, BK Ha and SH Lee (2014) Genome sequence of mungbean and insights into evolution within *Vigna* species. *Nat. Commun.* 5(1): 5443.
- Kendal E, M Karaman, S Tekdal and S Doğan (2019) Analysis of promising barley (*Hordeum vulgare* L.) lines performance by AMMI and GGE biplot in multiple traits and environment. *Appl. Ecol. Environ. Res.* 17(2): 5219-5233.
- Kohli M, H Bansal, GP Mishra, HK Dikshit, SB Reddappa, A Roy and M Aski (2024) Genome-wide association studies for earliness, MYMIV resistance, and other associated traits in mungbean (*Vigna radiata* L. Wilczek) using genotyping by sequencing approach. *PeerJ.* 12: e16653.
- Kona P, BC Ajay, K Gangadhara, N Kumar, RR Choudhary, MK Mahatma and KD Solanki (2024) AMMI and GGE biplot analysis of genotype by environment interaction for yield and yield contributing traits in confectionery groundnut. *Sci. Rep.* 14(1): 2943.
- Kumar H, V Chugh, M Kumar, V Gupta, S Prasad, S Kumar and M Kumar (2023a) Investigating the impact of terminal heat stress on contrasting wheat cultivars; a comprehensive analysis of phenological, physiological, and biochemical traits. *Front. Plant Sci.* 14: 1189005.
- Kumar S, H Kumar, V Gupta, A Kumar, CM Singh, M Kumar and R Kumar (2023b) Capturing agro-morphological variability for tolerance to terminal heat and combined heat-drought stress in landraces and elite cultivar collection of wheat. *Front. Plant Sci.* 14: 1136455.
- Kumar S, CM Singh, M Kumar, S Purwar, H Kumar, A Mishra and A Pratap (2024) Identification and characterization of potential sources of waterlogging tolerance in *Vigna* species as revealed by physiological, biochemical and gene expression profiling. *J. Plant Biochem Biotechnol.* 33(4): 657-673.
- Kyu KL, CM Taylor, CA Douglas, AI Malik, TD Colmer, KH Siddique and W Erskine (2024) Genetic diversity and candidate genes for transient waterlogging tolerance in mungbean at the germination and seedling stages. *Front. Plant Sci.* 15: 1297096.
- Ma C, C Liu and Z Ye (2024) Influence of genotype × environment interaction on yield stability of maize hybrids with AMMI model and GGE biplot. *J. Agron.* 14(5): 1000.
- Mahajan G, K Wenham and BS Chauhan (2023) Mungbean (*Vigna radiata*) growth and yield response in relation to water stress and elevated day/night temperature conditions. *J. Agron.* 13(10): 2546.
- Mahalingam A and N Manivannan (2023) Interspecific hybridization between *Vigna radiata* and *Vigna mungo* towards the broadening of genetic base for MYMV disease resistance and generating variability. *Legume Res.* 46(4): 395-402.
- Mohammadi S, M Lillemo, Å Ergon, S Shafiee, S Zanutto, JA Dieseth and AK Uhlen (2024) Unveiling phenotypic and environmental dynamics; exploring genetic stability and adaptability of faba bean cultivars in Norwegian climates. *Legume Sci.* 6(4): e70008.
- Nair RM, AK Pandey, AR War, B Hanumantharao, T Shwe, AKMM Alam and R Schafleitner (2019) Biotic and abiotic constraints in mungbean production progress in genetic improvement. *Front. Plant Sci.* 10: 1340.
- Ogiso-Tanaka E, S Chankaew, Y Yoshida, T Isemura, R Marubodee, A Kongjaimun and N Tomooka (2023) Unique salt-tolerance-related QTLs, evolved in *Vigna riukiuensis* (Na⁺ includer) and *V. nakashimae* (Na⁺ excluder), shed light on the development of super-salt-tolerant azuki bean (*V. angularis*) cultivars. *Plants.* 12(8): 1680.
- Oral E (2018) Effect of nitrogen fertilization levels on grain yield and yield components in triticale based on AMMI and GGE biplot analysis. *Appl. Ecol. Environ. Res.* 16(4): 4865-4878.
- Ortiz R, F Reslow, J Huicho, R Vetukuri and J Crossa (2024) Adaptability, stability, and productivity of potato breeding clones and cultivars at high latitudes in Europe. *Discov. Life.* 54(1): 13.
- Parihar AK, S Gupta, KK Hazra, A Lamichaney, D Sen Gupta, D Singh and S Das (2022) Multi-location evaluation of mungbean (*Vigna radiata* L.) in Indian climates; ecophenological dynamics, yield relation, and characterization of locations. *Front. Plant Sci.* 13: 984912.
- Pookhamsak P, T Girdthai, A Jomsangawong, A Masari, S Ngampongsai, A Tharapreuksapong and PA Tantasawat (2026) Development of a new powdery mildew and *Cercospora* leaf spot-resistant mungbean cultivar using marker-assisted backcross breeding. *J. Genet. Eng. Biotechnol.* 24(2): 100698.
- Pratap A, I Dhaliwal, CM Singh, A Mahalingam, N Manivannan, T Basavaraja and A Tripathi (2022) Biofortification of mungbean. In: *Biofortification of Staple Crops*. pp 295-333, Springer, Singapore.
- Pratap A, S Gupta, PS Basu, R Tomar, S Dubey, M Rathore and G Kumari (2019) Towards development of climate smart mungbean; challenges and opportunities. In: *Genomic Designing of Climate-Smart Pulse Crops*, pp 235-264.
- Pratap A, CM Singh, S Gupta, AK Gupta, RS Birader, U Prajapati and NP Singh (2021) Genetic enhancement in mungbean (*Vigna radiata*) as revealed by genome-wide mapped microsatellite markers. *Agric. Res.* 10(3): 369-377.
- Purwar S, CM Singh, M Kumar, AK Singh, A Pratap, P Singh and NP Singh (2023) Genome-wide identification and analysis of NBS-LRR-encoding genes in mungbean (*Vigna radiata* L. Wilczek) and their expression in two wild non-progenitors reveal their role in MYMIV resistance. *J. Plant Growth Regul.* 42(10): 6667-6680.
- Raizada A and J Souframanien (2019) Transcriptome sequencing, de novo assembly, characterisation of wild accession of blackgram (*Vigna mungo* var. *silvestris*) as a rich resource for development of molecular markers and validation of SNPs by high resolution melting (HRM) analysis. *BMC Plant Biol.* 19(1): 358.
- Rao PJM, N Sandhyakishore, S Srinivasan, S Sandeep, G Praveen, G Neelima and GA Kumar (2023) AMMI and GGE stability analysis of drought tolerant chickpea (*Cicer arietinum* L.) genotypes for target environments. *Legume Res.* 46(9): 1105-1116.
- Reddappa SB, MS Aski, GP Mishra, S Gupta, KM Shivaprasad, SK Sinha and HK Dikshit (2025) QTL mapping for yield contributing traits in mungbean (*Vigna radiata* L.) using a

- RIL population. *Sci. Rep.* 15(1):20795.
- Sadaiah K, GE Reddy, K Parimala, A Saritha, G Madhuri, VD Rani and L Krishna (2026) Genotype $\times$ environment interaction insights for yield and yield components in castor via AMMI and GGE biplot models. *Sci. Rep.* DOI: 10.1038/s41598-026-44030-5
- Sahu TK, SK Verma, Gayacharan, NP Singh, DC Joshi, DP Wankhede and AK Singh (2023) An associative transcriptomics study on rice bean (*Vigna umbellata*) provides new insights into genetic basis and candidate genes governing flowering, maturity and seed weight. *bioRxiv*: 2023-02.
- Salgotra RK and CN Stewart Jr (2022) Genetic augmentation of legume crops using genomic resources and genotyping platforms for nutritional food security. *Plants*, 11(14): 1866.
- Sao A and S Nair (2025) Identification of bacterial leaf blight resistance (*Xanthomonas oryzae* pv. *oryzae*) genes in advanced breeding lines of rice (*Oryza sativa* L.) through molecular markers. *Int. J. Bio-resour. Stress Manag.* 16(1): 1-7.
- Schreinemachers P, T Sequeros, S Rani, MA Rashid, NV Gowdru, MS Rahman and RM Nair (2019) Counting the beans: quantifying the adoption of improved mungbean varieties in South Asia and Myanmar. *Food Secur.* 11(3): 623-634.
- Sengupta J and CM Hussain (2025) MXene-based electrochemical biosensors: advancing detection strategies for biosensing (2020–2024). *Biosensors*. 15(3): 127.
- Shobeiri SS, P Pezeshkpour and B Naseri (2024) Evaluation of seed yield stability of lentil genotypes by linear mixed-effects models and multitrait stability index. *Legume Sci.* 6(3): e245.
- Singh CM, SB Mishra, A Pandey and M Arya (2015) Multivariate analysis in mungbean [*Vigna radiata* (L.) Wilczek] to identify the genetic donors for pubescence and agro-morphological traits. *Legume Res.* 38(6): 767-771.
- Singh CM, U Prajapati, S Gupta and A Pratap (2021) Microsatellite-based association mapping for agronomic traits in mungbean (*Vigna radiata* L. Wilczek). *J. Genet.* 100(2): 87.
- Singh CM, A Pratap, H Kumar, S Singh, BK Singh, D Prasad and M Kumar (2022) Recent advances in omics approaches for mungbean improvement. In: *Technologies in Plant Biotechnology and Breeding of Field Crops*, pp 181-200.
- Singh CM, S Purwar, AK Singh, BK Singh, M Kumar, H Kumar and KH Baek (2023) Analysis of auxin-encoding gene family in *Vigna radiata* and its cross-species expression modulating waterlogging tolerance in wild *Vigna umbellata*. *Plants* 12(22): 3858.
- Singh J, A Sharma, P Sharma and N Kumar (2023a) Genetic variability and association studies in mid-late and late group of cauliflower (*Brassica oleracea* L. var. botrytis). *Indian J. Plant Genet. Resour.* 36(1): 45-51.
- Singh PK, N Kumari, RK Jha and R Sadhukhan (2023b) Assessment of phenotypic variations and correlation among yield attributing and biochemical traits in mutant population of grasspea (*Lathyrus sativus* L.). *Indian J. Plant Genet. Resour.* 36(1): 77-84.
- Singh P, B Pandey, S Purwar, CM Singh, A Pratap and AK Mishra (2025) Exogenous application of γ -aminobutyric acid alleviates temperature stress in mungbean (*Vigna radiata*) and its wild non-progenitor (*Vigna glabrescens*) by regulating heat shock protein genes. *BMC Plant Biol.* 25(1): 1651.
- Singh V and M Collins (2024) Genotypic variability in response to heat stress and post-stress compensatory growth in mungbean plants [*Vigna radiata* (L.) Wilczek]. *Crops* 4(3): 270-287.
- Subhashree P, JP Sahoo, M Mahapatra, BP Das, A Parween and SK Sunani (2024) Morpho-biochemical and molecular characterization of new mungbean [*Vigna radiata* (L.) Wilczek] landraces for Cercospora leaf spot (CLS) disease resistance. *Physiol. Mol. Plant Pathol.* 134: 102448.
- Torres-Ordoñez LH, JD Valenzuela-Cobos, F Guevara-Viejo, P Galindo-Villardón and P Vicente-Galindo (2024) Effect of genotype $\times$ environment interactions on the yield and stability of sugarcane varieties in Ecuador: GGE biplot analysis by location and year. *Appl. Sci.* 14(15): 6665.
- Tran MTT and NT Truong (2023) Evaluation of genetic characteristics of introduced mung bean varieties based on agronomic traits. *Eng. Proc.* 56(1): 233.
- Tripathi A, CM Singh, M Kumar, S Purwar, A Mishra, D Kumar and NP Singh (2023a) Identification of potential sources of mungbean yellow mosaic India virus resistance in black gram (*Vigna mungo*) and expression of antioxidants and R-genes modulating resistance response in cultivated and its two wild relatives. *Plant Breed.* 142(5): 668-681.
- Tripathi K, RK Pamarthi, DP Semwal, V Tyagi, P Brahmi and A Pandey (2023b) Efforts for the revival of *Vigna subterranea* (L.) Verdc. (Bambara groundnut) through germplasm introduction in India. *Indian J. Plant Genet. Resour.* 36(1): 116-121.
- Wodebo KY, T Tolemariam, S Demeke, W Garedew, T Tesfaye, M Zeleke and M Sharma (2023) AMMI and GGE biplot analyses for mega-environment identification and selection of some high-yielding oat (*Avena sativa* L.) genotypes for multiple environments. *Plants* 12(17): 3064.
- Xue B, Z Liang, Y Liu, D Li and C Liu (2024) Genome-wide identification of the RALF gene family and expression pattern analysis in *Zea mays* (L.) under abiotic stresses. *Plants* 13(20): 2883.
- Yan W and MS Kang (2002) *GGE Biplot Analysis: A Graphical Tool for Breeders, Geneticists, and Agronomists*. CRC Press, pp 288.
- Yan W and NA Tinker (2006) Biplot analysis of multi-environment trial data: principles and applications. *Can. J. Plant Sci.* 86(3): 623-645.
- Zafar SH, M Umair and M Akhtar (2023) Nutritional evaluation, proximate and chemical composition of mungbean varieties/cultivars pertaining to food quality characterization. *Food Chem. Adv.* 2: 100160.

Supplementary Table1: Performance of mungbean promising lines for yield and yield component traits over the three years

Genotype	Source	Days to maturity	Number of pods per plant	Pod length (cm)	Number of seeds per pod	Seed index (g)	Seed yield per plant (g)
PDM 139	IIPR, Kanpur	59.56	24.07	6.72	10.40	3.48	5.85
PDM 54	IIPR, Kanpur	71.22	26.48	6.72	10.41	3.42	8.23
PDM 178	IIPR, Kanpur	65.11	33.96	6.87	10.61	3.70	6.18
IPM 2-14	IIPR, Kanpur	59.78	12.28	6.14	9.20	2.71	5.99
IPM 06-5	IIPR, Kanpur	60.11	22.95	8.39	11.23	4.27	8.99
IPM 409-4	IIPR, Kanpur	65.33	22.33	7.76	10.40	3.14	5.36
IPM 312-43K	IIPR, Kanpur	59.78	21.75	7.29	10.38	3.03	7.27
IPM 03-1	IIPR, Kanpur	60.44	24.01	8.10	10.32	3.24	6.35
IPM 03-3	IIPR, Kanpur	60.89	34.47	6.77	10.50	3.34	6.49
IPM 2-17	IIPR, Kanpur	60.78	37.74	7.96	10.18	3.29	8.56
IPM 02-3-2	IIPR, Kanpur	58.78	25.64	6.80	10.15	3.23	9.90
IPM 2-19	IIPR, Kanpur	60.44	21.80	6.65	9.67	3.09	9.97
IPM 05-3-22	IIPR, Kanpur	64.56	18.57	7.59	10.68	3.65	6.77
IPM 306-6	IIPR, Kanpur	62.34	23.07	7.00	10.01	3.27	6.64
Yellow Selection	IIPR, Kanpur	64.45	20.62	6.82	10.98	3.19	7.93
ML 1257	IIPR, Kanpur	74.78	25.09	6.91	9.67	3.70	6.12
Shikha	IIPR, Kanpur	59.11	25.15	7.83	10.51	3.24	11.14
TARAM 18	BARC, Mumbai	61.44	28.23	7.22	10.18	2.94	10.65
TMB 37	BARC, Mumbai	60.22	22.77	7.46	9.70	2.61	8.46
SPS-5	IIPR, Kanpur	69.67	10.84	9.01	11.35	4.23	8.13
SM 48	IIPR, Kanpur	59.56	13.15	8.44	11.39	4.17	10.36
PM-4	IIPR, Kanpur	59.78	17.35	8.34	11.41	3.74	9.74
UPM 98-1	IIPR, Kanpur	59.33	14.78	8.21	8.64	3.15	8.50
UPM02-17	IIPR, Kanpur	60.78	26.48	8.63	11.06	3.20	8.42
HUM 12	BHU, Varansi	67.78	16.84	6.96	9.94	3.59	8.74
BMS 18-3	BUAT, Banda	74.00	26.40	9.83	11.38	5.11	9.91
BMS 18-4	BUAT, Banda	75.45	15.48	8.29	11.09	4.73	9.87
EC 391178(y)	IIPR, Kanpur	66.66	15.08	7.51	9.86	3.43	10.46
BMS 18-5	BUAT, Banda	84.00	12.94	7.25	10.68	3.91	7.04
EC 520034	IIPR, Kanpur	68.22	12.06	8.32	10.02	3.70	10.20
GM 4	GAU, Gujrat	59.22	8.98	7.28	9.57	3.61	9.38
HW 421	IIPR, Kanpur	59.33	7.92	8.09	10.88	4.16	9.00
V1133	AVRDC, Hyderabad	61.33	22.43	7.44	10.56	4.10	9.33
China Mung 2	IIPR, Kanpur	61.11	10.76	7.86	10.33	3.85	9.13
IPOI 539	IIPR, Kanpur	60.66	22.43	6.77	9.50	3.08	7.55
F8(MXU) Extra Early	IIPR, Kanpur	60.11	20.96	7.61	10.14	3.76	8.02
MS 03-18	--	59.56	8.43	7.34	10.68	3.07	9.19
MH 805	CCSHAU, Hissar	60.22	28.55	8.36	11.03	4.14	9.86
SML 1815	PAU, Ludhiana	61.11	14.83	8.23	10.40	3.38	10.43
AKM/NP/8/9	PDKV, Akola	59.22	21.08	8.17	10.91	3.14	7.97
IPM 312-19	IIPR, Kanpur	58.22	27.46	7.34	8.62	3.08	7.89
BMS 19-3	BUAT, Banda	75.44	12.65	10.12	11.82	4.87	11.71
DGGV-2	UAS, Dharwad	63.78	16.69	9.37	11.39	5.16	8.02
BMS 18-17	BUAT, Banda	62.89	15.59	9.47	10.35	4.59	6.20
IC 296672	IIPR, Kanpur	62.78	26.56	6.20	8.85	3.09	9.43

Brazil	Exotic collection	61.22	17.88	8.84	10.90	5.10	9.29
MGG-353	ARS, Mandira	60.22	20.34	7.20	10.38	3.37	8.63
EC 520022	IIPR, Kanpur	64.11	28.05	10.25	10.67	4.53	8.99
ML 134	PAU, Ludhiana	60.11	34.35	7.91	10.86	3.31	10.99
ML 1299	PAU, Ludhiana	64.56	31.36	7.01	10.42	3.09	8.97
HUM 16	BHU, varansi	63.22	24.06	7.54	10.09	3.27	7.19
VIRAT	IIPR, Kanpur	56.89	21.74	6.61	9.28	3.06	6.66
IPM 2-3	IIPR, Kanpur	71.00	15.10	7.22	9.91	3.70	7.43
SML 668	PAU, Ludhiana	71.33	17.21	7.53	9.96	3.79	7.41
IPM 99-125	IIPR, Kanpur	70.44	23.01	7.06	9.38	3.42	6.91

Supplementary Table 2: List of primers used in DNA fingerprinting

S. No.	Primers	Type of marker	Forward	Reverse	Annealing Temperature (°C)
a)	VrSnRK2.6c	Functional	GGTGTGAACGAGGATGTCGTA	CTCCTTCAGAGTCTCGTATCGT	56
b)	VrSnRK2.1	Functional	GTGAGGATGAGGCACGCTT	AACGAGGAGCTGGACTACCA	56
c)	VrHSfA6a	Functional	GCTTGCTATGAACATGGAGGA	TCATTATCATCATTTTCCAATGC	56
d)	VrHSfA6b	Functional	GAGCAAATTGGAAGCCAGAA	AGTTGCCGAGCCAATACATC	54
e)	Mchr3-76	SSR	GAAATAAGCCTAAAGACCAG	GAAGTGCATCTGTCTATCCC	53
f)	HAAS-VR-1437	SSR	CCCCAGAATCAAGAAATCCA	GCTCTCGTTACTGGAGGCAC	56
g)	VrDREB5	Functional	CAATGCCACCATTCCCCCAT	TCCGAAGGAGAATAGGCCGT	57
h)	VrDREB12	Functional	TCACTTCTGTACCTCCTCCA	AACGCCTCCAAGTCATACCA	57
i)	VrDREB13	Functional	ATTCGTCACCCTTTGCTGAAG	TTGTCTTGGCATTTCGACCA	55
j)	VrDREB22	Functional	CCAATCCATTTCCTCTGCC	CCACGCTAACCCAATTCCCT	57
k)	VrDREB30	Functional	TCAACTTTCCCGACGAGGTG	ACTGCCATCCGCTTTCATCA	56
l)	SKPI	Functional	TGGCTGCCAACTACCTGAACA	ACATGTTCAACGACACCCACCT	59

RESEARCH ARTICLE

Aroma and Agro-morphological Diversity in Traditional Rice (*Oryza sativa*) Germplasm from Eastern India

Pooja Belwal^{1,2}, Amrita Banerjee¹, Anima Mahato², Priyamedha¹, Sougata Bhattacharya², Nabaneeta Basak³, Nimai Prasad Mandal¹ and Somnath Roy^{1*}

Abstract

Traditional aromatic and quality rice (*Oryza sativa* L.) landraces from eastern India harbour valuable grain quality traits, yet their agro-morphological diversity and fragrance genetics remain insufficiently characterized. This study evaluated 183 accessions from the Indian National GeneBank to characterize phenotypic variation and assess allelic diversity at the *BADH2* fragrance locus. Twenty-two agro-morphological traits were recorded, aroma was assessed using the KOH test, and allele-specific PCR was employed to detect the 8-bp functional deletion in exon 7 of *BADH2*. The accessions exhibited wide phenotypic variation, particularly in grain length, grain width, kernel shape, and pigmentation traits. Aroma profiling classified 42% of accessions as strongly aromatic and 36% as moderately aromatic. Multiple-factor analysis and clustering showed weak state-wise structuring, indicating considerable overlap among accessions, likely reflecting historical seed exchange. Genotyping showed that 31% of accessions carried the homozygous aromatic allele; however, a substantial proportion of aromatic accessions lacked the deletion, suggesting that additional genetic factors may contribute to fragrance in this germplasm. These findings highlight the diversity of agro-morphological and aroma-related traits in eastern Indian rice and provide a useful baseline for future genetic studies and quality-oriented breeding programmes, as well as for the conservation and utilization of the aromatic landraces.

Keywords: Aromatic rice, Agro-morphological diversity, *BADH2* gene, Eastern India germplasm, Landraces

¹Central Rainfed Upland Rice Research Station, ICAR-Central Rice Research Institute, Hazaribag, Jharkhand, 825301, India.

²ICAR-Indian Agricultural Research Institute, Jharkhand Campus, Gauria Karma, Jharkhand, 825405, India.

³Crop Physiology and Biochemistry Division, ICAR-Central Rice Research Institute, Cuttack, Odisha, 753006, India.

*Authors for correspondence:

Somnath.Roy@icar.org.in; sroypr@gmail.com

Received: 30/03/2026

Revised: 10/04/2026

Accepted: 13/04/2026

How to cite this article: Belwal P, A Banerjee, A Mahato, Priyamedha, S Bhattacharya, N Basak, NP Mandal and S Roy (2026) Aroma and Agro-morphological Diversity in Traditional Rice (*Oryza sativa*) Germplasm from Eastern India. *Indian J. Plant Genet. Resour.* 39(2): 20-28.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.177538>

Introduction

Rice is cultivated in more than 117 countries and remains central to global food security, particularly across Asia, Africa, and Latin America. India, a secondary centre of origin for *Oryza sativa*, conserves remarkable genetic variability in both cultivated rice and its wild relatives (Bisne and Sarawgi, 2008; Roy *et al.*, 2020). Among the major varietal groups, aromatic rices occupy a premium niche due to their distinctive fragrance, superior cooking quality, and strong consumer preference. Globally traded types such as Basmati rice from the Indian subcontinent and Jasmine rice from Thailand command high market value, with India exporting over 5.2 million metric tons of Basmati rice valued at USD 5.9 billion in 2024-25 (<https://apeda.gov.in/BasmatiRice>), underscoring the economic importance of fragrant rice.

The key biochemical compound responsible for fragrance, 2-acetyl-1-pyrroline (2-AP), accumulates due to loss-of-function mutations in the *BADH2* gene (Bradbury *et al.*, 2005a). The most widespread allele is the 8-bp deletion in exon 7 (*badh2.1*), originally identified in *Japonica* germplasm and now common across aromatic *Indica* groups (Kovach *et al.*, 2009).

Additional *BADH2* mutations, including G-to-T SNP (single nucleotide polymorphism) in exon 10 (*badh2.7*) and a 1-bp insertion in exon 14 (*badh2.6*), are reported particularly in eastern and north-eastern India (Kovach *et al.*, 2009; Chandanshive *et al.*, 2024). However, *BADH2* variation does not always correlate perfectly with sensory aroma scores, reflecting environmental influences and the possible involvement of other genetic loci (Sakthivel *et al.*, 2009; Prodhan and Qingyao, 2020).

Beyond Basmati, India hosts a wealth of non-Basmati aromatic rice, especially short- and medium-grain landraces cultivated in eastern states such as West Bengal, Bihar, Jharkhand, Odisha, Chhattisgarh, and Assam. Popular examples include 'Gobindabhog', 'Tulaipanji', 'Badshabhog', 'Katarni', 'Sonachur', 'Jeeraphool', and 'Kalajeera'. These landraces are deeply embedded in local food traditions and valued for grain appearance, aroma, and cooking quality. Despite their cultural and economic relevance, they have received limited attention in systematic breeding and characterization efforts. Only a few traditional aromatic varieties, such as 'Gobindabhog', 'Tulaipanji', 'Marcha Dhan', 'Jeeraphool', 'Nagri Dubraj', and 'Koraput Kalajeera', have attained Geographical Indication (GI) status, and none from Jharkhand have yet been recognized. However, many non-basmati landraces have been reported from Jharkhand, including 'Prasad bhog', 'Nanhia', 'Tulshimanjar', 'Baans Phul' and 'Shyam jeera' (Sinha *et al.*, 2016). Lack of formal morphological and genetic characterization constrains their documentation, legal protection, and utilization in quality-oriented breeding.

Aromatic landraces from eastern India exhibit wide morphological diversity, shaped by farmer-driven selection across heterogeneous landscapes. Significant variability has been reported for grain size and shape, awning, pigmentation, plant stature, and flowering behaviour (Ray *et al.*, 2013; Roy *et al.*, 2015; Roy *et al.*, 2016; Behera and Panda, 2023). However, this variation is often poorly structured geographically due to frequent seed exchange and inconsistent naming conventions. Landraces with identical names may be genetically unrelated, whereas distinct names may refer to closely related types. As a result, phenotypic identity and distinctiveness remain unclear, underscoring the need for integrated morphological and molecular approaches.

Despite long-standing interest, comprehensive studies that combine multi-trait agro-morphological evaluation with functional genotyping of the *BADH2* fragrance allele remain scarce in this region. Moreover,

emerging evidence suggests that a substantial proportion of mildly or strongly aromatic eastern Indian cultivars lack the common 8-bp deletion allele (Prodhan and Qingyao, 2020; Chandanshive *et al.*, 2024), indicating the possible presence of additional *BADH2* haplotypes or alternative fragrance mechanisms that remain unexplored.

To address these gaps, the present study evaluates 183 traditional rice accessions from the Indian National GeneBank, predominantly short-grain aromatic and quality landraces originating from Jharkhand, West Bengal, Bihar, Odisha, Chhattisgarh, and Assam. Using 22 agro-morphological descriptors, sensory aroma assessment, and allele-specific PCR targeting the 8-bp deletion in *BADH2*, this study aims to: (i) document the extent and pattern of agro-morphological diversity, (ii) determine the prevalence of the 8-bp FNP in *BADH2* and its association with phenotypic aroma, and (iii) identify unique or distinct landraces with potential for registration, conservation, and quality-breeding programmes. By integrating morphological and molecular evidence, this work provides a clearer understanding of the diversity, fragrance genetics, and distinctiveness of eastern Indian aromatic rice, offering a foundation for GI protection, conservation strategies, and future genomic investigation of fragrance-related loci.

Materials and Methods

Plant material

A total of 219 traditional rice accessions originating from eastern Indian states and Assam were procured from the Indian National GeneBank, ICAR-National Bureau of Plant Genetic Resources (NBPGR), New Delhi. Passport information and collection site coordinates were retrieved from the NBPGR PGR Portal. Initial accession selection was based on vernacular names reported in literatures to represent aromatic or high-quality short-grain landraces. Because accessions with the same vernacular name may differ across collection sites, multiple entries of some landrace names were intentionally included. Before detailed characterization, all 219 accessions were subjected to a preliminary germination test and seed lot examination. Of these, 183 accessions were retained for analysis based on two criteria: (i) satisfactory germination, and (ii) minimal visible seed mixture within the accession (Supplementary Table 1). Accessions showing very low (<5%) germination and/or obvious within-accession seed heterogeneity were excluded. The final set represented collections from Jharkhand, West Bengal,

Bihar, Odisha, Chhattisgarh, and Assam (Fig. 1).



Fig. 1: Collection sites of rice accessions in eastern India, with the inset depicting the count per state.

The map was generated from site coordinates obtained from ICAR-NBPGR and visualized in Google Earth.

Experimental site and growth conditions

The agro-morphological evaluation was conducted during the summer season (February–June 2025) at the phenotyping facility (net-house) of the Central Rainfed Upland Rice Research Station (CRURRS), ICAR–Central Rice Research Institute (CRRI), Hazaribag, Jharkhand. The experiment was laid out in a completely randomized design (CRD). For each accession, three healthy seedlings were transplanted individually into separate 10-kg soil-filled pots and maintained as three biological replicates. Seedlings were first raised in nursery trays and 21-day-old uniform seedlings were transplanted into pots. Pots were randomly arranged within the phenotyping facility (under natural conditions) and periodically repositioned during crop growth to minimize positional or micro-environmental bias. Standard agronomic management was followed, including fertilizer applications of 60:40:40 kg N:P₂O₅:K₂O/ha, irrigation and need-based plant protection measures.

Agro-morphological characterization

A total of 22 phenotypic traits were recorded following the Standard Evaluation System (SES) for Rice (IRRI, 2013), comprising 18 qualitative descriptors and four quantitative grain traits. The qualitative traits included leaf blade pubescence, leaf blade colour, basal leaf sheath colour, leaf angle, pigmentation traits (ligule, collar, and auricle colour), apiculus colour, lemma–palea colour, sterile lemma colour, awn presence and pigmentation, pericarp colour, kernel shape and aroma intensity. The grain traits included

grain length, grain width, kernel length, and 100-grain weight. Quantitative grain traits were measured across the three biological replicates, using 10 grains or kernels per accession per replicate. The 100-grain weight was recorded using well-filled dried grains with an approximate moisture content of 14%.

Aroma was assessed using the 1.7% KOH sensory test following Nagaraju *et al.*, (1991) and IRRI SES guidelines. In short, 2 g of dehusked grains were treated with 10 mL of 1.7% KOH and aroma was scored after 10 min relative to the aromatic check 'Pusa Basmati 1'. Aroma intensity was classified into three categories: 1, absent or very weak; 2, weak; and 3: strong. For accessions that did not flower, aroma was assessed from leaf tissue using the same KOH-based procedure. To minimize scorer bias and olfactory fatigue, samples were evaluated in small batches by three independent evaluators, and the scores were cross-checked before final classification.

Genotyping for BADH2 exon 7, 8-bp deletion

The aroma-linked 8-bp deletion in exon 7 of the *BADH2* gene was detected using the allele-specific amplification (ASA) PCR assay developed by Bradbury *et al.*, (2005b). Genomic DNA was extracted from 21-day-old seedlings using the DNeasy Plant Mini Kit (Qiagen, India). DNA concentration and quality were assessed using a NanoDrop spectrophotometer, and samples were diluted to a working concentration of 50 ng/μl. PCR amplification was performed using four primers: external sense primer (ESP), external antisense primer (EAP), internal fragrant antisense primer (IFAP), and internal non-fragrant sense primer (INSP), following the thermal cycling conditions described by Bradbury *et al.*, (2005b). This primer combinations enable allele-specific amplification of the *BADH2* exon 7 region, producing three diagnostic fragments: a 257 bp fragment corresponding to the aromatic allele (ESP + IFAP), a 355 bp fragment corresponding to the non-aromatic allele (EAP + INSP), and a 580 bp fragment serving as a positive control (ESP + EAP). PCR products were separated on 2% agarose gels stained with DNA-Safedye (SRL, India) and visualized using a Bio-Rad gel documentation system. Based on the observed banding patterns, genotypes were classified as: (i) homozygous non-aromatic (580+355 bp), (ii) heterozygous aromatic/non-aromatic (580+355+257 bp), or (iii) homozygous aromatic (580+257 bp).

Statistical analyses

Descriptive statistics for quantitative traits, including mean, range, standard deviation, coefficient of

variation, and skewness, were computed in R v4.4.3. Differences among accessions for quantitative traits were tested using one-way analysis of variance under the model: $Y_{ijk} = \mu + G_i + R_j + E_{ijk}$ where, Y_{ijk} is trait value, μ is overall mean, G_i is effect of the i^{th} genotype, R_j is the effect of replication, and E_{ijk} is the residual error. Multiple Factor Analysis (MFA) was performed using the FactoMineR package, treating quantitative variables as scaled continuous traits and qualitative variables as categorical traits. Phenotypic dissimilarity among accessions was estimated using Gower's distance, followed by hierarchical clustering with average linkage (UPGMA). The optimal number of clusters was identified using the goodness-of-split criterion implemented in the maptree package. To evaluate genotype–phenotype association between aroma class and *BADH2* genotypic category, a chi-square test of independence was performed using the contingency table of aroma score classes and marker genotypes.

Results and Discussion

Agro-morphological variation among accessions

The 183 traditional rice accessions exhibited substantial phenotypic diversity across the 22 agro-morphological traits evaluated. Qualitative traits showed wide variability, reflecting the heterogeneous evolutionary and cultural histories of these landraces. Traits such as leaf blade pubescence, leaf blade colour, and basal sheath colour demonstrated multiple expression states. Leaf blade colour was predominantly green (71%)

followed by dark green (23%), whereas basal leaf sheath colour ranged from green (85%) to purple-tinged phenotypes, suggesting underlying variation in anthocyanin pigmentation pathways. Leaf angle (LA) also showed even distribution, including erect, horizontal, and droopy forms, suggesting variation in plant architecture possibly linked to environmental adaptation. Other pigmentation traits, such as ligule colour (LC), collar colour (CC), and auricle colour (AC), showed limited variation, with most accessions being light green.

Furthermore, seed coat colour (SCC) and aroma are key grain quality parameters with both consumer and breeding relevance (Zhou *et al.*, 2020). While white pericarp was predominant (70%), the presence of red, brown, and light brown pericarps in a significant proportion of accessions indicates nutritional and phytonutrient diversity, which could be leveraged for biofortification efforts (Rathna Priya *et al.*, 2019). Aroma assessment using the KOH method revealed that, among the 183 accessions, 42% were strongly aromatic, 36% were mildly aromatic, and 23% were non-aromatic. Strongly aromatic landraces included several well-known traditional cultivars such as 'Gobindabhog', 'Katarni', 'Sonachur', 'Samjeera', 'Kalajeera', 'Kalanamak', and 'Tulaipanji'. Notably, many lesser-known cultivars, including 'Nanhia', 'Kusumphul', 'Prasad Bhog', 'Tushi Manjar', and 'Jeeraphul', also exhibited strong aroma. This highlights the underexplored aromatic potential within eastern Indian rice germplasm, particularly from Jharkhand (Table 1).

Table 1: List of highly aromatic landraces from the state of Jharkhand

Sl.No.	Landrace	Origin	BADH2 exon 7, 8-bp polymorphisms	Kernel shape
1	Gopalbhog (IC0385940)	Jharkhand	(580 bp & 257 bp)	1-Slender
2	Sonachur (IC0282478)	Jharkhand	(580 bp & 257 bp)	3-Medium
3	Kusum phool (IC0282466)	Jharkhand	(580 bp & 257 bp)	5-Bold
4	Dhania phool (IC0282519)	Jharkhand	(580 bp & 257 bp)	3-Medium
5	Sonachur (IC0282476)	Jharkhand	(580 bp & 257 bp)	1-Slender
6	Tulsi Manjar (IC0342874)	Jharkhand	(580 bp & 257 bp)	3-Medium
7	Kalajeera (IC0203154)	Bihar	(580 bp & 257 bp)	1-Slender
8	Jeeraphul (IC0267977)	Chattisgarh	(580 bp & 257 bp)	3-Medium
9	Badsha bhog (IC0299955)	Chattisgarh	(580 bp & 257 bp)	1-Slender
10	Katarni (IC0385729)	Bihar	(580 bp & 257 bp)	1-Slender
11	Badshabhog (IC0470970)	West Bengal	(580 bp & 257 bp)	3-Medium
12	Adamchini (IC0553800)	Uttar Pradesh	(580 bp & 257 bp)	3-Medium
13	Shyam Jeera (IC0553879)	Uttar Pradesh	(580 bp & 257 bp)	1-Slender
14	Badsha Pasand (IC0554635)	Uttar Pradesh	(580 bp & 257 bp)	1-Slender
15	Kasturi (IC0574880)	Bihar	(580 bp & 257 bp)	3-Medium
16	Sonachur (IC0574927)	Bihar	(580 bp & 257 bp)	1-Slender

17	Sitabhog (IC0594012)	West Bengal	(580 bp & 257 bp)	3-Medium
18	Sabour Surbhit (IC0595579)	Bihar	(580 bp & 257 bp)	3-Medium
19	Katari bhog (IC0618972)	West Bengal	(580 bp & 257 bp)	3-Medium
20	Radhuni pagol (IC0619255)	West Bengal	(580 bp & 257 bp)	1-Slender
21	Kala jeera (IC0619256)	West Bengal	(580 bp & 257 bp)	3-Medium
22	Kola joha (IC0619302)	Assam	(580 bp & 257 bp)	3-Medium
23	Mohan mala (IC0099217)	Jharkhand	(580 bp & 355 bp)	1-Slender
24	Kohraphool (IC0276799)	Jharkhand	(580 bp & 355 bp)	1-Slender
25	Prasad bhog (IC0282462)	Jharkhand	(580 bp & 355 bp)	1-Slender
26	Nanhia dhan (IC0342866)	Jharkhand	(580 bp & 355 bp)	1-Slender
27	Basmati (IC0282465)	Jharkhand	(580 bp & 355 bp)	3-Medium
28	Prasad bhog (IC0282406)	Jharkhand	(580 bp & 355 bp)	3-Medium
29	Bansbuda (IC0621734)	Jharkhand	(580 bp & 355 bp)	3-Medium
30	Prasad bhog (IC0621769)	Jharkhand	(580 bp & 355 bp)	1-Slender
31	Kamod (IC0396060)	Chattisgarh	(580 bp & 355 bp)	3-Medium
32	Samjeera (IC0332049)	Jharkhand	Heterozygous	1-Slender
33	Tulsi Manjar Dhan (IC0417979)	Jharkhand	Heterozygous	3-Medium
34	Lalmati (IC0640882)	Jharkhand	Heterozygous	3-Medium
35	Sakkar chini (IC0554637)	Uttar Pradesh	Heterozygous	3-Medium
36	Chini Karpoor (IC0554650)	Uttar Pradesh	Heterozygous	3-Medium

Grain-related qualitative traits were considerably diverse (Fig. 2). The apiculus colour was mostly straw (76%) but included brown and black variants, while lemma–palea colour exhibited a wide spectrum from gold and straw to rare reddish or purple forms. Such pigmentation variability aligns with findings in other aromatic landraces, highlighting the polygenic basis of grain colour traits (Ray *et al.*, 2013; Behera and Panda, 2023). Most accessions lacked awns (84%), although several carried long coloured awns, indicating selective retention of primitive traits in certain regional landraces. Pericarp colour was predominantly white (70%), with remaining accessions showing red or brown pericarps, traits associated with nutritional and antioxidant value (Rathna Priya *et al.*, 2019). This diversity is consistent with reports from Bangladesh and eastern India (Islam *et al.*, 2018).



Fig. 2: Grain morphological variation in representative rice accessions.

Quantitative grain traits also displayed considerable variation. Grain length ranged from 5.4–11.0 mm (CV = 16%), with the longest grains recorded in 'Kendu Manja' (IC0438534), 'Kasturi' (IC0438655), 'Kadam Phul' (IC0086591) and 'Vishun bhog' (IC0300024). Grain width showed moderate variation (CV = 15%), while kernel length ranged from 3.9–8.6 mm (CV = 17%). Kernel shape was dominated by medium types (50%), followed by slender (44%) and bold (5%). The widest variation was observed for 100-seed weight (CV = 33%), which ranged from 0.7 g ('Tulsi Manjar' IC0342874) to 3.2 g ('Kala bhuri' IC0421349), reflecting broad differences in endosperm development. These ranges corroborate earlier studies in Indo-Gangetic aromatic rice (Bairwa *et al.*, 2024).

Multivariate analysis revealed weak geographical structuring

To examine the overall pattern of phenotypic variation, Multiple Factor Analysis (MFA) integrating qualitative and quantitative traits was performed. The first five dimensions explained 26% of the total phenotypic

variation. Although this proportion appears modest, it is typical for mixed-data analyses involving a large number of categorical descriptors, where variance is distributed across multiple dimensions rather than concentrated in the leading axes. In the present study, qualitative traits contributed strongly to Dimensions 1–4, while quantitative grain traits such as grain length, grain weight, and kernel length were major contributors to Dimensions 1 and 2, indicating that grain morphology is a primary driver of phenotypic differentiation (Fig. 3A), consistent with earlier studies in aromatic rice landraces (Ray *et al.*, 2013; Roy *et al.*, 2016; Chandraker *et al.*, 2024).

Scatterplots of Dimensions 1 vs. 2 and 3 vs. 4 revealed weak clustering by state, with substantial overlap among accessions (Fig. 3B). This diffuse structure suggests that phenotypic variation is not geographically partitioned, likely reflecting historical seed exchange, cultural diffusion, and farmer-mediated selection across regions. However, future studies involving molecular markers will clarify these aspects. Similar patterns of weak geographic structuring have




Fig. 3: Multivariate analysis of 183 rice accessions.

(A) Percent contributions of qualitative and quantitative traits in MFA (B) Scatterplot depicting the spatial distribution of the accessions (C) Dendrogram showing grouping of rice accessions in seven clusters.

been reported in aromatic rice germplasm from eastern India and neighbouring regions (Prasad *et al.*, 2020).

Hierarchical clustering based on Gower's distance revealed pairwise dissimilarities ranging from 0.04 to 0.76. The Mantel correlation between the dissimilarity matrix and the dendrogram was significant ($r = 0.62$; $P = 0.001$), indicating a good representation of the underlying phenotypic relationships. Based on the goodness-of-split criterion, seven clusters were identified (Fig. 3C). However, the clustering pattern was highly uneven, with Cluster 3 comprising the majority of accessions ($n = 143$), while the remaining clusters contained relatively few genotypes. This pattern indicates that most accessions share a common morphological core defined by similar combinations of grain and vegetative traits, whereas a smaller subset represents phenotypic extremes or unique trait combinations. Clusters 1, 5, and 6 captured such outlier groups characterized by distinct grain shapes or pigmentation traits. Notably, Cluster 5 contained a single accession ('Tulsi Bhog'), highlighting its exceptional phenotypic distinctiveness within the panel. Clusters 2 and 4 exhibited higher values for grain length and brown rice length, whereas Cluster 6 showed the lowest values. Clusters 3 and 4 had comparatively lower median grain width and grain

weight. Previously, six clusters within aromatic rice accessions from West Bengal have been reported based on agro-morphology (Mondal *et al.*, 2024).

From a breeding perspective, the minor clusters (Clusters 1, 5, and 6) represent valuable sources of unique phenotypic variation, while the large Cluster 3 reflects a broadly adapted and morphologically conserved group. The absence of clear state-wise aggregation further indicates that local nomenclature does not reliably correspond to phenotypic distinctiveness, emphasizing the need for systematic characterization in germplasm management and utilization. However, a comprehensive molecular analysis will be required to fully resolve the underlying population structure and genetic relationships among these accessions.

Allelic variation at the *BADH2* exon 7 polymorphism and its association with aroma

Genotyping of the 8-bp deletion in exon 7 of the *BADH2* gene revealed three genotype classes among the 183 accessions: homozygous non-aromatic allele (580 + 355 bp; 101 accessions, 55%), homozygous aromatic allele (580 + 257 bp; 57 accessions, 31%), and heterozygous (580 + 355 + 257 bp; 25 accessions, 14%) (Fig. 4A). Integration of genotypic data with sensory aroma




Fig. 4: Genotyping for aroma-associated 8bp-deletion in exon 7 of *BADH2* gene.

(A) Representative gel photograph showing the allelic variations, (B) Polymorphisms detected in 183 accessions, (C) Scatter plot depicting distribution of rice accessions based on variation in 22 phenotypic traits.

classification demonstrated a strong but incomplete association between the *BADH2* exon 7 deletion and aroma phenotype (Fig. 4B). A chi-square test of independence confirmed a highly significant association between genotype and aroma class ($\chi^2 = 56.77$, $df = 4$, $P < 0.001$), indicating that variation at this locus is a major determinant of fragrance expression in the studied germplasm. Accessions carrying the homozygous aromatic allele were predominantly strongly aromatic, whereas those with the homozygous non-aromatic allele were largely non-aromatic or weakly aromatic. Heterozygous accessions exhibited intermediate distributions across aroma classes.

Despite this strong association, notable discordance between genotype and phenotype was observed. Among strongly aromatic accessions, only 56% carried the homozygous aromatic allele, while approximately 43% of strongly or mildly aromatic accessions lacked the exon 7 deletion. This discrepancy may partly reflect environmental influences on aroma expression, as 2-acetyl-1-pyrroline (2-AP) accumulation is affected by temperature, moisture, and nutrient status during crop growth (Guo et al., 2026). As the study was conducted under a single environment, genotype $\times$ environment interactions could not be assessed. Conversely, a small proportion of accessions carrying the aromatic allele exhibited weak or no aroma. Together, these findings indicate that the 8-bp deletion alone does not fully explain aroma variation in this germplasm, consistent with earlier reports (Fitzgerald et al., 2008; Shi et al., 2014).

Previous studies have reported multiple non-functional *BADH2* alleles associated with fragrance, including mutations in exon 10 (*badh2.7*) and exon 14 (*badh2.6*), particularly in *indica* rice from South and Southeast Asia (Kovach et al., 2009; Chandanshive et al., 2024). In the present study, genotyping was done for the exon 7 8-bp deletion (*badh2.1*), as it represents the most widely reported and diagnostically robust allele associated with fragrance across diverse rice germplasm (Bradbury et al., 2005b; Kovach et al., 2009). The allele-specific PCR assay for *BADH2* exon 7 8-bp deletion is simple, easy to score reliably and suitable for screening large germplasm sets. However, this approach does not capture other known functional variants such as *badh2.6* and *badh2.7*. Therefore, the presence of additional fragrance-associated alleles in accessions lacking the exon 7 deletion could not be resolved in this study and warrants further investigation through targeted sequencing or haplotype-based analyses.

Mapping *BADH2* genotypes onto MFA plots

(Fig. 4C) revealed moderate alignment between aromatic allele carriers and specific phenotypic clusters, suggesting that the exon 7 deletion contributes to phenotypic differentiation but does not fully define overall morphological structure. This partial correspondence reinforces the view that aroma, although influenced by *BADH2*, is only one component of the broader phenotypic diversity observed among traditional aromatic rice landraces.

Overall, this study revealed substantial agro-morphological and fragrance-related diversity among 183 traditional rice cultivars from eastern India, with 42% classified as highly aromatic, 36% as moderately aromatic, and 23% as non-aromatic based on the KOH test. Key grain traits, particularly grain length, brown rice length, and grain width, were major contributors to phenotypic differentiation across the panel. Genotyping indicated that accessions carrying the *BADH2* exon 7, 8-bp deletion were generally associated with stronger aroma expression; however, a considerable proportion of aromatic accessions lacked this allele, suggesting that additional genetic factors may contribute to fragrance in these landraces. These findings highlight the complexity of the genetic basis of aroma in eastern Indian rice germplasm and underscore the importance of integrating phenotypic and molecular approaches for its characterization. The results provide a useful baseline for future research aimed at elucidating the genetic basis of aroma, as well as for the conservation and protection of unique aromatic landraces in the region.

Acknowledgements

This work was supported by the Rashtriya Krishi Vikash Yojana (RKVY), Government of Jharkhand, under the project "Mainstreaming traditional aromatic quality rice varieties of Jharkhand: Evaluation, conservation and popularization through Farmer Producer Organizations" (F. No. 04/RKVY-13/2023-34), implemented at ICAR-CRRI, CRURRS, Hazaribag. The authors gratefully acknowledge ICAR-NBPGR, New Delhi, for providing rice germplasm seeds under a Material Transfer Agreement (MTA). We also thank the technical staff and field staff members at CRURRS for their valuable assistance in field and laboratory activities.

References

- Behera PK and D Panda (2023) Germplasm resources, genes and perspective for aromatic rice. *Rice Sci.* 30(4): 294-305.
- Bisne R and AK Sarawgi (2008) Agro-morphological and quality characterization of Badshah Bhog group from aromatic rice germplasm of Chhattisgarh. *Bangladesh J. Agric. Res.* 33(3):

- 479-492.
- Bradbury LM, RJ Henry, Q Jin, RF Reinke and DL Waters (2005b) A perfect marker for fragrance genotyping in rice. *Mol. Breed.* 16(4): 279-283.
- Bradbury LM, TL Fitzgerald, RJ Henry, Q Jin and DL Waters (2005a) The gene for fragrance in rice. *Plant Biotechnol. J.* 3(3): 363-370.
- Chandanshive S, S Mathure and A Nadaf (2024) Multiple mutations in BADH2 gene reveal the novel fragrance allele in indica rice (*Oryza sativa* L.). *J. Genet.* 103(1): 13.
- Fitzgerald TL, DLE Waters and RJ Henry (2008) The effect of salt on betaine aldehyde dehydrogenase transcript levels and 2-acetyl-1-pyrroline concentration in fragrant and non-fragrant rice (*Oryza sativa*). *Plant Sci.* 175(4): 539-546.
- Guo J, J Miao, J Chen, D Huang, C Wang and J Wen (2026) Environmental regulation of 2-Acetyl-1-pyrroline biosynthesis in fragrant rice: From metabolic pathways to sustainable quality management. *Genes* (Basel) 17(3): 349.
- IRRI (2013) *Standard Evaluation System (SES) for Rice*, 5th edn. International Rice Research Institute, Manila, 57 p.
- Kovach MJ, MN Calingacion, MA Fitzgerald and SR McCouch (2009) The origin and evolution of fragrance in rice (*Oryza sativa* L.). *Proc. Natl. Acad. Sci. USA* 106(34): 14444-14449.
- Mondal S, P Pradhan, B Das, D Kumar, B Paramanik, R Yonzon, R Barman, Saha D, J Karforma, A Basak, P Dey and MF Seleiman (2024) Genetic characterization and diversity analysis of indigenous aromatic rice. *Heliyon* 10(10): e31232.
- Nagaraju M, KK Mohanty, D Chaudhary and CA Gangadharan (1991) Simple technique to detect scent in rice. *Oryza* 28: 109-110.
- Prodhan ZH and SHU Qingyao (2020) Rice aroma; A natural gift comes with price and the way forward. *Rice Sci.* 27(2): 86-100.
- Rathna Priya TS, ARL Eliazar Nelson, K Ravichandran and U Antony (2019) Nutritional and functional properties of coloured rice varieties of South India; a review. *J. Ethnic Foods* 6(1): 1-11.
- Rathna Priya TS, ARL Eliazar Nelson, K Ravichandran and U Antony (2019) Nutritional and functional properties of coloured rice varieties of South India: A review. *J. Ethnic Foods* 6(1): 1-11.
- Ray A, D Deb, R Ray and B Chattopadhyay (2013) Phenotypic characters of rice landraces reveal independent lineages of short-grain aromatic indica rice. *AOB Plants* 5: plt032.
- Roy PS, R Samal, GJN Rao, SSC Patnaik, NN Jambhulkar, A Patnaik and T Mohapatra (2016) Differentiation and description of aromatic short grain rice landraces of eastern Indian state of Odisha based on qualitative phenotypic descriptors. *BMC Ecol.* 16(1): 36.
- Roy S, A Banerjee, B Mawkhlieng, AK Misra, A Pattanayak, GD Harishand KC Bansal (2015) Genetic diversity and population structure in aromatic and quality rice (*Oryza sativa* L.) landraces from North-Eastern India. *PLoS One* 10(6): e0129607.
- Roy S, A Banerjee, N Basak, J Kumar and NP Mandal (2020) Aromatic rice. In: *The Future of Rice Demand: Quality Beyond Productivity*, pp 251-282.
- Sakthivel K, RM Sundaram, NS Rani, SM Balachandran and CN Neeraja (2009) Genetic and molecular basis of fragrance in rice. *Biotechnol. Adv.* 27(4): 468-473.
- Shi Y, G Zhao, X Xu and J Li (2014) Discovery of a new fragrance allele and development of functional markers for identifying diverse fragrant genotypes in rice. *Mol. Breed.* 33: 701-708.
- Sinha H (2016) Rediscovering the traditional paddy varieties in Jharkhand: Conservation priority in hybrid rice era. *J. Rural Develop.* 35(2): 285-307.
- Zhou H, D Xia and Y He (2020) Rice grain quality-traditional traits for high quality rice and health-plus substances. *Mol. Breed.* 40: 1

RESEARCH ARTICLE

Assessment of Molecular Diversity and Phylogenetic Analysis Among Chilli Genotypes Using Start Codon Targeted (SCoT) Markers

Hanuman Ram¹, DK Samadia¹, Chet Ram^{1*} and AK Verma¹

Abstract

Assessment of genetic diversity is pivotal to identify genotypes for the crop improvement programme in chilli. Therefore, the present study was carried out to identify and characterise genetic variation among 45 chilli genotypes using Start Codon Targeted Polymorphism (SCoT) markers. Out of 18 primers, ten markers revealed significant polymorphism among the chilli accessions studied, with amplification ranging from 280 bp to 2.0 kb. A total of 81 alleles were obtained, ranging from 5 to 10 alleles, with an average of 8.1 alleles per primer. Out of 81 alleles, sixty-two (76.54%) were found polymorphic in nature. The genetic similarity among the analysed genotypes ranged from 24.2 to 100%. The UPGMA dendrogram was constructed to establish the genetic relationship among the chilli genotypes using Jaccard's coefficient. Cluster analyses grouped 45 genotypes into two major groups, which was also supported by principal coordinate analysis (PCoA). Analysis of molecular variance (AMOVA) showed a significant estimated value at a degree of 999 permutations. The percentage of variability was higher among populations (55%) than within populations (45%). The mean Nei's gene diversity (*h*) was 0.2233 and the mean Shannon's Information Index (*i*) was 0.3470, with the highest of both given by the SCoT-7 marker. The SCoT marker-based distinguishable genetic variation in the chilli genotypes may serve as a potential aid in chilli improvement programme. It demonstrates the utility of SCoT markers for estimating molecular diversity and genotype identification in chilli. This study also suggests the potential of SCoT markers in further association studies in chilli.

Keywords: Chilli, Diversity, PCoA, Polymorphism, SCoT markers

¹ICAR-Central Institute for Arid Horticulture, Bikaner, Rajasthan, 334006, India.

***Author for correspondence:**
chet.ram2@icar.org.in

Received: 24/12/2025

Revised: 19/03/2026

Accepted: 13/04/2026

How to cite this article: Ram H, DK Samadia, C Ram and AK Verma (2026) Assessment of Molecular Diversity and Phylogenetic Analysis Among Chilli Genotypes Using Start Codon Targeted (SCoT) Markers. *Indian J. Plant Genet. Resour.* 39(2): 29-37.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.174522>

Introduction

Chilli, scientifically categorized under the genus *Capsicum*, is one of the important members of the Solanaceae family. Native to Central and South America, chilli was domesticated more than 6,000 years ago and is now used extensively in industry, medicine, and cooking all over the world (Bosland and Votava, 2012). Among *Capsicum* spp., *Capsicum annum*, *C. chinense*, *C. frutescens*, *C. baccatum*, and *C. pubescens* are the main domesticated species (Pickersgill, 2007). Chillies are also nutritionally dense, containing high levels of vitamin C, minerals, amino acids, provitamin A (β -carotene), flavonoids, and phytochemicals, making them useful for pharmacological and dietary applications (Materska and Perucka, 2005; Olatunji and Afolayan, 2018; Palma-Orozco *et al.*, 2021). Capsaicinoids have been studied for their involvement in pain relief, metabolism stimulation, and cardiovascular protection in addition to their antioxidant and antimicrobial activities (Reyes-Escogido *et al.*, 2011). Due to their agronomic

importance, biological diversity, and therapeutic potential, chilli continues to be a focal point of research in plant science, nutrition, and pharmacology. Chillies are still focused on by plant science, pharmacology, and nutritional studies due to their genetic diversity, therapeutic potential, and agronomic importance.

Plant adaptation, resilience, and evolution are significantly influenced by genetic diversity. Genetic diversity at the molecular level includes differences across and within plant populations in terms of DNA sequences, gene architectures, allelic frequencies, and genome organization (Frankham *et al.*, 2002). Phenotypic traits diversity, including disease resistance, drought tolerance, yield, and quality are based on these molecular differences and is essential for crop improvement and breeding. In addition to managing genetic resources and identifying unique genotypes, an understanding of molecular genetic diversity is necessary for mapping genes associated with desired agronomic qualities using marker-assisted selection (MAS) and association mapping (Gupta *et al.*, 2001).

The researchers have characterized several vegetable germplasm, including chilli, using morphological markers (Rahevar *et al.*, 2021; Samadia *et al.*, 2024). However, the genetic diversity inferred by morphological markers is not precise due to the influence of environmental factors. Thus, molecular markers can provide accurate diversity at the molecular level. Maintaining and leveraging genetic variability at the molecular level is pivotal for sustainable agriculture and food security in the face of habitat loss and climate change. The evaluation of genetic diversity in plants has been completely transfigured by the advancement of molecular markers. Numerous methods have been executed to measure and examine genetic variability, including RAPD (Random Amplified Polymorphic DNA), AFLP (Amplified Fragment Length Polymorphism), SSR (Simple Sequence Repeats), and SNP (Single Nucleotide Polymorphisms) (Powell *et al.*, 1996; Semagn *et al.*, 2006). In continuation, the short, conserved areas that surround the ATG translation initiation codon in plant genes serve as the basis for the new class of gene-targeted molecular markers known as Start Codon

Targeted (SCoT) markers (Collard and Mackill, 2009). The functional nature of SCoT markers, which target conserved gene areas, makes them more trustworthy for identifying functional genetic variation related to the trait of interest than other anonymous markers like RAPD or AFLP. They are cost-effective and adaptable to a variety of plant species; they are useful in genetic diversity analysis, DNA fingerprinting, population structure assessment, phylogenetic studies, and QTL mapping; they are highly reproducible because longer primers and higher annealing temperatures are needed; and they don't require prior sequence information; able to identify polymorphisms in coding areas, which are frequently linked to adaptability and expressed traits. These markers' broad applicability in plant breeding, conservation genetics, and cultivar identification has been demonstrated by their successful use in a range of crops, including rice (*Oryza sativa*), wheat (*Triticum aestivum*), grapes (*Vitis vinifera*), and potato (*Solanum* spp.) (Xiong *et al.*, 2011; Gorji *et al.*, 2011). However, the utilization of SCoT markers for assessing genetic diversity is limited in chilli (Tsaballa *et al.*, 2015; Gupta *et al.*, 2019; Igwe *et al.*, 2019). The existing research gap limits the utility of gene-targeted molecular diversity for enhancing chilli breeding programmes. Therefore, the current investigation was formulated with the aims to (1) harness the SCoT marker-based genetic potential in chilli, (2) identify the unique genotype and (3) phylogenetic relationship among the chilli genotypes. Thus, the study was performed to assess molecular diversity and phylogenetic relationship among 45 chilli genotypes by employing 10 SCoT markers. The study highlights the promising potential of SCoT markers for advancing chilli crop improvement programmes.

Materials and Methods

Plant materials

The experimental material for the present investigation comprised 45 chilli genotypes, including four adapted varieties (JCA-283, Pusa Jwala, Pusa Sadabahar and RCH-1) (Table 1). The seeds of each genotype were sown

Table 1: List of genotypes used for molecular diversity and phylogenetic analysis

S. No.	Accession/ collector number	IC number	Source	S. No.	Accession/ collector number	IC number	Source
1.	AHM-05	IC-347001	ICAR-CIAH, Bikaner	6.	AHM-93	IC-347089	ICAR-CIAH, Bikaner
2.	AHM-31	IC-347027	ICAR-CIAH, Bikaner	7.	AHM-115	IC-347111	ICAR-CIAH, Bikaner
3.	AHM-35	IC-347031	ICAR-CIAH, Bikaner	8.	Chilli thin-1	--	ICAR-CIAH, Bikaner
4.	AHM-42	IC-347038	ICAR-CIAH, Bikaner	9.	Chilli thin-2	--	ICAR-CIAH, Bikaner
5.	AHM-55	IC-347051	ICAR-CIAH, Bikaner	10.	Chilli thick-1	--	ICAR-CIAH, Bikaner

S. No.	Accession/ collector number	IC number	Source	S. No.	Accession/ collector number	IC number	Source
11.	Chilli-dwarf	--	ICAR-CIAH, Bikaner	29.	AHM-192	IC-446364	ICAR-CIAH, Bikaner
12.	KSB-9	IC-261862	ICAR-CIAH, Bikaner	30.	AHM-193	IC-446365	ICAR-CIAH, Bikaner
13.	AHM-154	IC-446326	ICAR-CIAH, Bikaner	31.	AHM-194	IC-446366	ICAR-CIAH, Bikaner
14.	AHM-162	IC-446334	ICAR-CIAH, Bikaner	32.	AHM-197	IC-446369	ICAR-CIAH, Bikaner
15.	AHM-164	IC-446336	ICAR-CIAH, Bikaner	33.	AHM-198	IC-446370	ICAR-CIAH, Bikaner
16.	AHM-165	IC-446337	ICAR-CIAH, Bikaner	34.	AHMH-01	--	ICAR-CIAH, Bikaner
17.	AHM-167	IC-446338	ICAR-CIAH, Bikaner	35.	MC-2	--	ICAR-CIAH, Bikaner
18.	AHM-172	IC-446344	ICAR-CIAH, Bikaner	36.	MC-1	--	ICAR-CIAH, Bikaner
19.	AHM-175	IC-446347	ICAR-CIAH, Bikaner	37.	JCA-283	--	JNKVV, Jabalpur
20.	AHM-177	IC-446349	ICAR-CIAH, Bikaner	38.	Pusa Jwala	--	ICAR-IARI, New Delhi
21.	AHM-179	IC-446351	ICAR-CIAH, Bikaner	39.	Pusa Sadabahar	--	ICAR-IARI, New Delhi
22.	AHM-180	IC-446352	ICAR-CIAH, Bikaner	40.	RCH-1	--	AU, Jodhpur
23.	AHM-181	IC-446353	ICAR-CIAH, Bikaner	41.	AHMH-02	--	ICAR-CIAH, Bikaner
24.	AHM-182	IC-446354	ICAR-CIAH, Bikaner	42.	AHMH-03	--	ICAR-CIAH, Bikaner
25.	AHM-183	IC-446355	ICAR-CIAH, Bikaner	43.	AHMH-04	--	ICAR-CIAH, Bikaner
26.	AHM-184	IC-446356	ICAR-CIAH, Bikaner	44.	AHMH-05	--	ICAR-CIAH, Bikaner
27.	AHM-189	IC-446361	ICAR-CIAH, Bikaner	45.	AHMH-06	--	ICAR-CIAH, Bikaner
28.	AHM-191	IC-446363	ICAR-CIAH, Bikaner				

in a protective vegetable nursery bed (Ram *et al.*, 2024) at the vegetable research field of ICAR-Central Institute for Arid Horticulture, Bikaner, Rajasthan, India (28.1115° N and 73.3497° E). The seedlings were watered thrice a week with a drip irrigation system. The healthy and disease-free leaf samples were collected from nursery-staged seedlings (28-days-old). The leaf samples were wrapped in the aluminium foil and immediately frozen in liquid nitrogen (-196°C) and further stored at a deep freezer (-80°C) for DNA isolation.

DNA isolation and SCoT marker profiling

Molecular work was undertaken at the Molecular unit, Biotechnology laboratory, ICAR-CIAH, Bikaner. The genomic DNA was isolated from 100 mg of fresh young leaf tissue following the CTAB method in the presence of liquid nitrogen with some minor modifications (Murray and Thompson, 1980). DNA was precipitated with chilled iso-propanol, and subsequently, the DNA pellet was rinsed with 70% ethanol for 10-15 min. to remove excess amount of CTAB and other salt precipitates. The pellet was dried at room temperature for up to 1 h and dissolved in TE buffer (pH=8). The purified DNA was quantified electrophoretically on 0.8% agarose gel. The genomic DNA was diluted to 10 ng/μl as a working stock for PCR analysis. PCR reaction was carried out using a Thermal Cycler machine (Bio-Rad Laboratories Inc., USA) in a 15 μl reaction volume, composed of 2.0 μl of 10 ng genomic DNA, 7.5 μl of 2X PCR master mixture (Takara Bio Inc., Japan), 0.5 μl MgCl₂, 2.0 μl SCoT primer and 3.0 μl nuclease-free water (Hi-

media, India). PCR tubes containing the reaction components were gently tapped and given a brief spin to allow proper settling of the reaction mixture. PCR amplification was performed with initial denaturation at 94°C for 3 min. followed by 40 cycles of denaturation at 94°C for 30 s, annealing step at 45°C for 30 s, primer extension at 72°C for 2 min. and a final extension at 72°C for 10 min.

Gel electrophoresis and allele scoring

The amplicons were resolved electrophoretically on 1.5% agarose gel prepared in 1XTBE buffer. The DNA fragments (SCoT alleles) were visualized by staining with ethidium bromide (10 mg/ml) and finally documented using a gel documentation system (Syngene, USA). The amplicons were scored as alleles for each of the SCoT marker loci. The alleles were scored manually as a binary matrix (0 and 1, where 0 and 1 are considered as absence and presence of the alleles, respectively). The allele sizes (base pairs) were determined by comparing with a 100 bp standard DNA ladder (Bio-Helix, Taiwan). The allele data (binary matrix) from all the markers were used for statistical analysis.

Clustering and population structure analysis

The binary matrix data were further used for clustering analysis among the genotypes. The clustering was inferred using Jaccard's coefficient, and a tree was obtained using the UPGMA model in NTSys 2.0 software (Rohlf, 2000). To determine the number of population structures present among the 45 chilli genotypes, the

binary data were analysed in STRUCTURE 2.0 software with the admixture model (Pritchard *et al.*, 2000).

Statistical analysis

Total number of alleles, major allele frequency, gene diversity, heterozygosity and polymorphic information content (PIC) were estimated using PowerMarker V3.25 (Liu and Muse, 2005). Any allele appearing in only one genotype was considered a unique allele, whereas an allele with a frequency of <0.05 was considered a rare allele. Genetic dissimilarity was calculated for pairwise comparison of genotypes using Jaccard's coefficient with 1000 bootstraps. Principal coordinate analysis (PCoA) was calculated to supplement the clustering pattern (Gower, 1966). SCoT allelic composition for each genotype at every marker locus was determined by counting the number of alleles per locus and the allele frequencies and polymorphism information content (PIC) were determined according to the formula given by Botstein *et al.*, (1980).

$$PIC = \left(1 - \sum_{i=1}^k \hat{p}_i^2\right) \frac{2n}{2n-1} - \sum_{i=1}^{k-1} \sum_{j=i+1}^k 2\hat{p}_i^2 \hat{p}_j^2$$

Where p_i was the estimated allele frequencies of k alleles ($i=1$ to k) and n was the number of individuals. Analysis of Molecular Variance (AMOVA) was carried out using GenAlEx 6.5 (Peakall and Smouse, 2012). In addition, genetic indices such as observed number of alleles (n_a), number of effective alleles (n_e), Shannon's information index (i), Nei's observed heterozygosity (h) and expected heterozygosity (H_e) were also calculated using GenAlEx v6.503 (Peakall and Smouse, 2012).

Results and Discussion

SCoT allele characteristic and polymorphism

To assess the SCoT marker-based polymorphism among the chilli genotypes, eighteen SCoT markers were




Fig. 1: Representative gel image of SCoT7 amplification on genomic DNA of chilli genotypes.

The M and numerical 1-45 indicate the 100 bp standard DNA ladder and chilli genotypes, respectively.

profiled on the genomic DNA of chilli genotypes. Consequently, ten SCoT markers were amplified polymorphic alleles among the genotypes (Fig. 1) and used for the genetic diversity assessment among chilli genotypes. Eight SCoT markers were excluded from the analysis due to their monomorphic nature of amplification. A total of 81 alleles were obtained, ranging from 5 to 10 alleles, with an average of 8.1 alleles per locus. Out of 81 alleles, 62 (76.54%) were found polymorphic in nature. The allele size ranged from 280 bp to 2.0 kb, thereby revealing a significant level of genetic diversity among the chilli genotypes. Similarly, the genetic criteria pertain to efficiency of the markers, such as percentage of polymorphism, PIC, H_e and resolution power, which also showed significance values among the chilli genotypes. The moderate level of PIC value ranged from 0.25 to 0.58, with a mean value of 0.40 generated by the SCoT markers. Likewise, the H_e value ranged from 0.30 to 0.66 with a mean of 0.46. Additionally, SCoT markers produced resolution power ranging from 1.82 to 4.58 among chilli genotypes. On the contrary, the four SCoT markers, namely SCoT2, SCoT3, SCoT10 and SCoT11, were unable to produce resolutions among the alleles. Statistics of the polymorphism are presented in Table 2. The SCoT

Table 2: Polymorphism and marker efficiency of analysed SCoT markers in chilli

Marker	Sequence (5' – 3')	Amplicon size (bp)	No. of alleles	%Polymorphism	H_e	PIC value	Rp
SCoT2	CAACAATGGCTACCACCC	300-1100	10	70.00	0.49	0.44	--
SCoT3	CAACAATGGCTACCACCG	500-1600	10	90.00	0.62	0.54	--
SCoT4	CAACAATGGCTACCACCT	450-1000	5	60.00	0.30	0.25	1.82
SCoT5	CAACAATGGCTACCACGA	500-1000	6	83.33	0.49	0.37	2.76
SCoT6	CAACAATGGCTACCACGC	400-1000	8	62.50	0.47	0.36	3.64
SCoT7	CAACAATGGCTACCACGG	380-2000	10	90.00	0.42	0.33	3.91
SCoT8	CAACAATGGCTACCACGT	300-1350	9	88.89	0.47	0.36	4.58
SCoT9	CAACAATGGCTACCAGCA	280-870	7	85.7	10.44	0.34	2.53
SCoT10	CAACAATGGCTACCAGCC	550-1200	10	0.00	0.66	0.58	--
SCoT11	AAGCAATGGCTACCACCA	400-750	7	71.43	0.55	0.46	--
Mean	---	---	8.1	60.00	0.49	0.40	2.41

markers, as new gene-targeted marker systems, have been substantially utilised for characterizing germplasm of numerous crop plants, including solanaceous crops (Shahlaei *et al.*, 2014; Gupta *et al.*, 2019; Kulyan *et al.*, 2023; Altaf *et al.*, 2025). The SCoT marker has been considered as an effective and reproducible marker and resulted in high polymorphism correlated with traits of biological interest as compared to other random marker systems (Collard and Mackill, 2009; Shahlaei *et al.*, 2014). The SCoT marker analysed in the present study generated a substantial level of polymorphism among the chilli genotypes, indicating a wide level of genomic divergence among the accessions. Out of eighteen SCoT primers screened, ten primers were found polymorphic, while eight primers were excluded from the analysis due to their monomorphic allele pattern. The SCoT alleles are in accordance with the inherent nature of gene-targeted SCoT marker systems, where amplification efficiency varies according to primer–genome specificity (Collard and Mackill, 2009). The total of 81 alleles obtained with an average of 8.1 alleles per locus and 76.54% polymorphism indicated a higher resolution capacity of SCoT markers in chilli, demonstrating their power in revealing genomic variability in chilli in the present study. Similar high allele richness and polymorphism using SCoT markers have been reported earlier in *Capsicum* spp. (Gupta *et al.*, 2019; Igwe *et al.*, 2019; Abou-Sreea *et al.*, 2021). The moderate PIC values (0.25–0.58) and *He* values (0.30–0.66) obtained in this experiment further confirm the efficiency and informativeness of SCoT markers as powerful molecular tools for genetic diversity assessment among chilli genotypes. The findings of the present investigation were consistent with previous findings in *Capsicum* and other Solanaceae crops (Feng

et al., 2018; Gupta *et al.*, 2019; Igwe *et al.*, 2019; Shilpa *et al.*, 2021). The resolution power (*Rp*), ranging from 1.82 to 4.58, additionally indicated that these primers possess strong discriminatory potential in differentiating chilli genotypes at the molecular level. However, some markers (SCoT2, SCoT3, SCoT10 and SCoT11) have not produced allelic resolutions, which might be due to the low level of polymorphism in the present study. The similar discriminatory potential of SCoT markers in terms of resolution power has been observed in various crops (Bhawna *et al.*, 2017; Igwe *et al.*, 2017; Rai, 2023).

Genetic diversity analysis among chilli genotypes

Based on the genetic diversity indices such as observed number of alleles (*na*), number of effective alleles (*ne*), Shannon's information index (*i*) and Nei's observed heterozygosity (*h*), all the chilli genotypes showed a significant level of genetic diversity (Table 3). The observed and effective number of alleles generated by 10 SCoT markers in 45 chilli genotypes ranged from 1.40 (SCoT4) to 2.00 (SCoT5) and 1.23 (SCoT4) to 1.52 (SCoT8), with means of 1.75 and 1.36, respectively. The mean of Nei's observed heterozygosity was found to be 0.22, with ranging from 0.15 (SCoT4) to 0.30 (SCoT8). Similarly, Shannon's information index ranged from 0.22 (SCoT4) to 0.46 (SCoT8) with a mean of 0.34. The genetic diversity analysis identified the SCoT8 as a more informative marker to characterise 45 chilli genotypes. Whereas the SCoT4 markers have generated the least genetic diversity value in all the parameters. The number of alleles (*na* = 1.40 - 2.00; mean 1.75) and number of effective alleles (*ne* = 1.23 - 1.52; mean 1.36) align with earlier reports in *Capsicum* and tomato, where SCoT markers generally produce moderate to high allelic richness (Shahlaei *et al.*, 2014; Gupta *et al.*,

Table 3: SCoT markers based on allelic diversity in 45 chilli genotypes

Marker	<i>na</i>	<i>ne</i>	<i>h</i>	<i>i</i>
SCoT2	1.80	1.25	0.18	0.30
SCoT3	1.90	1.26	0.20	0.34
SCoT4	1.40	1.23	0.15	0.22
SCoT5	2.00	1.29	0.21	0.36
SCoT6	1.63	1.34	0.21	0.31
SCoT7	1.80	1.38	0.23	0.36
SCoT8	1.89	1.52	0.30	0.46
SCoT9	1.71	1.44	0.23	0.34
SCoT10	1.67	1.42	0.25	0.37
SCoT11	1.71	1.44	0.25	0.37
Mean	1.75	1.36	0.22	0.34

2019). Shannon's information index (i), ranging from 0.22 to 0.46 (mean 0.34), and Nei's heterozygosity ($h = 0.15\text{--}0.30$; mean 0.22), further confirm the presence of appreciable intra-specific variability. Such diversity estimates are comparable to those documented in chilli and other vegetable crops using SCoT and related co-dominant markers like SSRs (Gupta *et al.*, 2019; Ram *et al.*, 2021).

Structure-based diversity analysis

To determine the genetic populations present in 45 chilli genotypes, the binary matrix data were analysed in the STRUCTURE software. The delta K value revealed three population structures present in 45 chilli genotypes (Fig. 2). Population structure I consisted of 21 chilli genotypes, out of which 04 genotypes were found




Fig. 2: Eigen value (Delta K) indicating the number of populations among 45 chilli genotypes.

to be admixture type. Similarly, population cluster II contained 12 genotypes with 3 admixture types. The population structure III consisted of 08 highly porous, 3 admixture type and one genotype as a hybrid type (Fig. 3). The low level of admixture among the genotypes might be due to self-pollination nature of the crop. Population structure analysis identified three distinct genetic populations among the 45 chilli genotypes studied, supported by delta K analysis, which agrees with multiple population structure studies in chilli revealing 2–4 major genetic groups due to narrow reproductive behaviour and region-specific domestication (Taranto *et al.*, 2016). The comparatively




Fig. 3: Population structure analysis among 45 chilli genotypes.

lower proportion of admixture genotypes recorded might be attributed to the often-cross-pollinated nature of *Capsicum annuum*, thereby restricting intensive gene flow, as also suggested earlier by Pérez-Martínez *et al.*, (2022).

Clustering study

The UPGMA cluster analysis grouped 45 genotypes into three major clusters, some of them having sub-clusters (Fig. 4). Cluster one emerged as the largest cluster,




Fig. 4: UPGMA clustering analysis among 45 chilli genotypes.

possessing twenty-one genotypes with a 0.48 to 0.90 similarity index. Similarly, cluster 2 was found second largest cluster, having thirteen genotypes, including two varieties (JCA-283 and Pusa Sadabahar) with a 0.58 to 0.87 similarity index. Cluster 3 consisted of eleven chilli genotypes, including two varieties (Pusa Jwala and RCH-1). In cluster 3, five genotypes AHM-172, AHM-173, RCH-1, Pusa Jwala and AHMHR-3 were observed closest to each other in the phylogeny analysis, suggested the 100% similar based on the genetic level. The UPGMA-based clustering of genotypes also validated three primary genetic groups, which were further confirmed and supported by the STRUCTURE results. The clustering of commercial varieties such as Pusa Jwala, RCH-1, Pusa Sadabahar and their close grouping with specific accessions suggests shared allelic origin and similar genetic background, as also observed by Nagy *et al.*, (2007) and Dhaliwal *et al.*, (2014) in phylogenetic studies of chilli cultivars.

Molecular variability and coordination among chilli genotypes

An analysis of molecular variance (AMOVA) study was performed to analyse the distribution of genetic variations among or within populations of 45 chilli genotypes. Consequently, the analysis showed a

Table 4: AMOVA showing genetic variation among and within populations of 45 chilli genotypes

Source	df	SS	MS	Est. Var.	%
Among Pops	2	287.123	143.561	9.443	55%
Within Pops	4	2327.4	337.796	7.796	45%
Total	4	4614.556		17.239	100%

significant estimated value at a degree of 1000 permutations (Table 4). The results revealed that most of the variation (55%) was found among the populations rather than within populations (45%). Principal coordinate analysis (PCoA) was further performed to assess the cluster coordinates according to the genetic variability among 45 chilli genotypes. The findings of coordinate analysis revealed four clusters among 45 chilli genotypes with distribution in four quadrangles; however, the distribution of the genotypes across the quadrangles was not uniform. The maximum number of genotypes (21 genotypes) was presented by quadrangle 1, followed by quadrangle 4 (12 genotypes). Whereas, the quadrangle 2 presented the minimum number of genotypes clustered (4 genotypes). The coordinates 1 and 2 explained per cent variation of 54.56% and 19.94% with a cumulative per cent variation of 81.72% (Fig. 5). AMOVA indicated that maximum molecular variation resided among the populations rather than within populations, which clearly supports a structured genetic architecture in chilli. Similar trend of higher inter-population variance has been documented in chilli across different geographic origin-based studies (Yumnam *et al.*, 2012; Abhinaya *et al.*, 2016). The PCoA further supported cluster-based grouping of chilli genotypes, where differential spatial distribution of genotypes across quadrants indicated heterogeneous genetic divergence among the chilli genotypes studied. Thus, the combined marker efficiency statistics, clustering


pattern, population structure, and AMOVA outputs confirmed that SCoT markers exhibit high potential for precise discrimination, population stratification and genetic divergence in chilli genotypes.

In a nutshell, the findings of this study revealed a substantial level of allelic variation and polymorphism in chilli genotypes. Out of 81, ten SCoT markers were found informative to generate 81 alleles with 60.00% mean polymorphism. The moderate level of PIC value, ranging from 0.25 to 0.58, reflecting informativeness of SCoT markers analysed in this study. Three clusters corroborated with three population structures also infer the utility of the SCoT markers to identify the genotypes according to their phylogenetic proximity. Similarly, the AMOVA showed significant genetic variation among and within populations of 45 chilli genotypes. Overall, SCoT markers in this study were highly informative, revealing significant genetic diversity among chilli genotypes, which provided sufficient genetic distance to identify promising contrasting parents for future heterosis breeding, trait pyramiding and molecular breeding program development.

Conclusion

The present study confirmed that SCoT markers are highly efficient and reliable for assessing molecular polymorphism and genetic diversity in chilli. The high mean of per cent polymorphism (60%), moderate mean PIC values (0.40), superior resolution power (mean 2.41) and clear genetic clustering obtained in this study collectively indicated that SCoT markers have strong discriminatory strength for genome differentiation in *Capsicum annum*. The STRUCTURE, UPGMA and PCoA-based grouping consistently classified the germplasm into three genetically distinct populations, which suggests the existence of structured genetic divergence within the evaluated collection. AMOVA further validated this pattern by demonstrating that the majority of genetic variance in chilli lies among populations (55%) rather than within populations (45%), signifying clear differentiation and limited gene flow among the genotype groups.

From a breeding perspective, identification of genetically distant clusters and diverse genotypes is

**Fig.5:** Principal coordinate analysis (PCoA) of 45 chilli genotypes.

extremely valuable for parental selection in hybrid, heterosis and trait-specific breeding programmes. The genotypes belonging to genetically distant clusters can be strategically used to maximise heterotic response, broaden allelic spectrum, enhance recombination potential and pyramid desirable traits effectively in next-generation breeding. Therefore, the diversity detected through SCoT markers analysis in this study provides a valuable molecular basis and genetic resource framework for designing future chilli improvement, hybrid development and molecular breeding strategies.

References

- Abhinaya M, K Modha, RK Patel and HB Parmar (2016) Genetic diversity analysis for dry fruit yield, its attributes and quality traits in chilli (*Capsicum annuum* L.). *Electron. J. Plant Breed.* 7(4): 1200-1207.
- Abou-Sreea AIB, CR Azzam, S Al-Taweel, RM Abdel-Aziz, HEE Belal, MM Rady, EF Ali, A Majrashi and KAM Khaled (2021) Natural biostimulant attenuates salinity stress effects in chili pepper by remodelling antioxidant, ion, and phytohormone balances, and augments gene expression. *Plants* 10:2316.
- Altaf MT, MA Nadeem, A Ali, W Liaqat, M Bedir, N Baran, A Ilić, MK Ilyas, A Ghafoor, H Dogan, M Aasim and FS Baloch (2025) Applicability of Start Codon Targeted (SCoT) markers for the assessment of genetic diversity in bread wheat germplasm. *Genet. Resour. Crop Evol.* 72: 1205-1218. <https://doi.org/10.1007/s10722-024-02016-0>
- Bhawna, MZ Abidin, L Arya and M Verma (2017) Use of SCoT markers to assess the gene flow and population structure among two different population of bottle gourd. *Plant Genet.* 9: 80-86.
- Bird KA, H An, E Gazave, MA Gore, JC Pires and LD Robertson (2017) Population structure and phylogenetic relationships in a diverse panel of *Brassica rapa* L. *Front. Plant Sci.* 8: 321. doi:10.3389/fpls.2017.00321
- Bosland PW and EJ Votava (2012) *Peppers: Vegetable and Spice Capsicums*, 2nd edn. CABI Publishing.
- Botstein D, RL White, MH Skalnick and RW Davies (1980) Construction of a genetic linkage map in man using restriction fragment length polymorphism. *Am. J. Hum. Genet.* 32(3): 314-331.
- Collard BCY and DJ Mackill (2009) Start codon targeted (SCoT) polymorphism; a simple, novel DNA marker technique for generating gene-targeted markers in plants. *Plant Mol. Biol. Rep.* 27(1): 86-93. <https://doi.org/10.1007/s11105-008-0060-5>
- Dhaliwal MS, A Yadav and SK Jindal (2014) Molecular characterization and diversity analysis in chilli pepper using simple sequence repeats (SSR) markers. *Afr. J. Biotechnol.* 13(31): 3137-3143.
- Feng S, Y Zhu, CYu, K Jiao, M Jiang, J Lu, C Shen, Q Ying and H Wang (2018) Development of species-specific SCAR markers, based on a SCoT analysis, to authenticate *Physalis* (Solanaceae) species. *Front. Genet.* 9: 192. doi:10.3389/fgene.2018.00192
- Frankham R, JD Ballou and DA Briscoe (2002) *Introduction to Conservation Genetics*. Cambridge University Press.
- Gorji AM, P Poczar, Z Polgar and J Taller (2011) Efficiency of arbitrarily amplified dominant markers (SCoT, ISSR, and RAPD) for diagnostic fingerprinting in tetraploid potato. *Am. J. Potato Res.* 88(3): 226-237. <https://doi.org/10.1007/s12230-011-9197-7>
- Gower JC (1966) Some distance properties of latent root and vector methods used in multivariate analysis. *Biometrika* 53: 325-388.
- Gupta PK, RK Varshney, PC Sharma and B Ramesh (2001) Molecular markers and their applications in wheat breeding. *Plant Breed.* 120(3): 179-204.
- Gupta V, PK Jatav, SU Haq, KS Verma, VK Kaul, SL Kothari and S Kachhwaha (2019) Translation initiation codon (ATG) or SCoT markers-based polymorphism study within and across various *Capsicum* accessions; insight from their amplification, cross-transferability and genetic diversity. *J. Genet.* 98(2): 61.
- Igwe DO, CA Afiukwa, G Acquah and GN Ude (2019) Genetic diversity and structure of *Capsicum annuum* as revealed by start codon targeted and directed amplified minisatellite DNA markers. *Hereditas* 156: 32. <https://doi.org/10.1186/s41065-019-0108-6>
- Igwe DO, CA Afiukwa, BE Ubi, KI Ogbu, OB Ojuederie and GN Ude (2017) Assessment of genetic diversity in *Vigna unguiculata* L. (Walp) accessions using inter-simple sequence repeat (ISSR) and start codon targeted (SCoT) polymorphic markers. *BMC Genet.* 18: 98.
- Kulyan R, L Samarina, R Shkhalakhova, A Kuleshov, Y Ukhatoeva, O Antonova, N Koninskaya, A Matskiv, V Malyarovskaya and A Ryndin (2023) InDel and SCoT markers for genetic diversity analysis in a citrus collection from the western caucasus. *Int. J. Mol. Sci.* 24: 8276. <https://doi.org/10.3390/ijms24098276>
- Liu K and SV Muse (2005) PowerMarker: an integrated analysis environment for genetic marker analysis. *Bioinform.* 21(9): 2128-2129.
- Materska M and I Perucka (2005) Antioxidant activity of the main phenolic compounds isolated from hot pepper fruit (*Capsicum annuum* L.). *J. Agric. Food Chem.* 53(3): 1750-1756.
- Murray MG and WF Thompson (1980) Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Res.* 8(19): 4321-4325.
- Nagy I, A Stigel, Z Sasvari, M Roder and M Ganai (2007) Development, characterization, and transferability to other Solanaceae of microsatellite markers in pepper (*Capsicum annuum* L.). *Genome* 50: 668-688.
- Olatunji TL and AJ Afolayan (2018) The suitability of chili pepper (*Capsicum annuum* L.) for alleviating human micronutrient dietary deficiencies: A review. *Food Sci. Nutr.* 6: 2239-2251.
- Palma-Orozco G, C Orozco-Alvarez, AA Chavez-Villeda, A Mixtega-Martínez and R Castro-Munoz (2021) Capsaicin content in red habanero chilli (*Capsicum chinense* Jacq.) and its preservation after drying process. *Future Foods* 4: 1-11. <https://doi.org/10.1016/j.fufo.2021.100070>
- Peakall R and PE Smouse (2012) GenAlEx 6.5: Genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinform.* 28: 2537-2539.
- Pérez-Martínez AL, LE Eguarte, KL Mercer, NE Martínez-Ainsworth, L McHale, E Van der Knaap and L Jardón-Barbolla (2022) Genetic diversity, gene flow, and differentiation among wild, semiwild, and landrace chile pepper (*Capsicum*

- annuum*) populations in Oaxaca, Mexico. *Am. J. Bot.* 109: 1157-1176.
- Perrier X (2006) DARwin software. <http://darwin.cirad.fr/darwin>
- Pickersgill B (2007) Domestication of plants in the Americas: Insights from Mendelian and molecular genetics. *Ann. Bot.* 100(5):925-940.
- Powell W, M Morgante, C Andre, M Hanafey, J Vogel, S Tingey and A Rafalski (1996) The comparison of RFLP, RAPD, AFLP and SSR (microsatellite) markers for germplasm analysis. *Mol. Breed.* 2(3): 225-238.
- Pritchard JK, M Stephens and P Donnelly (2000) Inference of population structure using multilocus genotype data. *Genet.* 155: 945-959. <https://doi.org/10.1093/genetics/155.2.945>
- Rahevar PM, JN Patel, S Kumar and DJ Parmar (2021) Morphological characterization of 58 chilli genotypes. *Indian J. Plant Genet. Resour.* 34(02): 221-228. <https://doi.org/10.5958/0976-1926.2021.00021.8>
- Rai MK (2023) Start codon targeted (SCoT) polymorphism marker in plant genome analysis; current status and prospects. *Planta* 257(2):34. doi: 10.1007/s00425-023-04067-6
- Ram H, DK Samadia, AK Verma and PS Gurjar (2024) Surface covering protective vegetable nursery for hot arid climate. *Indian Hort.* 69(1): 26-28.
- Reyes-Escogido ML, EG Gonzalez-Mondragon and E Vazquez-Tzompantzi (2011) Chemical and pharmacological aspects of capsaicin. *Molecules* 16(2): 1253-1270.
- Rohlf FJ (2000) NTSYS-pc: Numerical Taxonomy and Multivariate Analysis System Version 2.1. Exeter Publishing Setauket, New York, USA.
- Samadia DK, AK Verma and H Ram (2024) Characterization and utilization of potential genetic resources for improvement of *Citrullus lanatus* under desert ecosystem. *Indian J. Plant Genet. Resour.* 37(03): 482-493. <https://doi.org/10.61949/0976-1926.2024.v37i03.11>
- Semagn K, A Bjornstad and MN Ndjioudjop (2006) An overview of molecular marker methods for plants. *Afr. J. Biotechnol.* 5(25): 2540-2568.
- Shahlaei A, S Torabi and M Khosroshah (2014) Efficacy of SCoT and ISSR markers in assesment of tomato (*Lycopersicon-esculentum* Mill.) genetic diversity. *Int. J. Biosci.* 5(2): 14-22.
- Shilpha J, S Pandian, MJ Largia, SI Sohn and M Ramesh (2021) Short-term storage of *Solanum trilobatum* L. synthetic seeds and evaluation of genetic homogeneity using SCoT markers. *Plant Biotechnol. Rep.* 15: 651-661.
- Taranto F, N D'Agostino, B Greco, T Cardi and P Tripodi (2016) Genome-wide SNP discovery and population structure analysis in pepper (*Capsicum annuum*) using genotyping by sequencing. *BMC Genom.* 17: 943.
- Tsabalala A, I Ganopoulos, A Timplalexi, X Aliki, I Bosmali, NO Irini, T Athanasios and P Madesis (2015) Molecular characterization of Greek pepper (*Capsicum annuum* L.) landraces with neutral (ISSR) and gene-based (SCoT and EST-SSR) molecular markers. *Biochem. Systemat. Ecol.* 59: 256-263. <https://doi.org/10.31883/pjfn/133690>
- Xiong F, R Zhong, Z Han, J Jiang, L He, W Zhuang and R Tang (2011) Start codon targeted polymorphism for evaluation of functional genetic variation and relationships in cultivated peanut (*Arachis hypogaea* L.) genotypes. *Mol. Biol. Rep.* 38(5): 3487-3494. <https://doi.org/10.1007/s11033-010-0459-6>
- Yumnam JS, W Tyagi, A Pandey, TM Ngasepam and M Rai (2012) Evaluation of genetic diversity of chilli landraces from Northeastern India based on morphology, SSR markers and the Pun1 locus. *Plant Mol. Biol. Rep.* 30(6): 1470-1479.

RESEARCH ARTICLE

Assessment of Antioxidant Activity and Quality Attributes in Indigenous Pran (Kashmiri Shallot) Genotypes

Gazala Nazir^{1*}, Suraya Ahsan¹, Mudasir Magray¹, Khursheed Hussain¹, Saima Tabasum¹, UH Masoodi¹, Syed Berges¹, Asima Amin¹, Abida Jabeen² and AA Lone³

Abstract

Pran is a condiment vegetable of the Kashmir region and belongs to family Alliaceae. Indigenous landraces of *Alliums* represent an important reservoir of nutritional and functional diversity; however, they remain poorly characterized in many mountain agro-ecosystems. Pran also known as Kashmiri shallot (*Allium x cornutum clementii* ex vis.), is widely cultivated for its distinctive pungency and culinary value, yet comprehensive information on its biochemical attributes is not available. The present study aimed to characterize the antioxidant potential, nutritional composition, and quality traits of indigenous Pran genotypes collected from diverse agro-ecological regions of Jammu and Kashmir. Thirty Pran genotypes along with a commercial check were evaluated for antioxidant activity (DPPH radical scavenging), vitamin C, total phenols, dry matter, total soluble solids (TSS), pyruvic acid, and total sugars. Significant genotypic variation ($p \leq 0.05$) was observed for all traits studied. Antioxidant activity ranged from 49.57 to 71.12%, vitamin C from 2.93 to 3.51 mg 100 g⁻¹, and total phenolics from 3.12 to 9.12 mg 100 g⁻¹. Dry matter and TSS varied from 13.20 to 23.89% and 9.79 to 22.14 °Brix, respectively. Pyruvic acid content ranged from 4.34 to 5.33%, while total sugars varied between 5.71 and 8.77%. Several genotypes exhibited superior performance across multiple biochemical and quality parameters, significantly outperforming the check. The pronounced biochemical diversity observed in Kashmiri Pran highlights its potential as a valuable genetic resource for nutritional improvement, the development of functional foods, and the sustainable utilisation of Pran germplasm. The identified elite genotypes may serve as promising candidates for breeding programmes and conservation initiatives in temperate and high-altitude regions

Keywords: Antioxidant activity, Biochemical diversity, Indigenous germplasm, Pran

¹Division of Vegetable Science, Sher-e-Kashmir University of Agricultural Sciences and Technology, Shalimar, Srinagar, Kashmir, 190025, India.

²Division of Food Science and Technology- Sher-e-Kashmir University of Agricultural Sciences and Technology, Shalimar, Srinagar, Kashmir, 190025, India.

³Department of Genetics and Plant Breeding, Dryland Agriculture Research Station, Rangreth, Srinagar, 191132, India.

*Author for correspondence:

gazalanazir6@gmail.com

Received: 15/01/2026

Revised: 15/04/2026

Accepted: 17/04/2026

How to cite this article: Nazir G, S Ahsan, M Magray, K Hussain, S Tabasum, UH Masoodi, S Berges, A Amin, A Jabeen and AA Lone (2026) Assessment of Antioxidant Activity and Quality Attributes in Indigenous Pran (Kashmiri Shallot) Genotypes. *Indian J. Plant Genet. Resour.* 39(2): 38-42.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.175204>

Introduction

Onion and shallot are among the most widely consumed vegetable crops globally and are valued not only for their culinary importance but also for their health-promoting properties. One notable species, Pran, is distinguished by its unique flavor and is widely used as a condiment in soups, meat dishes and salads (Suraya *et al.*, 2025). It holds a special place in Kashmiri cuisine, where the renowned “Wazwan” is considered incomplete without Pran. It is an important ingredient of chilli cakes commonly known as ‘Wari’. These chilli cakes are sold at exorbitant prices and even exported to Middle Eastern countries by certain private companies and can be stored up to a period of two to three years under ordinary conditions (Jabeen *et al.*, 2012). The simple paste, prepared from the Pran, not only enhances the taste but also holds a treasure trove of health benefits. As Pran is a type of shallot, shallots (*A. cepa* var. *aggregatum*) are rich in bioactive compounds such as phenolics, flavonoids, sulfur-containing compounds, and

vitamin C, which contribute to their antioxidant capacity and distinctive flavour (Griffiths *et al.*, 2002; Slimestad *et al.*, 2007). This fact led us to characterize the diverse Pran genotypes growing in parts of Jammu and Kashmir and find out their relative quality parameters. Pran is a triploid, viviparous onion with a moreslender stature than onion and more pinkish, flushed flowers. Pran is an integral component of Kashmiri cuisine (wazwan). It is typically propagated vegetatively and exhibits considerable variation in bulb size, pungency, and taste. Despite its local importance and adaptability to temperate and high-altitude conditions, Pran remains underutilized and scientifically under explored. Characterization of its biochemical and quality traits is essential for identifying nutritionally superior genotypes and for promoting its conservation and improvement. The present study aimed to study the extent of biochemical diversity among indigenous Kashmiri Pran genotypes with particular emphasis on antioxidant potential, nutritional composition, and quality attributes, and to identify elite genotypes suitable for breeding and value addition.

Materials and Methods

Plant material

Thirty indigenous diversePran genotypes were collected from different agro-ecological regions of Jammu and Kashmir. A commercially grown variety was included as a local check. The experiment was conducted at Urban Technological Park Habbak, Division of Vegetable Science, Sher-e-Kashmir University of Agricultural Science and Technology of Kashmir during *Rabi* 2023-2024. The experimental field of Urban Technological Park in Habbak is situated at an elevation of 1608 meters above sea level, between 34.16° N latitude and 74.83° E longitude. The single-factor experiment was set up in a Randomized Block Design (RBD) with three replications. The climate is temperate with mild summers. The bulbs were sown at depth of 3-4 cm, spacing of 20 cm × 15 cm, farmyard manure 25-30 tons per hectare and 100:80:60 Kg/ha of NPK were applied (Saleem *et al.*, 2023)

Biochemical analysis

Fresh bulb samples were analyzed for biochemical and quality traits using standard procedures. Antioxidant activity was estimated using the DPPH radical scavenging assay (Brand-William *et al.*, 1995). Vitamin C content was determined by titration method and expressed as mg 100/g fresh weight (Thimmaiah 1999). Total phenolic content was measured using the Folin–Ciocalteu method and expressed as mg 100/g (Bray and Thorpe, 1954). Dry matter content was determined by oven drying. Total soluble solids (TSS) were measured using a digital refractometer (Digital Abbe Refractometer, Kruss Optroic AOAC, 1980). Pyruvic acid content, an indicator of pungency, was estimated following Schwimmer and Weston (1961). Total sugars were estimated using standard biochemical methods (A.O.A.C.1980).

Statistical analysis

The experimental data obtained for different biochemical and quality traits were subjected to statistical analysis using analysis of variance (ANOVA) appropriate for the experimental design as described by Gomez and Gomez (1984). The significance of differences among genotypes was tested at 5% level of probability ($p \leq 0.05$).

The standard error of difference (SED) was calculated to estimate the variability among treatment means, and the critical difference (CD) values were worked out to compare the significance between genotypic means. Wherever the calculated difference between two means exceeded the CD value, the difference was considered statistically significant.

All observations were recorded on randomly selected samples, and mean values were used for statistical computation. The single-factor experiment was set up in a Randomized Block Design (RBD) with three replications.

Results

Antioxidant activity and Vitamin C content

The data presented in Table 1 revealed significant

Table1: Mean Performance of different genotypes of Pran with respect to antioxidant activity and quality traits

S.No.	Genotype	Antioxidant (% DPPH inhibit)	Vitamin-C (mg/100 g)	Phenols (mg/100 g)	Dry matter (%)	TSS (°Brix)	Pyruvic acid (mg/100 g fresh weight)	Total sugars (%)
1	SKAU-PRAN-H	51.25	2.93	7.51	18.41	10.24	4.65	5.71
2	SKAU-PRAN-Kup	50.16	2.96	9.11	19.78	13.9	5.11	6.23

S.No.	Genotype	Antioxidant (% DPPH inhibit)	Vitamin-C (mg/100 g)	Phenols (mg/100 g)	Dry matter (%)	TSS (°Brix)	Pyruvic acid (mg/100 g fresh weight)	Total sugars (%)
3	SKAU-PRAN-P	67.51	3.12	7.13	15.11	16	4.72	7.31
4	SKAU-PRAN-Gbl	61.27	3.08	4.12	23.77	19.22	4.81	6.16
5	SKAU-PRAN-Bo	67.06	3.48	5.23	13.2	13.89	5.29	8.77
6	SKAU-PRAN-K	49.57	2.94	6.31	18.1	18.09	4.34	6.13
7	SKAU-PRAN-Bnd	71.12	3.51	3.12	23.89	19.77	5.33	8.57
8	SKAU-PRAN-So	68.25	3.24	8.422	2.29	18.9	5.08	5.89
9	SKAU-PRAN-Q	65.15	3.34	8.13	15.23	16.89	5.13	8.42
10	SKAU-PRAN-So1	65.28	3.16	3.12	19.79	18.78	4.96	7.68
11	SKAU-PRAN-So2	65.97	2.98	3.21	19.33	17.89	4.89	7.55
12	SKAU-PRAN-L1	58.24	2.97	3.81	19.92	1.86	4.58	8.06
13	SKAU-PRAN-L2	59.13	2.93	3.43	19.51	20.88	4.62	8.12
14	SKAU-PRAN-L3	57.68	3.11	3.72	18.29	20.24	4.74	8.22
15	SKAU-PRAN-Sh	62.58	3.15	7.13	22.81	7.87	4.92	8.18
16	SKAU-PRAN-A	60.78	3.18	3.85	18.25	19.87	4.95	8.26
17	SKAU-PRAN-Kul	60.27	3.49	9.12	17.26	19.06	4.97	8.32
18	SKAU-PRAN-S1	68.99	3.32	3.81	19.2	29.82	4.99	7.69
19	SKAU-PRAN-Ch	59.15	3.42	5.51	17.92	2.14	5.24	7.54
20	SKAU-PRAN-S	269.82	3.49	3.51	22.71	17.97	4.89	7.45
21	SKAU-PRAN-S	368.45	3.45	4.12	21.24	18.81	5.18	7.76
22	SKAU-PRAN-S4	69.23	3.38	4.21	20.31	8.98	5.12	7.85
23	SKAU-PRAN-S5	68.85	3.44	5.52	19.24	17.68	5.24	7.33
24	SKAU-PRAN-S6	68.57	3.46	4.93	23.8	9.79	5.32	7.66
25	SKAU-PRAN-Bud	60.25	3.37	3.52	19.21	4.55	5.19	8.11
26	SKAU-PRAN-T	67.51	3.46	6.83	17.17	14.89	5.02	7.36
27	SKAU-PRAN-Ko	58.25	3.35	4.12	19.11	19.41	4.76	8.12
28	SKAU-PRAN-Qa	58.26	3.23	8.06	16.45	17.85	4.83	8.26
29	SKAU-PRAN-Brl	65.21	3.16	7.89	21.02	17.91	4.83	8.14
30	SKAU-PRAN-S7	69.23	3.41	3.78	20.49	10.78	5.17	7.53
	Local Check	66.33	3.14	7.28	22.13	18.17	5.02	6.26
	SED	0.172	0.0045	0.0045	0.121	0.0025	0.086	0.172
	CD	0.345	0.009	0.009	0.242	0.005	0.172	0.345

differences among Pran genotypes for antioxidant activity and vitamin C content. Antioxidant activity, expressed as per cent DPPH inhibition, ranged from 49.57 to 71.12%. The highest antioxidant activity was recorded in SKAU-PRAN-Bnd (71.12%), followed by SKAU-PRAN-S2 (69.82%), SKAU-PRAN-S4 and SKAU-PRAN-S7 (69.23%), all of which were significantly superior to the check. The lowest antioxidant activity was observed in SKAU-PRAN-K (49.57%). Vitamin C content varied from 2.93 to 3.51 mg/100 g. Maximum vitamin C was recorded in SKAU-PRAN-Bnd (3.51 mg/100 g), followed by SKAU-PRAN-S2 (3.49 mg/100 g).

Several genotypes showed significantly higher vitamin C content than the check, indicating substantial nutritional variability within the Pran germplasm.

Total phenols

Table 1 depicts the variation in total phenol content among Pran genotypes, which ranged from 3.12 to 9.12 mg/100 g. The highest phenol content was recorded in SKAU-PRAN-Kul (9.12 mg/100 g), closely followed by SKAU-PRAN-Kup (9.11 mg/100 g) and SKAU-PRAN-So (8.42 mg/100 g). Genotypes with lower phenolic content included SKAU-PRAN-Bnd, SKAU-PRAN-So1,

and SKAU-PRAN-So2. The observed variation suggests the presence of genotypes with enhanced nutraceutical potential suitable for functional food development.

Dry matter and total soluble solids (TSS)

Dry matter content exhibited wide variation among the genotypes, ranging from 13.20% to 23.89% (Table 1). The highest dry matter content was recorded in SKAU-PRAN-Bnd (23.89%), followed by SKAU-PRAN-S6 (23.80%) and SKAU-PRAN-Gbl (23.77%), whereas SKAU-PRAN-Bo (13.20%) recorded the lowest value. Total soluble solids (TSS) ranged from 9.79°Brix to 22.14°Brix. The highest TSS was observed in SKAU-PRAN-Ch (22.14°Brix), followed by SKAU-PRAN-L1 (21.86°Brix) and SKAU-PRAN-L2 (20.88°Brix). Higher TSS values indicate better sweetness and suitability for processing and culinary purposes.

Pyruvic acid and total sugar

Pyruvic acid content, an indicator of pungency, varied significantly among the genotypes (Table 1), ranging from 4.34 to 5.33%. The highest pyruvic acid content was recorded in SKAU-PRAN-Bnd (5.33%), followed by SKAU-PRAN-S6 (5.32%) and SKAU-PRAN-Bo (5.29%),

indicating their strong pungency compared to the check. Total sugar content ranged from 5.71% to 8.77%. The highest total sugars were observed in SKAU-PRAN-Bo (8.77%), followed by SKAU-PRAN-Bnd (8.57%), SKAU-PRAN-Kul (8.32%), and SKAU-PRAN-A (8.26%). Genotypes with higher sugar content are likely to possess better taste and consumer acceptability.

Genotypes such as SKAU-PRAN-Bnd, SKAU-PRAN-Bo, SKAU-PRAN-Kul, and SKAU-PRAN-S2 consistently ranked among the top performers across multiple biochemical and quality traits, making them promising candidates for conservation, varietal improvement, and value-added utilization.

Discussion

The present investigation revealed substantial biochemical and quality-related variability among Kashmiri Pran (shallot) genotypes, highlighting the richness of indigenous germplasm and its potential for nutritional enhancement and varietal improvement. Similar wide variability for biochemical traits has been earlier reported in onion and shallot genotypes, suggesting strong genetic control influenced by environmental adaptation (Griffiths *et al.*, 2002).

Table 2: Comparative summary of top-performing pran (Kashmiri Shallot) genotypes for biochemical and quality traits

Trait	Range Observed	Top Performing Genotype(s)	Value
Antioxidant activity (% DPPH inhibition)	49.57 – 71.12	SKAU-PRAN-Bnd	71.12%
		SKAU-PRAN-S2	69.82%
		SKAU-PRAN-S4 / S7	69.23%
Vitamin C (mg/100 g)	2.93 – 3.51	SKAU-PRAN-Bnd	3.51
		SKAU-PRAN-Bo / S2	3.49
Total phenols (mg/100 g)	3.12 – 9.12	SKAU-PRAN-Kul	9.12
		SKAU-PRAN-Kup	9.11
		SKAU-PRAN-So	8.42
Dry matter (%)	13.20 – 23.89	SKAU-PRAN-Bnd	23.89
		SKAU-PRAN-S6	23.80
		SKAU-PRAN-Gbl	23.77
TSS (°Brix)	9.79 – 22.14	SKAU-PRAN-Ch	22.14
		SKAU-PRAN-L1	21.86
		SKAU-PRAN-L2	20.88
Pyruvic acid (%) – pungency	4.34 – 5.33	SKAU-PRAN-Bnd	5.33
		SKAU-PRAN-S	65.32
		SKAU-PRAN-Bo	5.29
Total sugars (%)	5.71 – 8.77	SKAU-PRAN-Bo	8.77
		SKAU-PRAN-Bnd	8.57
		SKAU-PRAN-Kul	8.32

Antioxidant activity and vitamin C

Antioxidant activity varied widely among the Pran genotypes, with several genotypes recording values significantly higher than the check (Table 2). High antioxidant potential in *Allium* species is primarily attributed to the presence of phenolics, flavonoids, vitamin C, and sulfur-containing compounds (Slimestad *et al.*, 2007). Genotypes such as SKAU-PRAN-Bnd and SKAU-PRAN-S2, which exhibited superior antioxidant activity, may therefore serve as valuable sources of health-promoting compounds. Although onions are generally considered moderate sources of vitamin C, the observed variation among Pran genotypes aligns with earlier findings that genotype plays a crucial role in determining ascorbic acid content (Kumar *et al.*, 2015). Higher vitamin C levels in certain genotypes enhance their dietary value and contribute synergistically to antioxidant activity.

Total phenolic content

Phenolic compounds are major contributors to antioxidant capacity and are known to vary widely among onion cultivars (Lachman *et al.*, 2003). In the present study, phenolic content ranged considerably, with SKAU-PRAN-Kul and SKAU-PRAN-Kup recording the highest values. These findings corroborate earlier reports by Prakash *et al.*, (2007), who observed significant genotypic differences in total phenols among onion cultivars. High-phenol genotypes are particularly important for functional food development and may offer enhanced protection against oxidative stress-related disorders (Yang *et al.*, 2004).

Dry matter and total soluble solids (TSS)

Dry matter content is a critical quality trait influencing storage life, processing suitability, and pungency in onions and shallots. Genotypes with higher dry matter content, such as SKAU-PRAN-Bnd and SKAU-PRAN-S6, are considered desirable for dehydration and processing purposes. Similar associations between high dry matter content and better processing quality have been reported earlier. Total soluble solids, which are closely associated with sweetness and flavour intensity, also varied significantly among genotypes. High TSS values observed in certain Pran genotypes are consistent with earlier reports that attribute such variation to differences in carbohydrate accumulation and genetic background. Genotypes combining high dry matter and high TSS may thus possess superior culinary and commercial value.

Pyruvic acid and total sugars

Pyruvic acid content is widely used as an indicator of pungency in onions and shallots. The significant

variation observed among Pran genotypes reflects differences in sulfur metabolism and enzymatic activity, as also reported by Schwimmer and Weston (1961). Genotypes such as SKAU-PRAN-Bnd and SKAU-PRAN-S6, which recorded higher pyruvic acid content, may be preferred for consumers favouring strong pungency and traditional flavour profiles. Total sugars contribute directly to sweetness and overall palatability. Genotypes with higher sugar content, notably SKAU-PRAN-Bo and SKAU-PRAN-Kul, are likely to be more acceptable for fresh consumption. Previous studies have demonstrated an inverse or complex relationship between sugar content and pungency, depending on genotype and environmental conditions, which may explain the observed variation among Pran genotypes.

References

- AOAC (1980) *Official Methods of Analysis*. Association of Official Analytical Chemists, Washington, DC.
- Brand-Williams W, ME Cuvelier and C Berset (1995) Use of a free radical method to evaluate antioxidant activity. *LWT - Food Sci. Technol.* 28: 25-30.
- Bray HG and WV Thorpe (1954) *Methods of Biochemical Analysis*. Kalyani Publications, New Delhi.
- Gomez KA and AA Gomez (1984) *Statistical Procedures for Agricultural Research*, 2nd edn. John Wiley & Sons, New York, USA.
- Griffiths G, L Trueman, T Crowther, B Thomas and B Smith (2002) Onions; a global benefit to health. *Phytother. Res.* 16: 603-615.
- Jabeen N, B Afroza, SH Khan and MI Makhdoomi (2012) Kashmiri chilli; a gold mine. *Indian J. Arecanut Spices Med. Plants* 14(2): 33-36.
- Kumar S, M Imtiyaz and R Singh (2015) Effect of genotypes on quality parameters of onion (*Allium cepa* L.). *Indian J. Agric. Sci.* 85: 123-127.
- Lachman J, D Proněk, A Hejtmánková, J Dudjak, V Pivec and K Faitová (2003) Total polyphenol and main flavonoid antioxidants in different onion varieties. *Hort. Sci. (Prague)* 30: 142-147.
- Prakash D, BN Singh and G Upadhyay (2007) Antioxidant and free radical scavenging activities of phenols from onion (*Allium cepa*). *Food Chem.* 102: 1389-1393.
- Saleem T, SH Khan, N Jabeen and T Sultan (2023) Pran: Endemic edible *Allium* of Kashmir. *Agric. Articles* 3(1).
- Schwimmer S and WJ Weston (1961) Enzymatic development of pyruvic acid in onion as a measure of pungency. *J. Agric. Food Chem.* 9: 301-304.
- Slimestad R, T Fossen and IM Vågen (2007) Onions; a source of unique dietary flavonoids. *J. Agric. Food Chem.* 55: 10067-10080.
- Suraya Ahsan, G Nazir, AA Lone, S Tabasum, N Nazir, I Ahmad, K Hussain, U Masoodi and S Berjes (2025) Correlation coefficient and path analysis studies in native collections of Kashmiri pran (*Allium × cornutum*). *Plant Arch.* 25(1): 2282-2286.
- Thimmaiah SK (1999) *Standard Methods of Biochemical Analysis*. Kalyani Publications, New Delhi.
- Yang J, KJ Meyers, J van der Heide and RH Liu (2004) Varietal differences in phenolic content and antioxidant activity of onions. *J. Agric. Food Chem.* 52: 6787-6793.

RESEARCH ARTICLE

Morpho-biochemical Characterization and Multivariate Assessment of Genetic Diversity in Rajmah (*Phaseolus vulgaris* L.) Genotypes from Northeast India

Bedashruti Saikia¹, Rupam Borgohain^{2*}, Gayatree Goswami², Jutika Das² and Anubrat Borah²

Abstract

The present study assessed morphological, yield, and biochemical diversity among forty rajmah (*Phaseolus vulgaris* L.) genotypes collected from five north eastern states of India, representing both vine and dwarf growth habits. Field experiments were conducted during the *rabi* seasons of 2022–23 and 2023–24 at Assam Agricultural University, Jorhat, using a randomized block design with three replications. Eighteen traits were studied, including ten morphological and eight biochemical parameters and were evaluated through univariate and multivariate analyses. Analysis of variance revealed highly significant genotypic differences for all traits, indicating substantial genetic variability. Vine types exhibited greater plant height, branching, pod number, and seed yield, whereas dwarf types recorded higher 100-seed weight and superior nutritional attributes, including fat, fibre, niacin, vitamin B₆, and folic acid. Correlation analysis showed strong positive associations between yield and key plant architecture traits. Principal component, cluster, and heatmap analyses grouped genotypes primarily by growth habit and regional origin. The outcomes of the study signify rich genetic diversity in the rajmah germplasm of Northeast India and identify key traits for simultaneous improvement of yield and nutritional quality.

Keywords: Biochemical traits, Genetic diversity, Multivariate analysis, Northeast India, *Phaseolus vulgaris*

¹Department of Plant Breeding and Genetics, Assam Agricultural University, Jorhat, Assam, 785013, India.

²Directorate of Research (Agri), Assam Agricultural University, Jorhat, Assam, 785013, India.

***Author for correspondence:**

rupam.borgohain@aau.ac.in

Received: 02/01/2026

Revised: 22/04/2026

Accepted: 24/04/2026

How to cite this article: Saikia B, R Borgohain, G Goswami, J Das and A Borah (2026) Morpho-biochemical Characterization and Multivariate Assessment of Genetic Diversity in Rajmah (*Phaseolus vulgaris* L.) Genotypes from Northeast India. *Indian J. Plant Genet. Resour.* 39(2): 43-52.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.174765>

Introduction

The common bean (*Phaseolus vulgaris* L.) is among the most important grain legumes globally, serving as a major source of dietary protein, complex carbohydrates, fibre, and essential micronutrients such as iron and zinc (Broughton *et al.*, 2003; Blair *et al.*, 2010). Owing to its adaptability and nitrogen-fixing ability, it plays a vital role in food and nutritional security, particularly for smallholder farmers (Beebe *et al.*, 2010). In India, especially in the North-Eastern Hill Region, rajmah (dry beans) is a key component of local diets and cropping systems, cultivated across diverse altitudes and micro-climates (Singh *et al.*, 2014; Rai *et al.*, 2020). Despite its importance, indigenous rajmah germplasm of Northeast India remains poorly characterized for morphological, yield, and biochemical diversity.

Originating in the Americas (Kwak & Gepts, 2009), common bean was likely introduced into India through early trade and colonial routes during the 16th century (Indian Farming, 2017). Subsequent diversification under varied Himalayan micro-climates, local selection, and regional isolation fostered a secondary centre of diversity in Northeast India (Dutta *et al.*, 2015; Banoo *et al.*, 2020).

Previous studies have reported substantial variability in *P. vulgaris* for plant height, pod length, seed weight, and maturity period (Raturi *et al.*, 2017; Chahota *et al.*, 2020). Multivariate tools such as cluster and principal component analyses have been effective in identifying divergent groups and key discriminating traits (Andrade *et al.*, 2019; Girish *et al.*, 2021). Biochemical profiling for protein, fat, fibre, and vitamins has also provided valuable inputs for nutritional improvement in common bean (Celmeli *et al.*, 2018; Khan *et al.*, 2021). However, integrated multivariate evaluations combining morphological and biochemical traits remain limited for rajmah germplasm.

Multivariate approaches, including principal component analysis, hierarchical clustering, and correlation analysis, efficiently reveal trait interrelationships and major dimensions of variation (Jain *et al.*, 2019; Singh *et al.*, 2021) and have differentiated growth habits, regional adaptation, and nutritional quality groups in legumes (Oliveira *et al.*, 2019; Mwale *et al.*, 2020). Given the ecological and cultural diversity of Northeast India, rajmah genotypes are expected to exhibit wide variation shaped by environment and farmer-driven selection; however, systematic quantitative analyses remain scarce. Therefore, the present study aimed to assess variability among rajmah genotypes through integrated analyses of morphological, yield, and biochemical traits, to identify key traits for genetic improvement and nutritional enhancement.

Materials and Methods

Experimental site and plant materials

The study was conducted using 40 rajmah genotypes collected from five north-eastern states of India viz., Assam, Meghalaya, Mizoram, Arunachal Pradesh, and Nagaland represent both dwarf and vine growth habits (Supplementary Table 1). Genotypes were labelled using state-specific prefixes followed by numeric codes: AS (Assam), MR (Mizoram), NL (Nagaland), AR (Arunachal Pradesh), and ML (Meghalaya). Field experiments were carried out during the *rabi* seasons of 2022–23 and 2023–24 at the Instructional-cum-Research (ICR) Farm, Assam Agricultural University (AAU), Jorhat (26.75° N, 94.22° E; 87 m asl). The experiment followed a Randomized Block Design with three replications as per Fisher (1935). Each plot comprised two rows (3 m length, 45 cm × 20 cm spacing), and standard agronomic practices were adopted to ensure uniform crop growth.

Trait measurement and biochemical estimation

Eighteen traits were studied, including ten morphological (plant height, branches per plant, chlorophyll content index, days to 50% flowering, days to maturity, pods per plant, pod length, seeds per pod, 100-seed weight, and yield t/ha) and eight biochemical parameters (crude protein, fat, fibre, cooking time, riboflavin, niacin, vitamin B₆, and folic acid). Chlorophyll Content Index (CCI) was measured using a SPAD meter by taking non-destructive readings from the fully expanded upper canopy leaves of each genotype. For each replication, SPAD readings were recorded from 5 plants, and the values were averaged to obtain a representative chlorophyll estimate.

Crude protein, fat, and fibre contents were analysed following AOAC (2016) procedures. Vitamins were estimated spectrophotometrically using Lowry *et al.*, (1951) with modifications for legume seeds. Each analysis was performed in triplicate, and mean values were used for statistical computation.

Statistical analysis

Descriptive statistics (mean, range, SD, CV %) were computed to assess variability. Data normality was verified using Shapiro–Wilk tests and Q–Q plots. Analysis of variance was performed using the randomized block design model:

$$Y_{ij} = \mu + G_i + R_j + \epsilon_{ij}$$

Where Y_{ij} represents the observed value of the i^{th} genotype in the j^{th} replication, μ is the overall mean, G_i is the fixed effect of the i^{th} genotype, R_j is the random effect of the j^{th} replication, and ϵ_{ij} is the random error term associated with each observation.

Significant differences were separated using Tukey's HSD test ($\alpha = 0.05$), while non-parametric data were tested using Kruskal–Wallis and Dunn's post-hoc tests (Gomez & Gomez, 1984). A Welch's t-test was used to compare dwarf and vine growth types. All analyses were performed in R software (version 4.4.2; R Core Team, 2024) using *rstatix*, *ggplot2*, and *dplyr* packages (Wickham, 2016).

Multivariate and correlation analyses

Pearson's correlation coefficients were calculated to evaluate inter-trait associations and visualized through heatmaps (*corrplot*, *ggcorrplot*; Friendly, 2002). Principal Component Analysis (PCA) was carried out on standardized data (mean = 0, SD = 1) following Jolliffe (2002) to identify key traits contributing to variability. Eigenvalues > 1 were retained (Kaiser's criterion), and scree plots and biplots were generated to visualize trait

loadings and genotype grouping.

Hierarchical clustering was performed on principal component scores using Euclidean distance and Ward's linkage (Sneath & Sokal, 1973) to classify genotypes by plant type and region, thereby reducing multicollinearity among traits and ensuring that clustering was driven by the major, independent sources of phenotypic variation; regional variation in standardized mean trait values was visualized using heatmaps (Gu, 2016).

Data visualization

All graphical outputs, including PCA biplots, dendrograms, and heatmaps, were generated in R using the packages ggplot2, factoextra, ggdendro, and pheatmap for reproducibility and visual clarity.

Results and Discussion

Variation among genotypes

Analysis of variance revealed highly significant differences ($P < 0.001$) among the 40 rajmah genotypes for all morphological and biochemical traits, except seeds per pod, which also differed significantly ($P < 0.01$). This pronounced variability reflects the broad genetic base of germplasm collected from diverse agro-ecological regions of Northeast India and indicates substantial scope for selection and trait-based genetic improvement (Dutta *et al.*, 2015; Banoo *et al.*, 2020; de Paula *et al.*, 2024).

Descriptive statistics (Table 1) revealed appreciable variation for several agronomic traits, with plant height ranging from 30.00 to 192.33 cm ($CV = 15.75\%$), branches per plant from 10.67 to 30.33 ($CV = 7.59\%$), and pods per plant from 14.33 to 70.00 ($CV = 4.57\%$). Days to maturity varied from 110.33 to 132.00 days, indicating the presence of both early- and late-maturing genotypes. Seed yield ranged from 1.03 to 2.77 t/ha, demonstrating substantial scope for yield improvement. Among quality traits, 100-seed weight

varied widely from 20.67 to 105.33 g, reflecting marked diversity in seed size and market class preferences. Crude protein ranged from 14.03 to 23.47%, fat from 0.60 to 4.07%, crude fibre from 4.75 to 23.26%, and folic acid from 242.81 to 336.37 μg , indicating significant nutritional variability among genotypes.

Welch's t-test (data not shown) revealed clear phenotypic differentiation between growth habits. Vine types recorded significantly higher plant height, branching, pod number, seeds per pod, phenological duration, seed yield, and cooking time, reflecting vigorous growth and longer crop duration. In contrast, dwarf types exhibited significantly higher 100-seed weight and superior nutritional attributes, including crude fibre, fat, niacin, vitamin B₆, and folic acid contents. Traits such as chlorophyll content index, pod length, crude protein, and riboflavin did not differ significantly between growth habits, indicating selective rather than uniform effects of growth habit on trait expression. Similar growth habit-driven patterns have been reported in common bean germplasm (Gelaw, 2017; Shah *et al.*, 2020).

Following the detection of significant genotypic differences in ANOVA, Tukey's HSD post-hoc test was applied to identify specific pairwise differences among genotypes for each trait. Table 1 integrates descriptive statistics with trait desirability criteria and superior-performing genotypes identified through HSD comparisons. Vine genotypes were prominent for growth and yield-related traits, whereas several dwarf genotypes excelled in seed size, earliness, cooking quality, and nutritional attributes. Early-maturing genotypes were particularly relevant for short-duration and multiple cropping systems, while nutrient-dense entries may serve as valuable donors for biofortification. Overall, growth habit emerged as a major source of variability, and the evaluated genotypes displayed broad agronomic and nutritional diversity, consistent with earlier reports in common bean (Ahmad, 2018; de Paula *et al.*, 2024).

Table 1: Descriptive statistics, trait desirability, and superior genotype performance based on Tukey's HSD analysis for major traits of rajmah

Trait	Mean $\pm$ SD	Range (Min–Max)	CV (%)	HSD (0.05)	Nature of desirable performance	Desirable genotypes with mean values	Remarks
Plant height (cm)	83.23 $\pm$ 52.50	30–192.33	15.75	43.29	Vine: Taller plants (180–250cm) desirable Dwarf: Short plants (30–40cm) desirable	Vine: NL3 (192.33); AR7 (183.33); AR13 (182.00) Dwarf: AS2 (36.33); AS1 (35.33); NL5 (30.00)	Height determines plant architecture and suitability for vine or dwarf production systems.

Trait	Mean $\pm$ SD	Range (Min–Max)	CV (%)	HSD (0.05)	Nature of desirable performance	Desirable genotypes with mean values	Remarks
Branches/plant	19.44 $\pm$ 5.08	10.67–30.33	7.59	4.88	Higher desirable in both types	Vine: AR13 (30.33); NL3 (28.67); AR4 (27.67) Dwarf: AS7 (20.67); AS3 (20.33); AS4 (19.67)	Higher branching may increase reproductive sites and yield potential.
Days to maturity	120.32 $\pm$ 7.08	110.33–132	1.28	5.09	Lower desirable in both types	Vine: NL3 (128.00); AR11 (127.00); AR9 (127.00) Dwarf: MR1 (115.00); AS3 (114.33); AR12 (114.00)	Early maturity is useful for short-duration and multiple cropping systems.
Pods/plant	38.32 $\pm$ 14.73	14.33–70.0	4.57	5.78	Higher desirable in both types	Vine: AR7 (70.00); NL3 (68.33); AR6 (67.67) Dwarf: AS11 (37.33); AS3 (37.00); AS10 (36.00)	More pods per plant directly support higher productivity.
100-seed weight (g)	58.98 $\pm$ 16.93	20.67–105.33	2.21	4.30	Higher desirable (50–100 g)	Vine: AR2 (105.33); Dwarf: AR3 (88.33); AR12 (85.00)	Bold seeds are preferred for market value and consumer acceptance.
Seed yield (t ha ¹)	1.79 $\pm$ 0.47	1.03–2.77	12.47	0.74	Higher desirable (> 1.5 t/ha)	Vine: AR6 (2.77); AR7 (2.66); NL3 (2.37) Dwarf: AS7 (2.21); AS11 (1.82); AS3 (1.80)	High-yielding genotypes are useful for varietal improvement and cultivation.
Cooking time (min)	25.90 $\pm$ 3.32	20.33–36.67	4.89	4.19	Lower desirable (20–25 min)	Vine: AR2 (36.67) Dwarf: AS10(20.33); AS5 (20.67); ML3 (21.30)	Shorter cooking time indicates softer texture after cooking and reduces fuel consumption.
Crude protein (%)	17.70 $\pm$ 2.20	14.03–23.47	4.74	2.77	Higher desirable (> 20%)	Vine: AR9 (23.47) Dwarf: MR2 (21.43); MR4 (21.03); AS3 (20.50)	High protein genotypes are valuable for nutritional improvement.
Fat (%)	1.93 $\pm$ 0.93	0.6–4.07	9.04	0.58	Higher desirable (1.5–3%)	Vine: AR8 (3.83); AR11 (3.60) Dwarf: MR2 (4.07); AS6 (3.80)	Higher fat may improve energy value and nutritive quality.
Crude fiber (%)	8.19 $\pm$ 3.60	4.75–23.26	2.53	0.68	Moderate desirable (6–9%)	Vine: AR8(7.7); AR6(7.23); AR3(7.01) Dwarf: As6(8.85); ML3(8.21);ML4(7.01)	Suitable fibre content contributes to dietary and health benefits.
Folic acid (μ g)	287.16 $\pm$ 22.43	242.81–336.37	1.30	12.36	Higher desirable (> 300 μ g)	Vine: ML3 (335.92) Dwarf: MR1 (336.37); AS2 (333.21)	High folic acid genotypes are useful for biofortification.

Correlation analysis

The Pearson correlation analysis revealed strong interrelationships among morpho-agronomic and quality traits (Fig. 1). Plant height showed strong positive correlations with branches per plant ($r = 0.75$), pods per plant ($r = 0.83$), days to flowering ($r = 0.89$), days to maturity ($r = 0.89$), and seed yield ($r = 0.73$), indicating that taller and more vigorous genotypes tended to be later maturing and more productive. Branches per plant were positively associated with chlorophyll content index ($r = 0.86$), pods per plant ($r = 0.64$), days to flowering ($r = 0.75$), days to maturity ($r = 0.71$), and seed yield ($r = 0.81$), confirming the importance of branching and canopy vigour in yield formation. Pods per plant also showed strong positive correlations with days to flowering ($r = 0.85$), days to maturity ($r = 0.84$), and seed yield ($r = 0.86$), highlighting pod number as a major yield component. Similar associations of pod number, branching, and plant vigour with seed yield have been reported in common bean by Raturi *et al.*, (2017) and Chahota *et al.*, (2020), supporting their use as indirect selection criteria. Seeds per pod had moderate positive associations with pod length ($r = 0.49$), days to flowering ($r = 0.48$), and yield ($r = 0.52$). In contrast, 100-seed weight was negatively associated with days to flowering ($r = -0.46$), indicating that larger-seeded genotypes tended to flower earlier, while its relationship with yield was weak ($r = 0.18$). Cooking time showed negative correlations with yield ($r = -0.18$), riboflavin ($r = -0.41$), and folic acid

($r = -0.31$), suggesting that faster-cooking genotypes may also possess favourable nutritional quality. Among biochemical traits, crude protein was positively correlated with fat ($r = 0.25$) and riboflavin ($r = 0.29$), whereas riboflavin, niacin, vitamin B₆, and folic acid showed moderate positive intercorrelations ($r = 0.22$ – 0.44), indicating coordinated accumulation of nutritional compounds. Overall, the results suggest that yield improvement in this germplasm can be effectively targeted through plant architecture traits, while nutritional traits remain partly independent, offering scope for simultaneous selection of high-yielding and nutrient-dense genotypes.

Principal component analysis (PCA)

Principal component analysis of 18 morphological and biochemical traits across 40 rajmah genotypes representing two growth habits and five Northeast Indian regions revealed that the first six principal components (eigenvalues > 1) explained 77.6% of the total variation, with PC1 (35.2%) and PC2 (11.2%) contributing most. The scree plot showed a sharp decline after PC2, indicating that the first two components captured the major variability (Fig. 2). PC1 was dominated by plant height, branches per plant, pods per plant, and seed yield, confirming their central role in productivity differentiation, whereas PC2 was primarily associated with 100-seed weight, crude protein, and cooking time, reflecting seed size and compositional traits. Nutritional traits such as fat, fibre, riboflavin, niacin, vitamin B₆, and folic acid loaded moderately on later components, indicating secondary contributions to genotype differentiation. These patterns are consistent with previous multivariate analyses in common bean germplasm (Cichy *et al.*, 2019).




Fig. 1: Pearson correlation heatmap showing associations among 18 morphological and biochemical traits in 40 rajmah genotypes.




Fig. 2: Scree plot showing percentage variance explained by successive principal components.

In the PCA scatter plot based on plant type, vine genotypes clustered mainly along the positive axis of PC1 and were associated with taller plants, greater branching, and higher pod number, whereas dwarf genotypes grouped on the negative side of PC1, reflecting compact growth habit and comparatively lower yield potential. The limited overlap between groups indicated a few intermediate phenotypes (Fig. 3). Clear separation of growth habits in PCA space indicates that plant architecture contributes strongly to overall diversity, which agrees with observations of Andrade *et al.*, (2019) and Girish *et al.*, (2021), who also reported clustering of bean genotypes according to growth form and yield components. The PC1 vs PC3 projection further supported this separation, with vine genotypes showing a wider spread along PC3, suggesting greater within-group variation for




Fig. 3: PCA scatter plot of genotypes based on 18 traits (PC1 vs PC2), color-coded by plant type (Dwarf and Vine).

secondary traits such as maturity, seed quality, or nutritional attributes, while dwarf genotypes remained more compact (Fig. 4). Similarly, the PC2 vs PC3 plot showed partial differentiation between plant types, with vine genotypes tending toward higher positive PC2 scores and dwarf genotypes clustering nearer the origin or negative PC2 region, indicating that PC2 and PC3 captured additional but smaller sources of variation beyond the dominant growth-habit effect represented by PC1 (Fig. 5).

Regional PCA plots indicated that genotypes from Meghalaya and Arunachal Pradesh aligned more closely with yield-related traits on the positive side of PC1, whereas those from Assam and Nagaland




Fig. 4: PCA scatter plot of genotypes based on 18 traits (PC1 vs PC3), color-coded by plant type (Dwarf and Vine).




Fig. 5: PCA scatter plot of rajmah genotypes based on 18 traits (PC2 vs PC3), grouped by plant type (Dwarf and Vine).

clustered toward compact growth and earlier maturity, with Mizoram genotypes occupying intermediate positions (Supplementary Fig. 1). Collectively, these patterns highlight the influence of growth habit and regional adaptation on phenotypic diversity in rajmah and agree with earlier reports in common bean (Long *et al.*, 2020). PCA biplots illustrated that vectors for plant height, branches per plant, seeds per pod, pods per plant, and seed yield aligned with vine-type genotypes, while vectors for 100-seed weight and crude protein, fibre, fat, and vitamin traits aligned in the opposite direction, highlighting contrasting trait emphasis across growth habits (Fig. 6).




Fig. 6: PCA biplot of genotypes and trait vectors coloured by plant type.

Regional variation in standardised mean trait values

The hierarchical heatmap of standardised mean trait values (Fig. 7) displayed distinctive regional patterns: accessions from Arunachal Pradesh recorded the highest values for yield-related traits (height, branching, pods per plant and seed yield), Nagaland genotypes showed moderate yield but heavier seeds and longer cooking time, Mizoram genotypes were richer in nutritional traits (crude protein, fat, vitamins) and accessions from Assam and Meghalaya combined early maturity with lower yield traits. These results corroborate the PCA findings and confirm that regional origin strongly influences the trade-off between yield potential and nutritional composition in this germplasm collection.




Fig. 7: Hierarchical heatmap showing standardized mean values of 18 morphological and biochemical traits across rajmah genotypes from five northeastern regions.

Cluster analysis and regional grouping of genotypes

Hierarchical cluster analysis based on PCA scores (Fig. 8) grouped the 40 rajmah genotypes into two major clusters broadly corresponding to dwarf and vine growth habits, with further sub-clusters reflecting regional origins. Accessions from Arunachal Pradesh

and Meghalaya formed one main cluster characterised by greater plant height and pod number, while accessions from Assam and Nagaland grouped separately, displaying compact growth and moderate yield. Mizoram accessions occupied intermediate positions between the two main clusters. This clustering confirms that growth habit is the principal determinant of phenotypic diversity in this collection, with regional adaptation contributing to intra-group variation. Similar clustering patterns based on morphological divergence have been reported in common bean germplasm from China, Ethiopia, and Brazil (Gelaw, 2017; Long et al., 2020; de Paula et al., 2024), supporting the robustness of the present findings.




Fig. 8: Hierarchical dendrogram showing clustering of 40 rajmah genotypes by region based on multivariate trait data.

Breeding implications

The findings suggest that selection for improvement of rajmah germplasm in Northeast India can be effectively guided by a concise set of traits like plant height, branches per plant, pods per plant, 100-seed weight and crude protein. The clear separation of vine and dwarf types with distinct trait profiles indicates the potential for targeted crossing to combine high-yield and high-nutritional quality. In particular, vine-type accessions with strong yield component traits and dwarf-type accessions with superior seed quality can be selected as parents for developing genotypes combining both agronomic performance and nutritional value. The strong region-based differentiation also suggests the value of exploiting local adaptation in breeding strategies. Many of the underlying associations (e.g., yield component trait correlations) have been emphasized in recent meta-QTL and genomic studies of common bean (Arriagada et al., 2023).

Conclusion

The present study demonstrated substantial morpho-biochemical diversity among rajmah genotypes

collected from Northeast India. Significant variation across agronomic, yield, and nutritional traits indicates strong potential for selection and genetic improvement. Multivariate analyses consistently revealed growth habit as the principal source of variation, with vine genotypes excelling in plant vigour and yield-related traits, while dwarf genotypes possessed superior seed size and several nutritional attributes. Regional grouping further suggested the influence of local adaptation on phenotypic expression. Traits such as plant height, branches per plant, pods per plant, 100-seed weight, and crude protein emerged as important selection criteria for breeding. The identified contrasting genotypes provide valuable parental resources for developing high-yielding, nutritionally enriched, and regionally adapted rajmah cultivars for the North-Eastern Hill Region and similar agro-ecologies.

Acknowledgements

The authors are grateful to the Assam Agricultural University, Jorhat, for providing the necessary facilities to conduct this study. Support received from the Department of Plant Breeding and Genetics and the Directorate of Research (Agri), AAU, during field and laboratory work is duly acknowledged.

References


- Ahmad KMS (2018) Genetic diversity of common bean (*Phaseolus vulgaris* L.) cultivars from different origins revealed by microsatellite markers. *J. Adv. Biol. Biotechnol.* 17(4): 1-9.
- Andrade RA, LM Lima and EV Rodrigues (2019) Genetic diversity and structure in common bean based on morpho-agronomic traits. *Genet. Resour. Crop Evol.* 66(5): 1107-1121.
- AOAC (2016) *Official Methods of Analysis*, 20th edn. Association of Official Analytical Chemists International, Gaithersburg, Maryland, USA.
- Arriagada O, B Arévalo, RA Cabeza, B Carrasco and AR Schwember (2023) Meta-QTL analysis for yield components in common bean (*Phaseolus vulgaris* L.). *Plants* 12(1): 117.
- Banoo A, N Ahmed, SH Khan and SM Zargar (2020) Genetic diversity assessment in common bean (*Phaseolus vulgaris* L.) germplasm using agro-morphological traits. *Legume Res.* 43(5): 674-680.
- Beebe S, AV González and J Rengifo (2010) Research on trace minerals in the common bean. *Food Nutr Bull* 31(1 Suppl 1): 67-70.
- Blair MW, LF González, PM Kimani and L Butare (2010) Genetic diversity, inter-gene pool introgression and nutritional quality of common bean (*Phaseolus vulgaris* L.) from Central Africa. *Theor. Appl. Genet.* 121(2): 237-248.
- Broughton WJ, G Hernández, M Blair, S Beebe, P Gepts and J Vanderleyden (2003) Beans (*Phaseolus* spp.)-model food legumes. *Plant Soil* 252(1): 55-128.
- Celmeli T, H Sari, H Canci, D Sari, A Adak, T Eker and C Toker (2018) Nutritional content of common bean (*Phaseolus vulgaris* L.) landraces in comparison to modern varieties. *Agronomy* 8(9): 166.
- Chahota RK, TR Sharma and SK Sharma (2020) Genetic variability and trait association studies in rajmah (*Phaseolus vulgaris* L.) under mid-hill conditions of Himachal Pradesh. *Legume Res.* 43(6): 773-778.
- Cichy KA, JA Wiesinger, M Berry, S Nchimbi-Msolla, D Fourie, TG Porch, D Ambechew and PN Miklas (2019) Role of genotype and environment in determining cooking time of dry beans (*Phaseolus vulgaris* L.). *Legume Sci.* 1: e13.
- de Paula E, RN Almeida, TO Santos, JDN Souza Neto, EMS Riva-Souza, SCP Posse, MN Souza, AFF Madella de Oliveira, AC Santos Júnior and JO Santos (2024) Genetic diversity of common bean (*Phaseolus vulgaris* L.) landraces based on morphological traits and molecular markers. *Plants* 13(18): 2584.
- Dutta M, A Saikia, N Gogoi and JD Nath (2015) Genetic variability studies in pole-type rajmah (*Phaseolus vulgaris* L.) genotypes of north-eastern hill region of India. *Legume Res.* 38(5): 631-635.
- Fisher RA (1935) *The Design of Experiments*, 1st edn. Oliver and Boyd, Edinburgh, Scotland, 252 p.
- Friendly M (2002) Corrgrams; exploratory displays for correlation matrices. *The American Statistician* 56: 316-324.
- Gelaw YM (2017) Genetic diversity in speckled bean (*Phaseolus vulgaris* L.) genotypes in Ethiopia. *J. Plant Breed. Crop. Sci.* 9(11): 200-207.
- Girish M, G Muniswamy and S Ramesh (2021) Multivariate analysis for genetic divergence in rajmah (*Phaseolus vulgaris* L.) genotypes. *J. Hort. Sci.* 16(1): 47-54.
- Gomez KA and AA Gomez (1984) *Statistical Procedures for Agricultural Research*, 2nd edn. John Wiley and Sons, New York, USA, 680 p.
- Gu Z (2016) Complex heatmaps reveal patterns and correlations in multidimensional genomic data. *Bioinformatics* 32(18): 2847-2849.
- Indian Farming (2017) Rajmah; an important pulse crop for hills. *Indian Farming* 67(9): 17-19.
- Jain A, P Sharma and V Kumar (2019) Morphological and biochemical characterization of common bean genotypes. *Indian J. Genet. Plant Breed.* 79(4): 737-744.
- Jolliffe IT (2002) *Principal Component Analysis*, 2nd edn. Springer-Verlag, New York, USA, 487 p.
- Khan S, A Singh and M Sharma (2021) Nutritional and biochemical evaluation of common bean genotypes for protein and micronutrient content. *Legume Res.* 44(6): 673-679.
- Kwak M and P Gepts (2009) Structure of genetic diversity in the two major gene pools of common bean (*Phaseolus vulgaris* L.). *Theor. Appl. Genet.* 118: 979-992.
- Long J, J Zhang, X Zhang, J Wu, H Chen, P Wang, Q Wang and C Du (2020) Genetic diversity of common bean germplasm resources based on morphological traits. *Front. Genet.* 11: 697.
- Lowry OH, NJ Rosebrough, AL Farr and RJ Randall (1951) Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193(1): 265-275.
- Mwale VM, JM Bokosi and MB Kwapata (2020) Morphological diversity of common bean landraces from Malawi. *Int. J. Plant Breed. Genet.* 14(1): 1-9.

- Oliveira JS, MF Pereira and PA de Carvalho (2019) Genetic diversity of common bean based on morphological and biochemical traits. *PesqAgropecu Bras* 54: e00013.
- R Core Team (2024) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna.
- Rai N, M Rai and R Singh (2020) Characterization of common bean germplasm under North-Eastern hill conditions. *Indian J. Agric.Sci.* 90(2): 378-383.
- Raturi A, R Sharma and A Sharma (2017) Variability and correlation studies in common bean (*Phaseolus vulgaris* L.) genotypes. *Legume Res.* 40(3): 464-469.
- Shah AH, I Khaliq, G Bashir and S Jannat (2020) Genetic diversity in common bean germplasm of Jammu & Kashmir. *Indian J. Agric. Res.* 54(5): 629-634.
- Singh S, M Sood and A Sharma (2014) Genetic divergence and variability studies in rajmah (*Phaseolus vulgaris* L.). *Legume Res.* 37(4): 439-444.
- Singh Y, SK Yadav and S Sharma (2021) Multivariate genetic analysis of common bean genotypes. *Plant Genet. Resour.* 19(5): 432-440.
- Sneath PHA and RR Sokal (1973) *Numerical Taxonomy: The Principles and Practice of Numerical Classification*. W.H. Freeman and Company, San Francisco, USA, 573 p.
- Wickham H (2016) *ggplot2: Elegant Graphics for Data Analysis*, 2nd edn. Springer International Publishing, Cham, Switzerland, 260 p.

Supplementary Table 1: Genotype designation, plant type, and state of origin of the rajmah genotypes evaluated

S. No.	Genotype designation	Plant type	State of origin
1.	ML1	Dwarf	Meghalaya
2.	ML2	Dwarf	Meghalaya
3.	ML3	Dwarf	Meghalaya
4.	ML4	Vine	Meghalaya
5.	ML5	Dwarf	Meghalaya
6.	NL1	Vine	Nagaland
7.	NL2	Vine	Nagaland
8.	NL3	Vine	Nagaland
9.	NL4	Dwarf	Nagaland
10.	NL5	Dwarf	Nagaland
11.	AR1	Dwarf	Arunachal Pradesh
12.	AR2	Vine	Arunachal Pradesh
13.	AR3	Dwarf	Arunachal Pradesh
14.	AR4	Vine	Arunachal Pradesh
15.	AR5	Vine	Arunachal Pradesh
16.	AR6	Vine	Arunachal Pradesh
17.	AR7	Vine	Arunachal Pradesh
18.	AR8	Vine	Arunachal Pradesh
19.	AR9	Vine	Arunachal Pradesh
20.	AR10	Vine	Arunachal Pradesh
21.	AR11	Vine	Arunachal Pradesh
22.	AR12	Dwarf	Arunachal Pradesh
23.	AR13	Vine	Arunachal Pradesh
24.	MR1	Dwarf	Mizoram
25.	MR2	Dwarf	Mizoram
26.	MR3	Vine	Mizoram

S. No.	Genotype designation	Plant type	State of origin
27.	MR4	Dwarf	Mizoram
28.	MR5	Vine	Mizoram
29.	AS1	Dwarf	Assam
30.	AS2	Dwarf	Assam
31.	AS3	Dwarf	Assam
32.	AS4	Dwarf	Assam
33.	AS5	Dwarf	Assam
34.	AS6	Dwarf	Assam
35.	AS7	Dwarf	Assam
36.	AS8	Dwarf	Assam
37.	AS9	Dwarf	Assam
38.	AS10	Dwarf	Assam
39.	AS11	Dwarf	Assam
40.	AS12	Dwarf	Assam

**Supplementary Fig. 1:** PCA scatter plot of genotypes grouped by region of origin.

RESEARCH ARTICLE

Fatty Acid Composition of Seeds of Three Native *Garcinia* Species from Andaman and Nicobar Islands, India.

Pooja Bohra^{1*} and Ajit Arun Waman¹

Abstract

Garcinia L. is an important genus, comprising of species revered for their edible fruits and exploited as sources of industrially valuable compounds. Andaman and Nicobar Islands in the Bay of Bengal have considerable diversity for the genus and a number of aspects are yet to be studied in most of the species. To understand the potential of native species of these islands, the fatty acid composition of seed fat of three *Garcinia* species, viz. *G. kurzii* Pierre, *G. cowa* Roxb. ex Choisy and *G. dulcis* (Roxb.) Kurz was determined using GC-MS-MS analysis. Results revealed that seeds of *G. cowa* and *G. dulcis* were rich in unsaturated fatty acids, while seeds of *G. kurzii* had higher saturated fatty acids content. Oleic acid (44.24%) and Stearic acid (34.94%) were dominant in *G. cowa*, Oleic acid (39.82%) and Palmitic acid (33.48%) in *G. dulcis*, whereas Stearic acid (41.34%) and Oleic acid (23.27%) were major fatty acids in *G. kurzii*. Arachidic, Eicosenoic, Linoleic and Palmitoleic acids content were also found in varied quantities. Thus, the present report opens up avenues for the utilisation of seeds of the studied species for different industrial purposes.

Keywords: *Garcinia cowa*, *Garcinia dulcis*, *Garcinia kurzii*, GC-MS

¹Division of Horticulture and Crop Improvement, ICAR-Central Island Agricultural Research Institute, Sri Vijaya Puram, Andaman and Nicobar Islands, 744105, India.

***Authors for correspondence:**

poojabohra24@gmail.com

Received: 22/10/2025

Revised: 23/03/2026

Accepted: 15/04/2026

How to cite this article: Bohra P and AA Waman (2026) Fatty Acid Composition of Seeds of Three Native *Garcinia* Species from Andaman and Nicobar Islands, India. *Indian J. Plant Genet. Resour.* 39(2): 53-57.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.178052>

Introduction

The genus *Garcinia* L. is one of the important genera of the Clusiaceae family, species of which have been valued for their multifaceted uses, including as fresh fruits, natural colourants, acidulants, functional foods, saturated and unsaturated fats, varnishes, etc. (Baruah *et al.*, 2012; Parthasarathy *et al.*, 2013; Bohra *et al.*, 2020). Identification of bioactive compounds of high pharmaceutical and industrial significance, such as hydroxycitric acid, flavonoids, xanthenes, benzophenones, biphenyls, acyl phloroglucinols, etc., has further increased their utilization potential (Aravind *et al.*, 2016).

As per the Plants of the World Online database, ca. 414 accepted species of the genus *Garcinia* have been reported so far (<https://powo.science.kew.org>). Of these species, a few are being deliberately cultivated, and a larger number of it is yet to be brought into mainstream cultivation due to a lack of scientific studies, especially on the phytochemical potential of their plant parts (Aravind *et al.*, 2016). Most of the reported *Garcinia* species produce edible fruits; yet, considering the sour to highly sour taste, most of these species may not be suitable for direct consumption (Te-chato, 2007; Baruah *et al.*, 2012; Ellina *et al.*, 2015). Under such circumstances, identification of their non-fruit uses could be beneficial.

The Andaman and Nicobar Islands are one of the important agro-biodiversity regions representing the flora of Indo-Burma

and Sundaland (Singh *et al.*, 2016). This agrobiodiversity holds potential for the identification of novel sources of medicines, industrial compounds, genes, foods and novel crops, if studied systematically. However, when compared with the other biodiverse regions of the country, limited research work has been done on the species distributed in these islands, especially on the genus *Garcinia* (Parthasarathy *et al.*, 2013; Bohra *et al.*, 2021a). Hence, research on various aspects of these potential species is envisaged.

Seeds in fruit-bearing species have been the default material of perpetuation naturally and their utilization from an economic point of view is largely negligible. In fact, they largely remain a waste resource in a number of cases (Abraham *et al.*, 2006). Heavy bearing capacity of various *Garcinia* species and presence of multiple seeds in each fruit in most of the species makes their seeds a potential resource for utilization (Te-chato, 2007; Ellina *et al.*, 2015; Shameer *et al.*, 2016; Bohra and Waman, 2019). Traditionally, in the Western Ghats, seeds of species such as *G. indica* and *G. gummi-gutta* have been utilized as sources of fats for cosmetic and medicinal purposes (Joshi, 2005; Baliga *et al.*, 2011). However, such traditional knowledge in other *Garcinia* species is largely unavailable.

Recently, *Garcinia* fat is being considered as an ingredient for various industries, including a substitute for cocoa butter, personal care products, pharmaceuticals, biofuels, edible oils etc. (Parthasarathy and Nandakishore, 2014; Ananthakrishnan and Ramesh kumar, 2016). Studies on fatty acid content in Indian *Garcinia* species are limited (Parthasarathy and Nandakishore, 2014; Patil *et al.*, 2016). Considering this, the present study was undertaken to understand the composition of seed fat of three native *Garcinia* species of the Andaman and Nicobar Islands of the Bay of Bengal.

Materials and Methods

Field surveys were conducted in different parts of the Andaman group of Islands, India and ripe fruits of



Fig. 1: Fruit of *Garcinia kurzii*.



Fig. 2: Fruit of *Garcinia cowa*.



Fig. 3: Fruit of *Garcinia dulcis*.

Garcinia kurzii (Ghat kau phal, Fig. 1), *G. cowa* (Kau phal, Fig. 2), *G. dulcis* (Tok phal, Fig. 3) were collected from the identified locations of Middle Andaman Island. Fruits were brought to the ICAR-Central Island Agricultural Research Institute, Sri Vijaya Puram and used for the studies. Morphological parameters, including fruit colour, fruit surface, fruit length (cm), fruit width (cm), peel thickness (mm) and number of seeds per fruit were recorded using the standard procedures. Seeds were extracted, the adhering pulp was washed, and seeds were air-dried to facilitate the removal of the seed coat. Such seeds were used for measurement of seed length (cm), seed width (cm) and seed thickness (cm).

In order to carry out fatty acid profiling, air-dried seeds were immediately transported to the ICAR-Indian Institute of Horticultural Research, Hessarghatta, Bengaluru, India. The procedure for lipid fractionation, esterification and subsequent Gas chromatography-mass spectrometry (GC-MS/MS) described elsewhere (Bohra *et al.*, 2021a) was followed. For identification of fatty acid compounds, relative retention times of FAME peaks were compared with those of reference standards (Sigma-Aldrich, USA) along with spectral comparison with reference (Wiley and NIST-2007). Analysis was carried out in duplicate, and data were presented as mean $\pm$ standard error of the mean.

Results and Discussion

General description

G. kurzii is an endemic species distributed in the Andaman and Nicobar Islands (Nissar *et al.*, 2025). Fruits were 3.2-3.8 cm $\times$ 2.9-3.2 cm in dimension, purple to black with an inedible rind of 1.5 to 2.5 mm thick. Each

Table 1: Basic information about the studied *Garcinia* species

Parameter	<i>G. kurzii</i>	<i>G. cowa</i>	<i>G. dulcis</i>
Edible portion	Pulp	Peel and pulp, young leaves	Peel and pulp
Fruit colour	Purple to black	Orange	Yellow
Fruit surface	Smooth	Ridged	Smooth
Fruit length (cm)	3.2 - 3.8	3.5 - 4.6	4.5 - 5.6
Fruit width (cm)	2.9 - 3.2	3.8 - 5.8	5.5 - 6.4
Fruit weight (g)	13.8 - 16.0	40.1 - 66.0	68.5 - 111.2
Rind thickness (mm)	1.5 - 2.5	4.0 - 6.5	0.8 - 1.4
No. of seeds per fruit	1 - 4	1 - 6	3 - 5
Seed length (cm)	1.7 - 2.1	1.8 - 2.1	1.6 - 2.7
Seed width (cm)	0.7 - 0.9	1.1 - 1.5	1.4 - 1.7
Seed thickness (cm)	0.6 - 0.8	0.7 - 1.0	0.8 - 1.4

fruit weighed between 13.8 and 16.0 g. White, sweetish-sour pulp is eaten by local settlers and the Nicobarese tribe of these islands. Fruits had four segments with one seed each measuring 1.7-2.1 cm long, 0.7-0.9 cm wide and 0.6-0.8 cm thick (Table 1).

G. cowa had orange fruits with a ridged surface. Fruits are dried and used as an acidulant, while orange coloured pulp is eaten fresh. Further, green fruits are also used against food poisoning in Northeastern India (Hazarika *et al.*, 2016). In the present study, fruits weighed between 40.1 and 66.0 g, while dimensions were 3.5-4.6 cm × 3.8-5.8 cm with 4.0 to 6.5 mm thick rind. An earlier study from Bangladesh (Hossain *et al.*, 2023) has suggested comparatively smaller-sized fruits (18.26 to 31.29 g), whereas fruits of 3-6 cm size and 40-120 g weight have been reported from China (Liu *et al.*, 2005). In the present study, each fruit had 1 to 6 seeds with dimensions of 1.8-2.1 cm × 1.1-1.5 cm × 0.7-1.0 cm; while Hossain *et al.*, (2023) reported the presence of

5.00 to 7.33 seeds per fruit.

G. dulcis is a large fruited species, the fruits of which are suitable for preparation of value-added products such as jams and preserves; whereas bark exudates are used in dyeing (Maheshwari, 1964). A recent study has shown antioxidant and antielastase activities of this species (Ambarwati *et al.*, 2024). Fruits of *G. dulcis* were of 4.5-5.6 cm × 5.5-6.4 cm dimensions. Fruits were the heaviest among all the studied species and weighed between 68.5 and 111.2 g. Fruits had a smooth yellow surface with yellowish, sweetish, sour edible pulp. Fruits had 3-5 seeds each of 1.6-2.7 cm length, 1.4-1.7 cm width and 0.8-1.4 cm thickness.

Fatty acid composition

Seed fatty acid profiling suggested the presence of seven compounds, irrespective of the species (Table 2). These compounds contributed to 99.12% of total composition in *G. cowa*, 98.09% in *G. dulcis* and 97.81%

Table 2: Fatty acid composition of three *Garcinia* species

Fatty acid	C number	Class of compound	Peak area (%)		
			<i>G. kurzii</i>	<i>G. cowa</i>	<i>G. dulcis</i>
Palmitic acid	16:0	Saturated	11.08 ± 0.284	10.52 ± 0.051	33.48 ± 1.501
Stearic acid	18:0	Saturated	41.34 ± 0.470	34.94 ± 1.206	9.00 ± 1.081
Arachidic acid	20:0	Saturated	1.12 ± 0.794	0.45 ± 0.170	1.41 ± 0.608
		Σ Saturated fatty acid	53.54	45.91	43.89
Palmitoleic acid	16:1	Monounsaturated	0.32 ± 0.071	0.20 ± 0.062	7.20 ± 0.270
Oleic acid	18:1	Monounsaturated	23.27 ± 0.771	44.24 ± 1.606	39.82 ± 2.028
Eicosenoic acid	20:1	Monounsaturated	1.56 ± 0.933	1.13 ± 0.202	2.70 ± 0.751
		Σ Monounsaturated fatty acid	25.15	45.57	49.72
Linoleic acid	18:2	Polyunsaturated	19.12 ± 1.638	7.64 ± 0.072	4.48 ± 1.182
		Σ Polyunsaturated fatty acid	19.12	7.64	4.48
		Σ Unsaturated fatty acid	44.27	53.21	54.20
		Total	97.81	99.12	98.09

in *G. kurzii*. Of these seven compounds, three were saturated fatty acids (SFA), three were monounsaturated fatty acids (MUFA), and one was polyunsaturated fatty acid (PUFA). Relative abundance of these compounds varied among the species studied. In *G. cowa* and *G. dulcis*, proportion of unsaturated fats was higher (53.21% and 54.20%, respectively) than SFAs. Higher proportions of unsaturated fatty acids have been reported in *G. multiflora* (84.19%), *G. xanthochymus* (65.79%), *G. gummi-gutta* (59.64%) and *G. andamanica* (53.24%), as well (Manohar *et al.*, 2014; Patil *et al.*, 2016; Phuong *et al.*, 2018; Bohra *et al.*, 2021a), which support the findings of the present study. On the other hand, in *G. kurzii*, higher amounts of saturated fatty acids (53.54%) were present than the unsaturated ones. *G. dhanikhariensis*, the endemic species of the Andaman Islands has recently been reported to have higher proportions of SFAs (52.69%) apart from exotic *G. hanburyi* (53.11%) and *G. gaudichaudii* (61.15%) (Phuong *et al.*, 2018; Bohra *et al.*, 2021b). The composition of unsaturated fatty acids showed higher quantities of MUFAs than that of PUFAs. This observation was in line with the earlier reports in *Garcinia* species (Phuong *et al.*, 2018; Bohra *et al.*, 2021a; Bohra *et al.*, 2021b).

PUFAs are considered healthier compounds, and the highest amounts of PUFAs were reported in *G. kurzii* (19.12%), which was followed by *G. cowa* (7.64%) and the lowest (4.48%) in *G. dulcis*. As low as 25.15% MUFAs were observed in *G. kurzii*, while they were 49.72% in *G. dulcis*. Proven beneficial effects and health-promoting properties of the unsaturated fatty acids have been reported by earlier researchers (Ajayi *et al.*, 2007; Patil *et al.*, 2016). Hence, *G. cowa* and *G. dulcis*, with higher quantities of such compounds, could be utilized for blending purposes, in line with *G. gummi-gutta* that has been conventionally utilized for culinary preparations in Kerala state of India (Abraham *et al.*, 2006).

In case of *G. cowa*, Oleic acid was found to be the most abundant compound (44.24%), followed by Stearic acid (34.94%), Palmitic acid (10.52%), Linoleic acid (7.64%) and Eicosenoic acid (1.13%), while Arachidic acid and Palmitoleic acid were present in minute quantities. Similarly, Oleic acid was the most dominant compound (39.82%) in the seeds of *G. dulcis*. Palmitic acid (33.48%), Stearic acid (9.00%), Palmitoleic acid (7.20%), Linoleic acid (4.48%), Eicosenoic acid (2.70%) and Arachidic acid (1.41) were the other compounds present in *G. dulcis*.

Several reports have described the presence of Oleic acid in *Garcinia* seeds, which is an important constituent from a health point of view (Grund,

1989). *G. andamanica*, an endemic species of these islands, has been reported to contain 44.35% Oleic acid in its seed fat (Bohra *et al.*, 2021a), whereas another endemic species *G. dhanikhariensis* had 42.40% of it (Bohra *et al.*, 2021b). Higher quantities of Oleic acid in the presently studied species could be of great practical utility as healthy oil for commercial utilization as already advocated for the Vietnamese species (Phuong *et al.*, 2018). *Camellia oleifera*- an oleic acid-rich species has been reported to have high moisturizing properties (Sang *et al.*, 2023), useful for hair care products.

Stearic acid and oleic acid were the dominant compounds in *G. kurzii* (41.34% and 23.27%, respectively). Linoleic acid (19.12%), Palmitic acid (11.08%) and Eicosenoic acid (1.56%) were the other compounds observed in *G. kurzii*. These findings are in accordance with the reports on *G. dhanikhariensis*, *G. gaudichaudii* and *G. hanburyi*, with 43.62%, 50.12% and 58.87%, of stearic acid, respectively (Phuong *et al.*, 2018). Saturated fats such as Stearic acid and Palmitic acid are known to have surfactant, softening and reparative effects and hence, are used as key ingredients in soaps, shaving creams, and related cosmetic formulations (Parthasarathy and Nandakishore, 2014). *G. kurzii* and *G. dulcis* could also be promising alternative sources for these industries. *G. dulcis* in the present study exhibited a trend of Oleic acid > Palmitic acid > Palmitoleic acid, which was similar to *G. xanthochymus* (Manohar *et al.*, 2014).

Conclusion

In conclusion, fatty acid composition of seeds varied with species, in which *G. cowa* and *G. dulcis* had higher content of unsaturated fatty acids, while *G. kurzii* showed higher saturated fatty acids. Oleic acid and Stearic acid (*G. cowa*), Oleic acid and Palmitic acid (*G. dulcis*), and Stearic acid and Oleic acid (*G. kurzii*) were dominant compounds apart from other minor constituents. Considering the natural diversity present in different parts of the Andaman and Nicobar Islands, intra-specific variability needs to be screened for the identification of superior types.

Acknowledgements

Financial support to the first author from ICAR, New Delhi in the form of inhouse project is gratefully acknowledged. The authors are also grateful to the Director of the Institute for providing necessary facilities, SAIF, ICAR-IIHR, Bengaluru, for help in chromatography and Dr PS Shameer for confirming the botanical identity of samples.

References

- Abraham Z, SK Malik, GE Rao, SL Narayanan and S Biju (2006) Collection and characterisation of Malabartamarind [*Garcinia cambogia* (Gaertn.) Desr]. *Genet. Resour. Crop Evol.* 53: 401-406. <https://doi.org/10.1007/s10722-004-0584-y>.
- Ajayi IA, RA Oderinde, BO Ogunkoya, A Egunyomi and VO Taiwo (2007) Chemical analysis and preliminary toxicological evaluation of *Garcinia mangostana* seeds and seed oil. *Food Chem.* 101:999-1000.
- Ambarwati NSS, NT Sukma, Y Desmiaty, A Auliya, S Budi, M Arifuddin and I Ahmad (2024) *Garcinia dulcis* and *Garcinia forbesii* King fruit peel extract; secondary metabolite composition, antioxidant, and elastase inhibitory activity evaluation. *J. Adv. Pharm. Technol. Res.* 15(1): 8-12. doi: 10.4103/japtr.japtr_344_23.
- Ananthakrishnan R and KB Rameshkumar (2016) Phytochemicals and bioactivities of *Garcinia indica* (Thouars) Choisy; a review. In: KB Rameshkumar (ed) *Diversity of Garcinia species in the Western Ghats: Phytochemical Perspective*. JNTBGRI, Thiruvananthapuram, India, pp 142-150.
- Aravind APA, NM Lekshmi and KB Rameshkumar (2016) Structural diversity of secondary metabolites in *Garcinia* species. In: KB Rameshkumar (ed) *Diversity of Garcinia species in the Western Ghats: Phytochemical Perspective*. JNTBGRI, Thiruvananthapuram, India, pp 19-75.
- Baliga MS, HP Bhat, RJ Pai, R Bloor and PL Palatty (2011) The chemistry and medicinal uses of the underutilized Indian fruit tree *Garcinia indica* Choisy (kokum); a review. *Food Res. Intl.* 44: 1790-1799.
- Baruah S and SK Borthakur (2012) Studies on morphology and ethnobotany of six species of *Garcinia* L. (Clusiaceae) found in the Brahmaputra valley, Assam, India. *J. Nat. Prod. Plant Resour.* 2(3): 389-396.
- Bohra P and AA Waman (2019) Morphological and biochemical studies in *Garcinia gummi-gutta* (L.) Roxb. *Erwerbs-Obstbau* 61: 217-223. <https://doi.org/10.1007/s10341-019-00419-3>.
- Bohra P, AA Waman and RK Devi (2021b) Seed fatty acid composition and germination studies in *Garcinia dhanikhariensis* SK Srivastava (Clusiaceae); a novel tropical fruit species from Bay Islands, India. *Intl. J. Fruit Sci.* 21(1): 970-978. <https://doi.org/10.1080/15538362.2021.1951921>.
- Bohra P, AA Waman and NA Giri (2021a) *Garcinia andamanica* King (Clusiaceae); an important horticultural genetic resource from Indian islands. *Genet. Resour. Crop Evol.* 68(4): 1675-1689. doi: 10.1007/s10722-021-01136-1.
- Bohra P and AA Waman (2020) Morphological and biochemical studies in *Garcinia dhanikhariensis*; a potential natural colorant from the Bay Islands, India. In: Book of Abstracts of International Web-Conference on New Trends in Agriculture, Environmental & Biological Sciences for Inclusive Development, Rampur, India, 21-22 June 2020, pp 101-102.
- Ellina M, HS Edison and Noflindawati (2015) Preliminary studies on morphological and molecular characterization of wild *Garcinia* in Sijunjung, West Sumatra. *Indian J. Plant Genet. Resour.* 28(1): 31-38. DOI 10.5958/0976-1926.2015.00004.2
- Grundy SM (1989) Monounsaturated fatty acids and cholesterol metabolism; implication for dietary recommendations. *J Nutr.* 119: 529-533.
- Hazarika TK, M Siljea, D Mandal, K Upadhyaya, BA Nautiyal and AC Shukla (2016) Underutilized and unexploited fruits of Indo-Burma hot spot, Meghalaya, north-east India; ethno-medicinal evaluation, socio-economic importance and conservation strategies. *Genet. Resour. Crop Evol.* 63: 289-304. DOI 10.1007/s10722-015-0248-0.
- Hossain MM, MA Rahim, Y Afsana, AK Basunia, MR Haque and MJ Mohona (2023) Morphological characterization and principal component analysis of cowa (*Garcinia cowa* Roxb.) germplasm. *Intl. J. Minor Fruits Med. Arom. Plants* 9(2): 141-148.
- Joshi GD (2005) Traditional methods of kokum (*Garcinia indica* Choisy) fruits processing. In: *Proceedings and Compendium: Second National Conference on Kokum*, Taleigao, Goa University.
- Liu Y, Q Yu-Ping, Z Ling and J Chen (2005) Dormancy breaking and storage behaviour of *Garcinia cowa* Roxb. (Guttiferae) seeds; implications for ecological function and germplasm conservation. *J. Integr. Plant Biol.* 47(1): 38-49. <https://doi.org/10.1111/j.1744-7909.2005.00010.x>.
- Maheshwari JK (1964) Taxonomic study of Indian Guttiferae III. The genus *Garcinia* L. s.l. *Bull. Bot. Surv. India* 6: 107-135.
- Manohar SH, PM Naik, LM Patil, I Karikatti and HN Murthy (2014) Chemical composition of *Garcinia xanthochymus* seeds, seed oil, and evaluation of its antimicrobial and antioxidant activity. *J. Herbs Spices Med. Plants* 20: 148-155. DOI: 10.1080/10496475.2013.847886.
- Nissar VAM, K Pradheep, KJ John, I Jaisankar, BA Jerard, TPM Azharudheen, IPV Kumar, KV Saji, TE Sheeja, CM Harinarayanan and PS Shameer (2025) Reinstatement of *Garcinia kurzii*, an endemic species from Andaman and Nicobar Islands of India and notes on its distribution, conservation and economic importance. *Genet. Resour. Crop Evol.* <https://doi.org/10.1007/s10722-025-02460-6>.
- Parthasarathy U and OP Nandakishore (2014) Morphological characterisation of some important Indian *Garcinia* species. *Dataset Papers Sci.* <http://dx.doi.org/10.1155/2014/823705>
- Parthasarathy U, K Nirmal Babu, R Senthil Kumar, GR Ashis, S Mohan and VA Parthasarathy (2013) Diversity of Indian *Garcinia*; a medicinally important spice crop in India. *Acta Hortic.* 979: 467-476.
- Patil MM, AM Muhammed and KA Anu-Appaiah (2016) Lipids and fatty acid profiling of major Indian *Garcinia* fruit; a comparative study and its nutritional impact. *J. Am. Oil Chem. Soc.* 93: 823-836.
- Phuong DL, TT Nguyen, QL Pham, MQ Pham, TTT Tran, THM Pham, CK Ping and DT Tran (2018) Fatty acid, tocopherol, sterol compositions and antioxidant activity of three *Garcinia* seed oils. *Rec. Nat. Prod.* 12: 323-331.
- Sang SH, AA Gabriel, BL Kai, L Siew-Keah, K Jing-Wen, L Sue-Kei, Y Jessica-Ai-Lyn, ST Ching and C Yik-Ling (2023) Natural alternatives from your garden for hair care; revisiting the benefits of tropical herbs. *Heliyon* 9(11): e21876. <https://doi.org/10.1016/j.heliyon.2023.e21876>.
- Shameer PS, KB Rameshkumar and N Mohanan (2016) Diversity of *Garcinia* species in the Western Ghats. In: KB Rameshkumar (ed) *Diversity of Garcinia species in the Western Ghats: Phytochemical Perspective*, JNTBGRI, Thiruvananthapuram, India, pp 1-18.
- Singh S, AA Waman, P Bohra, RK Gautam and S Dam Roy (2016) Conservation and sustainable utilization of horticultural biodiversity in tropical Andaman and Nicobar Islands, India. *Genet. Resour. Crop Evol.* 63: 1431-1445. <https://doi.org/10.1007/s10722-016-0445-5>.
- Te-chato S (2007) Floral and fruit morphology of some species in *Garcinia* spp. *Songklanakarin J. Sci. Technol.* 29: 245-252.

RESEARCH ARTICLE

Evaluation of Barley (*Hordeum vulgare* L.) Germplasm for Heat Tolerance Using Germination Indices and Seedling Traits

Mohmad Amin Bhagat¹, Navjyot Kaur^{2*}, Prinka Goyal¹ and Simarjit Kaur³

Abstract

Heat stress is a major constraint in barley production. In the present study, 92 genotypes of barley along with 8 checks were evaluated for heat tolerance at the seedling stage under *in vitro* conditions at three temperature regimes viz., 25°C (control), 30°C and 35°C. The genotypes were assessed for germination indices viz., germination percentage, speed of germination and seedling traits viz., root length, shoot length, seedling vigour index I and II. Tolerant genotypes exhibited higher germination percentage, root length, shoot length, seedling vigour index I and II at 30 and 35°C. Based on the hierarchical cluster analysis, nine heat-tolerant genotypes viz., BL 1515, BL 1729, BL 1780, BL 1784, BL 1786, BL 1792, BL 1794, BL 1797 and IBYT-E24 and three heat-sensitive genotypes viz., BL 1723, IBON-23 and IBYT-E15 were identified. These heat-tolerant genotypes can be used as potential donors for developing heat-tolerant barley varieties in future breeding programs.

Keywords: Germination, Heat stress, Screening, Seedling vigour, Tolerant genotypes

¹Department of Botany, Punjab Agricultural University, Ludhiana, Punjab, 141004, India.

²Office of the Director (Seeds), Punjab Agricultural University, Ludhiana, Punjab, 141004, India.

³Department of Plant Breeding and Genetics, Punjab Agricultural University, Ludhiana, Punjab, 141004, India.

***Author for correspondence:**

navjyot_grewal@yahoo.com

Received: 01/04/2025

Revised: 22/03/2026

Accepted: 17/04/2026

How to cite this article: Bhagat MA, N Kaur, P Goyal and S Kaur (2026) Evaluation of Barley (*Hordeum vulgare* L.) Germplasm for Heat Tolerance Using Germination Indices and Seedling Traits. *Indian J. Plant Genet. Resour.* 39(2): 58-66.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.178163>

Introduction

The average temperature of the Earth has risen between 0.5 and 0.8°C over the past 100 years due to global warming (Ju *et al.*, 2013). Scientists of the Intergovernmental Panel on Climate Change (IPCC) have predicted that the global annual temperature increased at an average rate of 0.08°C per decade since 1880 and over twice that rate (0.18°C/0.32°F) since 1981 (IPCC, 2022). There have been considerable changes in weather events since the onset of the middle of the 20th century, which have put global food security in great threat (Lesk *et al.*, 2016). Greenhouse gases are major causes of global warming. Over the last 250 years, 150% and 300% rise in the concentration of CH₄ and CO₂, respectively, have been documented (Friedlingstein *et al.*, 2010). It has been predicted that for every 1°C rise in temperature, there is 6% decline in global wheat production (Asseng *et al.*, 2015). In barley, grain yield and grain protein reduction of up to 29% and 22%, have been reported under high temperature stress (Ingvordsen *et al.*, 2015). This emphasizes the need to frame breeding programs that can target the development of new genotypes by incorporating the traits for heat stress tolerance (Rollins *et al.*, 2013).

Barley (*Hordeum vulgare* L.) is the fourth major cereal grain crop in the world after maize, rice and wheat, belonging to the family Poaceae (To *et al.*, 2020). It is commonly used for animal feeding, human food and malting purposes due to the

presence of important nutrients such as proteins, niacin, vitamins, fibres, and minerals (Huang *et al.*, 2020). As compared to other cereals such as maize, rice and wheat, barley shows promising adaptive capabilities under heat, drought and salinity stresses (Munns *et al.*, 2006). Therefore, barley is considered an important staple food crop in those regions of the earth that are sensitive to climate change. Due to such features, barley has become a model crop for the researchers working on stress-responsive genes (Li *et al.*, 2013).

Heat stress is one of the abiotic stresses that limits the production and productivity of barley (Sallam *et al.*, 2018; Bhagat *et al.*, 2025). The metabolic processes of plants are affected by high temperatures, which ultimately determine the crop's productivity. Information regarding the genetic variations, changes in physiological processes and level of existing genetic diversity in barley germplasm is needed to produce new cultivars having high tolerance to heat stress. Heat stress significantly reduces barley growth and yield by decreasing kernel weight, accelerating grain filling activities, and inducing early senescence (Sehgal *et al.*, 2018). Genetic improvement is considered the best tool for developing high-yielding barley cultivars with increased heat tolerance. Hawker and Jenner (1993) have reported that high temperature caused the inhibition of seed germination and seedling growth in barley by causing hindrance in the availability of energy and nutrients by preventing the synthesis and activity of hydrolytic enzymes like α -amylase. Lachno and Baker (1986) have reported that an increase in temperature results in ABA accumulation in seeds, which inhibits the activity of α -amylase.

Understanding the response mechanism to heat stress at seed germination stage may partially help to address some of the concerns for improving tolerance to heat stress and minimize consequent impacts. The genotypes exhibiting heat tolerance at the seedling stage have been reported to possess heat tolerance at the flowering and grain-filling stages as well. Wu (2013) reported that screening for heat stress tolerance at the seedling stage under controlled conditions in the growth chamber can be used for selecting heat-tolerant genotypes in cotton (*Gossypium hirsutum* L.). Screening germplasm for heat stress tolerance at seedling stage under *in-vitro* conditions is a simple, rapid, economical and time-effective methodology, as screening large germplasm under field conditions puts constraints in terms of time, labour and field resources. Keeping these in view, the present study was conducted to screen the advanced breeding lines of barley at seedling stage for selecting the heat-tolerant lines that can be used as

heat-tolerant donors in future barley breeding programs for the development of heat-tolerant varieties.

Materials and Methods

Experimental area and plant material

A set of ninety-two genotypes of barley was screened for heat tolerance at the seedling stage. For comparative studies, eight checks viz., DWRB-123, DWRUB 52, PL 807, PL 891, DWRB-92, PL 426, DWRB-137 and DWRB-101 were also evaluated alongwith the complete set. These genotypes were procured from the Department of Plant Breeding and Genetics (PBG), Punjab Agricultural University (PAU), Ludhiana. The genotypes from 1 to 57 were advanced breeding lines developed through pedigree method by the Department of PBG (Table 1). The genotypes from 58 to 92 were high-yielding and introduced from Institute National de la Recherche Agronomique du Maroc, Morocco (ICARDA-Morocco). The study was carried out at the Seed Physiology Laboratory, Office of the Director (Seeds), PAU, Ludhiana, India.

Table 1: Pedigree of one hundred genotypes of barley

Sr. no.	Entry name	Pedigree	Nomenclature in ICARDA, Morocco Nursery
1	BL1510	IEBON-33 (2010-11)/ PL426	-
2	BL1515	BK9816/DWRUB52	-
3	BL1521	BCU323/PL807	-
4	BL1523	PL830/DWR91	-
5	BL1531	PL807/PL861	-
6	BL1534	PL172/BCU347	-
7	BL1537	DWR83/RD2798	-
8	BL1539	PL840/PL833	-
9	BL1540	PL840/PL833	-
10	BL1544	BL71/13TH-HBSN-4	-
11	BL1566	VJM315/31TH-IBON-181	-
12	BL1594	BISON 87/CANELA	IBON-(2014-15)-E57
13	BL1607	LIGNEE527/GERBEL/3/ BOY-B*2/SURB//CI12225.2D/ 4/M104/6/LEGACY/4/TOCTE// GOB/HUMAI10/3/ATAH92/ ALELI/5/ESMERALDA	IBON-(2014-15)- E131
14	BL1608	ESMERALDA/3/LEGACY// PENCO/CHEVRON-BAR/4/ LA MOLINA 96	IBON-(2014-15)- E133
15	BL1610	E.ACACIA/DEFRA/3/ SVANHALS-BAR/MSEL// AZAF/GOB24DH	IBON-(2014-15)- E147

Sr. no.	Entry name	Pedigree	Nomenclature in ICARDA, Morocco Nursery	Sr. no.	Entry name	Pedigree	Nomenclature in ICARDA, Morocco Nursery
16	BL1649	GLORIA-BAR/COPAL// PM5/BEN/3/SEN/4/ PETUNIA 1/5/PETUNIA 2// PENCO/CHEVRON-BAR/ 4/PETUNIA 2/3/CHAMICO/ TOCTE//CONGONA	IBON-(2015-16)- E132	55	BL1808	PL807/PL841	-
17	BL1712	DWRB116/PL172	-	56	BL1810	PL807/PL61	-
18	BL1714	PL807/ BCU 361	-	57	BL1812	PL838/BH918	-
19	BL1715	PL807/ BCU 361	-	58	DWRB123	CHAMICO/TOCTE// CONGONA/6/P.STO/3/ LBIRAN/UNA80//LIGNEE640/ 4/BLLU/5/PETUNIA 1	-
20	BL1717	PL807/ BCU 361	-	59	IBON-4 (2018-19)	DWRUB54/DWR51	IBON-4 (2018-19)
21	BL1719	PL807/ BCU 361	-	60	IBON-5 (2018-19)	CANELA//E.QUEBRACHO/ W9338	-
22	BL1723	PL807/ BCU 361	-	61	IBON-8 (2018-19)	CANELA//LIMON/ BICHY2000	-
23	BL1724	PL807/ BCU 361	-	62	IBON-14 (2018-19)	PFC92126/ BICHY2000	-
24	BL1725	PL807/ BCU 361	-	63	IBON-23 (2018-19)	GLORIA-BAR/COPAL/ 6/P.STO/3/LBIRAN/ UNA80//LIGNEE640/4/ BLLU/5/PETUNIA 1	-
25	BL1728	PL807//PL426/BL10	-	64	IBON-39 (2018-19)	BLLU/6/P.STO/3/LBIRAN/ UNA80//LIGNEE640/4/ BLLU/5/PETUNIA 1	-
26	BL1729	PL830//PL426/BL10	-	65	IBON-55 (2018-19)	LBIRAN/UNA80//LIGNEE 640/3/PUNGSANCH APSSALBORI	-
27	BL1730	DWR96//PL426/BL10	-	66	IBON-58 (2018-19)	LBIRAN/UNA80//LIGNEE 640/3/PUNGSANCH APSSALBORI	-
28	BL1733	DWRUB 73//PL426/BL10	-	67	IBON-70 (2018-19)	J09072 F3 10/030575	-
29	BL1735	DWRUB 73//PL426/BL10	-	68	IBON-75 (2018-19)	P.STO/3/LBIRAN/ UNA80//LIGNEE640/4/ BLLU/5/PETUNIA1/6/ P.STO/3/LBIRAN/UNA80// LIGNEE640/4/BLLU/5/ PETUNIA 1	-
30	BL1736	DWRUB 73//PL426/BL10	-	69	IBON-78 (2018-19)	GLORIA-BAR/IAR.H.485// ALOE/3/CABUYA	-
31	BL1740	INBON-15 (2010-11)/PL426	-	70	IBON-88 (2018-19)	CANELA//ATAH92/ GOB	-
32	BL1743	INBON-15 (2010-11)/PL426	-	71	IBON-89 (2018-19)	CANELA//LIMON/BICHY 2000	-
33	BL1745	INBON-15 (2010-11)/PL426	-	72	IBON-90 (2018-19)	BISON 136/CANELA	-
34	BL1748	INBON-15 (2010-11)/PL426	-	73	IBON-91 (2018-19)	OPS66/CANELA	-
35	BL1750	INBON15 (2010-11)/DWRUB 73	-	74	IBON-107 (2018-19)	CIRU/3/AGAVE/ SUMBARD400//MARCO	-
36	BL1753	IEBON-33 (2010-11)/PL426	-	75	IBON-108 (2018-19)	MAHIGAN/H00058	-
37	BL1754	IEBON-33 (2010-11)/PL426	-				
38	BL1756	IEBON-33 (2010-11)/PL426	-				
39	BL1761	IEBON-33 (2010-11)/PL426	-				
40	BL1766	IEBON-33 (2010-11)/PL426	-				
41	BL1770	Moroc-9-73/A.Aswad//DWR96	-				
42	BL1771	Moroc-9-73/A.Aswad//DWR96	-				
43	BL1776	Moroc-9-73/A.Aswad //DWR96	-				
44	BL1780	BK9816/DWRUB52	-				
45	BL1781	BK9816/DWRUB52	-				
46	BL1784	EC492253/PL807	-				
47	BL1786	EC492253/PL807	-				
48	BL1787	EC492253/PL807	-				
49	BL1792	EC492253/PL838	-				
50	BL1794	BCU323/PL807	-				
51	BL1797	BCU323/PL807	-				
52	BL1801	BCU323/PL807	-				
53	BL1804	PL830/DWR94	-				
54	BL1807	PL807/PL841	-				

Sr. no.	Entry name	Pedigree	Nomenclature in ICARDA, Morocco Nursery
76	IBYT-E1 (2018-19)	P.STO/3/LBIRAN/UNA80 //LIGNEE640/4/BLLU/5/PETUNIA1/6/P.STO/3/LBIRAN/UNA80//LIGNEE 640/4/BLLU/5/PETUNIA 1	-
77	IBYT-E2 (2018-19)	P.STO/3/LBIRAN/UNA80// LIGNEE640/4/BLLU/5/PETUNIA1/6/P.STO/3/LBIRAN/UNA80//LIGNEE 640/4/BLLU/5/PETUNIA 1	-
78	IBYT-E5 (2018-19)	CIRU/ZIGZIG	-
79	IBYT-E6 (2018-19)	CIRU/ZIGZIG	-
80	IBYT-E7 (2018-19)	CANELA//LIMON/ BICHY2000	-
81	IBYT-E8 (2018-19)	CIRU/TOCTE	-
82	IBYT-E11 (2018-19)	S95031002008N/ H87010008N	-
83	IBYT-E12 (2018-19)	J09049 F3 10/030552	-
84	IBYT-E13 (2018-19)	MSEL/LOGAN-BAR	-
85	IBYT-E15 (2018-19)	MSEL/LOGAN-BAR	-
86	IBYT-E16 (2018-19)	LEGACY/CHAMICO// ATAH92/GOB	-
87	IBYT-E17 (2018-19)	AJO 61/6/Vmorales	-
88	IBYT-E19 (2018-19)	VMorales/6/ORGE618	-
89	IBYT-E20 (2018-19)	ALELI/SCARLETT	-
90	IBYT-21 (2018-19)	GLORIABAR/COPAL// PM5/BEN/3/SEN/4/PETUNIA1/5/PETUNIA2//PENCO/CHEVRONBAR/4/PETUNIA2/3/CHAMICO/TOCTE//CONGONA	-
91	IBYT-E22 (2018-19)	BBSC/CONGONA// FRESA	-
92	RIHANE-03 (IBYTE-24)	As46//Avt/Aths	IBYT-E24 (2018-19)
93	DWRB-123	DWRUB54/DWR51	
94	DWRUB52	DWR17/K551	-
95	PL807	LENT/BLLU//PINON	-
96	PL891 (IBYTE-24)	IBON 343 /12TH HSBN-176	-

Sr. no.	Entry name	Pedigree	Nomenclature in ICARDA, Morocco Nursery
97	DWRB-92	DWR28/DWR45	-
98	PL426	KARAN92/PL101	-
99	DWRB-137	DWR28/DWRUB64	-
100	DWRB-101	DWR28/BH581	-

*IBON, International Barley Observation Nursery; IBYT, International Barley Yield Trial; BL, Barley Ludhiana; DWRB, Directorate Of Wheat Research Barley; DWRUB, Directorate Of Wheat Research United Breweries.

Experimental design

The experiment was conducted in a completely randomized block design (CRBD) in four replications. One hundred seeds of each genotype in each replication of each treatment were sown using paper method as described by ISTA (2019). The seeds were surface sterilized using 0.5% sodium hypochlorite solution for 1-2 minutes, followed by washings thrice with autoclaved distilled water. These seeds were air-dried and placed on moistened germination paper. The paper was folded vertically. The seeds were allowed to germinate in a BOD incubator (REMI CI-10S). For heat stress, 30°C and 35°C temperature was maintained in BOD incubator. An optimum temperature 25°C was used as control. Twenty representative seedlings were used to measure seedling traits at 7-days after sowing (DAS).

Germination percentage (GP): The seeds with visible radicle protrusion were considered as germinated seeds. Per cent germination was calculated using the following formula (ISTA, 2019):

$GP = (\text{Total no. of seeds germinated} / \text{Total no. of seeds placed}) \times 100$

Root and shoot length (RL and SL): Root and shoot length of twenty seedlings of each genotype of each replication and each treatment were measured using centimeter scale at 7 DAS. The mean RL and SL were recorded in centimetres. Seedling length was computed by adding SL and RL, which was used for the calculation of the seedling vigour index I.

Seedling dry weight (DW): Twenty seedlings of each genotype from each replication and each treatment were dried in an oven at $65 \pm 2^\circ\text{C}$ until a constant weight was obtained. The mean seedling dry weight was recorded in grams. Seedling dry weight was further used for the calculation of seedling vigour index II.

Vigour index (VI): Seedling vigour index I (SVI I) and seedling vigour index II (SVI II) were calculated using the following formula (Abdul-Baki and Anderson, 1973):

SVI I = Germination(%) × mean seedling length (cm)

SVI II = Germination(%) × mean seedling dry weight (g)

Speed of germination: Germination counts were made on 4th, 5th, 6th and 7th days after the start of the experiment, with the criterion for germination being visible protrusion of the radicle from seed. Speed of germination (SG) was calculated using the following formula as suggested by ISTA (2019):

$$\frac{n_1}{d_1} + \frac{n_2}{d_2} + \frac{n_3}{d_3} + \dots$$

where, n = number of seeds germinated, d = number of days.

Statistical analysis

The variations in germination and seedling traits of one hundred genotypes were represented using a colour gradient heat map in Prism Graphpad (v.9.0, Prism Inc, San Diego, CA, USA). The data were subjected to two-way analysis of variance (ANOVA) to find differences among genotypes and treatments at 1% and 5% level of significance using Tukey's test (Tukey 1949).

The variation of responses according to the temperature differences was evaluated using the Ward's algorithm and the Euclidean distance under the Hierarchical Cluster Analysis (HCA) (Cha-um *et al.*, 2010). Cluster analysis of the data was done using SPSS version 17.0 software (SPSS Inc., Chicago, IL, USA). The cluster ranking for the groups was obtained based on Ward's minimum variance cluster analysis using the differences in the values of morpho-physiological parameters viz., germination percentage (%), speed of germination, root and shoot length, seedling dry weight and seedling vigour index I and II (Zeng *et al.*, 2002; Ravikiran *et al.*, 2018).

Results and Discussion

Effects of heat stress on germination indices and seedling traits of barley genotypes

The morpho-physiological characteristics viz., germination percentage, speed of germination, root length (RL), shoot length (SL), seedling dry weight, seedling length and seedling vigour index I and II, significantly varied ($p < 0.01$) among genotypes and treatments. ANOVA revealed that temperature, genotypes and their interaction significantly contributed to germination percentage, speed of germination, root length, shoot length, seedling length, seedling dry weight and vigour (Table 2).

Germination percentage

There were significant differences in germination percentage among the genotypes and treatments at 1% level of significance. The per cent germination ranged from 45-97.7% with a mean value 82.8% at 25°C (control), 2.5%-94.7% with a mean value 52.3% at 30°C and 1.3-86.3% with mean value 37.6% at 35°C (Table 3). Mean germination percentage decreased by 36% and 55% at 30°C and 35°C, respectively, as compared to mean germination percentage at 25°C. Heat-tolerant barley genotypes exhibited minor reduction in germination percentage (12.7%) as compared to heat-sensitive genotypes (87.6%) at 35°C (Fig.1a). Temperature showed considerable effects on germination characteristics, including the initiation and the final germination percentage. It is therefore the most critical factor in the determination of the success or failure of plant establishment (Al-Ahmadi and Kafi, 2007). Reduction in germination due to heat stress in wheat has been attributed to impairments in the activities of α -amylase, which hampers starch breakdown and abolishes the delivery of nutrients to developing embryos (Essemine *et al.*, 2010). Reduction in germination is also attributed to a reduction in

Table 2: F-values of ANOVA for germination and seedling traits in one hundred genotypes of barley

Source of variance	GP	SOP	RL	SL	TSL	DW	SVI I	SVI II
Temperature	263.3 ^{**}	271.8 ^{**}	141.3 ^{**}	103 ^{**}	167.9 [*]	410.3 ^{**}	299.6 ^{**}	109.2 [*]
Genotype	669.4 ^{**}	54.88 ^{**}	20.36 ^{**}	8.9 ^{**}	13.05 ^{**}	16.03 ^{**}	141.94 ^{**}	76.42 [*]
T×G	85.35 ^{**}	10.16 ^{**}	5.79 ^{**}	5.25 ^{**}	5.61 ^{**}	5.27 ^{**}	36.93 ^{**}	12.38 ^{**}

^{**}significant at 0.01 level: GP, germination percentage; SOP, Speed of germination; RL, Root length; SL, Shoot length; TSL, Total seedling length; DW, Dry weight; SVI I, Seedling vigour index I; SVI II, Seedling vigour index II.

metabolic efficiency at high temperature than optimum temperature in *Eurotia lanata* (Thygersen et al., 2002). Heat stress caused inhibition of seed germination through the synthesis of ABA in wheat (Hasanuzzaman et al., 2013). Heat and salt stress induced decrease in germination has also been documented in other plant species, viz., *Triticum aestivum* (Hakim et al., 2012; Hossain and Teixeira da Silva, 2012; Singh et al., 2025a, 2025b), *Zea mays* (Chhatwal et al., 2025), *Moringa peregrine* (Sardooi et al., 2019) and *Brassica juncea* (Toosi et al., 2014).

Speed of germination

The increment in speed of germination of barley genotypes was observed at 30°C; but a reduction was recorded at 35°C as compared to control temperature (25°C). It ranged from 0.8-17.8 at 25°C, 1.9-18.2 at 30°C and 0.3-8.7 at 35°C (Fig. 1b). The interaction effect of temperature and genotypes was significant for speed of germination in barley genotypes. The F-value observed

for temperature on speed of germination was 271.8 and 54.88 for genotypes indicating higher contribution of temperature towards reducing the speed of germination (Table 2). These results showed agreement with findings of Toosi et al., (2014) in *Brassica juncea* and Lima et al., (2021) in *Dalbergia spruceana*, who also observed that increase in temperature up to 30°C caused an increase in speed of germination and declined with the further increase in temperature to 35°C. With an increase in temperature, the activities of genes for production of GA are upregulated and thus speed of germination is hastened in *Brassica* (Finch-Savage and Footitt, 2012).

Root and shoot length

Seedlings RL and SL varied from 10.35-21.55 cm and 11.7-19.85 cm, respectively at 25°C, 8.0-19.6 cm and 6.3-22.13 cm, respectively at 30°C and 0.5-5.85 cm and 2.5-6.8 cm, respectively at 35°C (Fig. 2a & b). RL and SL of seedlings decreased significantly at 35°C but meagre




Fig. 1: Heat maps depicting effect of three temperature regimes (25, 30 and 35°C) on (a) germination percentage and (b) speed of germination in one hundred genotypes of barley. Higher colour intensity corresponds to higher germination percentage and speed of germination in figures (a) and (b) respectively and vice-versa.




Fig. 2: Heat maps depicting effect of three temperature regimes (25, 30 and 35°C) on (a) shoot length (cm) and (b) root length (cm) in one hundred genotypes of barley. Higher colour intensity corresponds to higher shoot length and root length in figures (a) and (b) respectively and vice-versa.

reduction in RL and SL of seedlings was observed at 30°C. The RL ($F=141.3$) of seedlings showed more reduction as compared to SL ($F=103.0$) with the effects of heat stress being significant for both parameters ($p<0.01$). Reduction in shoot and root length in seedlings of wheat germplasm under heat stress environment was also reported by Lu *et al.*, (2022) and Hasan *et al.*, (2024).

Seedling vigour index I and II

Seedling vigour index I and II varied from 1419.8-3550.5 with mean value of 2635.0 and 10.3-23.4 with an average value of 16.4 respectively, under control conditions (25°C) (Table 3). Mean SVI I decreased by 89.2% at 35°C as compared to 25°C. The highest mean SVI II (16.3) was reported at 25°C trailed by at 30°C (7.8) and 35°C (3.7) (Fig. 3a, Table 4). The interaction effect between temperature and genotypes was also reported to be significant for both SVI I and II.

The results indicated that the SVI I ($F=299.6$) witnessed a higher reduction as compared to SVI II ($F=109.2$), with the effects of heat stress being significant for both parameters ($p<0.01$) (Table 2). Among selected barley genotypes, SVI II of sensitive genotypes ranged from 0.4 to 1.2 and tolerant genotypes from 6.1 to 11.2 at 35°C (Fig. 3b, Table 4). SVI II

Table 3: Mean values of morpho-physiological parameters at three temperature regimes of one hundred genotypes of barley

Trait	Temperature		
	25°C	30°C	35°C
Germination percentage	82.8	52.3	37.6
Speed of germination	6.6	10.0	72.5
Shoot length (cm)	15.5	12.8	4.0
Root length (cm)	16.3	13.4	2.9
Seedling length (cm)	31.8	26.3	7.0
Seedling vigour index I	2635.0	1436.5	285.7
Seedling vigour index II	16.3	7.8	3.7

Table 4: Mean values of germination and seedling parameters at three temperature regimes in heat-tolerant and heat sensitive genotypes of barley

Trait	Temperature					
	25°C		30°C		35°C	
	Heat-tolerant genotypes	Heat-sensitive genotypes	Heat-tolerant genotypes	Heat-sensitive genotypes	Heat-tolerant genotypes	Heat-sensitive genotypes
Germination percentage	95.0	74.9	91.8	14.3	82.9	9.3
Speed of germination	11.4	3.5	14.6	1.7	5.8	0.4
Seedling vigour index I	2922.3	2271.4	2558.8	297.3	623.0	50.7
Seedling vigour index II	18.2	15.1	13.6	1.7	8.5	0.8

of tolerant barley genotypes decreased by 37.5% at 35°C as compared to 30°C; while SVI II of sensitive genotypes decreased by 52.9% at 35°C as compared to 30°C. SVI I determines the fate of seedling establishment and incorporates the combined effects of germination and seedling growth. The higher vigour index of heat-tolerant genotypes could be attributed to higher germination rate and seedling length. On the




Fig. 3: Heat maps depicting effect of three temperature regimes (25, 30 and 35°C) on (a) seedling vigor index I and (b) seedling vigor index II in one hundred genotypes of barley. Higher colour intensity corresponds to higher seedling vigor index I and II in figures (a) and (b) respectively and vice-versa.




Fig. 4: Clustering of one hundred genotypes of barley using cluster dendrogram based on the furthest neighbour to determine the heat tolerance potential.

other hand, the sensitive ones had a lower germination rate and seedling length, and ultimately had a lower vigour index. The decline in germination and in turn, seedling vigour has also been observed due to high temperature stress during early seedling growth in wheat (Hasan *et al.*, 2013). Reduction in the seedling dry weight and vigour was a result of reduction in seed reserve mobilization due to heat stress in wheat (Essemine *et al.*, 2010).

Hierarchical clustering

Cluster ranking based on morphological and physiological changes in barley seedlings under heat stress classified the one hundred barley genotypes into three groups: highly heat tolerant (21), moderately heat tolerant (27) and heat sensitive (52) (Fig. 4). Out of these categorized genotypes, nine heat tolerant genotypes viz., BL1515, BL1729, BL1780, BL1784, BL1786, BL1792, BL1794, L1797 and IBYT-E24 can be used as potential donors for developing heat-tolerant barley varieties in future breeding programs.

Conclusion

Temperature is a detrimental factor for plant growth and productivity worldwide. In the present study, one hundred barley genotypes were evaluated for their heat tolerance at seedling stage on the basis of morpho-physiological parameters, i.e. speed of germination, root length, shoot length, seedling dry weight, germination percentage, seedling vigour index I and II. Further, HCA (hierarchical clustering analysis) segregated the barley genotypes into three categories, i.e. highly heat tolerant (21), moderately heat tolerant (27) and heat sensitive (52) based on their tolerance to heat stress. Seedling vigour index was recognized as

promising trait for selection of heat-tolerant genotypes. Selected heat-tolerant barley genotypes in the present study could be used for future breeding programmes for the introgression of heat tolerance in cultivated barley varieties.

References

- Abdul-Baki AA and JD Anderson (1973) Vigour determination in soybean seed by multiple criteria. *Crop Sci.* 13(6): 630-633.
- Al-Ahmadi MJ and M Kafi (2007) Cardinal temperatures for germination of *Kochia scoparia* (L.). *J. Arid Environ.* 68(2): 308-314.
- Asseng S, F Ewert and P Martre (2015) Rising temperatures reduce global wheat production. *Nat. Clim. Change* 5(2): 143-147.
- Bhagat MA, N Kaur, L Vij, P Goyal and S Kaur (2025) Comparison of physio-biochemical and yield traits in barley (*Hordeum vulgare* L.) germplasm under terminal heat stress conditions in Northern India. *Russian J. Plant Physiol.* 72: 197.
- Cha-um S, M Ashraf and C Kirdmanee (2010) Screening upland rice (*Oryza sativa* L. ssp. *indica*) genotypes for salt-tolerance using multivariate cluster analysis. *Afr. J. Biotech.* 9(30): 4731-4740.
- Chhatwal H, N Gupta, M Goyal, P Goyal, GK Gill and N Garg (2025) Modulation of antioxidant system by glutathione in maize seedlings under salt stress conditions. *Indian J. Ecol.* 52(4): 750-754.
- Essemine J, S Ammar and S Bouzid (2010) Impact of heat stress on germination and growth in higher plants; physiological, biochemical and molecular repercussions and mechanisms of defence. *J. Biol. Sci.* 10(6): 565-572.
- Finch-Savage WE and S Footitt (2012) To germinate or not to germinate; a question of dormancy relief not germination stimulation. *Seed Sci. Res.* 22(4): 243-248.
- Friedlingstein P, RA Houghton and G Marland (2010) Update on CO₂ emissions. *Nat. Geosci.* 3(12): 811-812.
- Hakim MA, A Hossain, JA Teixeira da Silva, VP Zvolinsky and MM Khan (2012) Yield, protein and starch content of 20 wheat (*Triticum aestivum* L.) genotypes exposed to high

- temperature under late sowing conditions. *J. Sci. Res.* 4(2): 477-89.
- Hasan MA, JU Ahmed, T Hossain, MAK Mian and MM Haque (2013) Evaluation of the physiological quality of wheat seed as influenced by high parent plant growth temperature. *J. Crop Sci. Biotech.* 16: 69-74.
- Hasan MM, MAB Mia, JU Ahmed, MA Karim, AKM Aminul Islam and MMohi-Ud-Din (2024) Heat stress tolerance in wheat seedling; clustering genotypes and identifying key traits using multivariate analysis. *Heliyon* 10(19): e38623.
- Hasanuzzaman M, K Nahar, MM Alam, R Roy chowdhury and M Fujita (2013) Physiological, biochemical, and molecular mechanisms of heat stress tolerance in plants. *Int. J. Mol. Sci.* 14(5): 9643-9684.
- Hawker JS and CF Jenner (1993) High temperature affects the activity of enzymes in the committed pathway of starch synthesis in developing wheat endosperm. *Aust. J. Plant Physiol.* 20: 197-209.
- Hossain A and JA Teixeira da Silva (2012) Phenology, growth and yield of three wheat (*Triticum aestivum* L.) varieties as affected by high temperature stress. *Not. Sci. Biol.* 14(3): 97-109.
- Huang H, X Gao, Y Li, P Tian, Y Nima, Z Laba and W Liao (2020) Content analysis of vitamins, dietary fibers and amino acids in a wide collection of barley (*Hordeum vulgare* L.) from Tibet, China. *Bioinformation* 16(4): 314.
- Ingvordsen CH, G Backes, MF Lyngkjær, P Peltonen-Sainio, JD Jensen, M Jalli, A Jahoor, M Rasmussen, TN Mikkelsen, A Stockmarr and RB Jørgensen (2015) GWAS of barley phenotypes established under future climate conditions of elevated temperature, CO₂, O₃ and elevated temperature and CO₂ combined. *Procedia Environ. Sci.* 29: 164-165.
- IPCC (2022) Climate Change 2022: Impacts, Adaptation, and Vulnerability. In: Pörtner HO, DC Roberts, M Tignor, ES Poloczanska, K Mintenbeck, A Alegría, M Craig, S Langsdorf, S Löschke, V Möller, A Okem and B Rama (eds.). Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK and New York, USA.
- ISTA (2019). International rules for seed testing. International Seed Testing Association, Sapporo, Japan.
- Ju H, M van der Velde, E Lin, W Xiong and Y Li (2013) The impacts of climate change on agricultural production systems in China. *Clim. Change* 120: 313-324.
- Lachno DR and DA Baker (1986) Stress induction of abscisic acid in maize roots. *Physiol. Plant* 68: 215-221.
- Lesk C, P Rowhani and N Ramankutty (2016) Influence of extreme weather disasters on global crop production. *Nature* 529: 84-87.
- Li WT, C Liu, YX Liu, ZE Pu, SF Dai, JR Wang and YM Wei (2013) Meta-analysis of QTL associated with tolerance to abiotic stresses in barley. *Euphytica* 189: 31-49.
- Lima CC, ESC Gurgel and EEDL Borges (2021) Antioxidant enzyme activity in germination of *Dalbergia spruceana* seeds under different temperatures. *J. Seed Sci.* 43: e202143006.
- Lu L, L Hui, W Yu and Y Guijun (2022) Wheat genotypes tolerant to heat at seedling stage tend to be also tolerant at adult stage; the possibility of early selection for heat tolerance breeding. *The Crop Journal* 10(4): 1006-1013.
- Munns R, RA James and A Läuchli (2006) Approaches to increasing the salt tolerance of wheat and other cereals. *J. Exp. Bot.* 57(5): 1025-1043.
- Ravikiran KT, SL Krishnamurthy, AS Warraich and PC Sharma (2018) Diversity and haplotypes of rice genotypes for seedling stage salinity tolerance analyzed through morpho-physiological and SSR markers. *Field Crops Res.* 220(8): 10-18.
- Rollins JA, B Drosse, MA Mulki, S Grando, M Baum, M Singh and M VonKorff (2013) Variation at the vernalisation genes *Vrn-H1* and *Vrn-H2* determines growth and yield stability in barley (*Hordeum vulgare*) grown under dry land conditions in Syria. *Theor. Appl. Genet.* 126: 2803-2824.
- Sallam A, A Amro, A El-Akhdar, MF Dawood, T Kumamaru and PS Baenziger (2018) Genetic diversity and genetic variation in morpho-physiological traits to improve heat tolerance in spring barley. *Mol. Biol. Rep.* 45: 2441-2453.
- Sardooi ER, A Hashemi, HE Damaneh, H Khosravi and S Barkhori (2019) Assessment of seed germination of *Moringa peregrina* under drought and salinity stress and its cardinal germination temperatures in laboratory environment. *Desert Ecosyst. Eng. J.* 2(1): 21-30.
- Sehgal A, K Sita, KH Siddique, R Kumar, S Bhogireddy, RK Varshney and H Nayyar (2018) Drought or/and heat-stress effects on seed filling in food crops; impacts on functional biochemistry, seed yields, and nutritional quality. *Front. Plant Sci.* 9: 1705.
- Singh P, P Goyal, S Singh and A Sharma (2025a) Salinity induced stomatal and physio-biochemical responses in 'Berbet' introgression lines of wheat (*Triticum aestivum* L.). *Russian J. Plant Physiol.* 72(4): 133.
- Singh P, P Goyal and A Sharma (2025b) Multi-dimensional evaluation of responses to salt stress of *Berbet* introgression lines of wheat (*Triticum aestivum* L.). *Plant Genet. Resour. Characterisation Util.* 23(5): 351-362.
- Thygeson T, JM Harris, BN Smith, LD Hansen, RL Pendleton and DT Booth (2002) Metabolic response to temperature for six populations of winter fat (*Eurotia lanata*). *Thermochimica Acta* 394: 211-217.
- To VT, Q Shi, Y Zhang, J Shi, C Shen, D Zhang and W Cai (2020) Genome-wide analysis of the GRAS gene family in barley (*Hordeum vulgare* L.). *Genes* 11(5): 553.
- Toosi AF, BB Bakar and M Azizi (2014) Effect of drought stress by using PEG 6000 on germination and early seedling growth of *Brassica juncea* var. *Ensabi* Scientific Papers Series A Agronomy 57: 360-363.
- Tukey JW (1949) Comparing individual means in the analysis of variance. *Biometrics* 5(2): 99-114.
- Wu T (2013) Identification of heat tolerant upland cotton (*Gossypium hirsutum* L.) germplasm utilizing chlorophyll fluorescence measurement. Doctoral Dissertation, Auburn University, Auburn, Alabama, USA.
- Zeng L, MC Shannon and CM Grieve (2002) Evaluation of salt tolerance in rice genotypes by multiple agronomic parameters. *Euphytica* 127(2): 235-245.

RESEARCH ARTICLE

Effect of Various Seed Priming Agents on Germination, Seedling Vigour and Grain Yield of Dry Direct-Seeded Rice in the Northwestern Indo Gangetic Plains of India

Sukham Madaan^{1*}, Mangat Ram¹, Lalita¹, Sumit Saini¹, Ashish Jain¹, Rakesh Kumar¹, Mahaveer Bochalya¹, Amit Kumar¹, Ashwani Kumar¹, Vishal Gandhi¹, Charan Singh¹, Parvesh Kumar¹, Sarita Devi² and Anita Kumari²

Abstract

Dry direct-seeded rice (DSR) grown in sub-tropical northwestern Indo-Gangetic Plains of India is often exposed to multiple environmental stresses during germination phase, leading to poor crop establishment and productivity. Seed priming may help overcome these stresses by ensuring better crop germination and establishment. The present study was carried out during 2022 and 2023 to assess the effectiveness of priming agents viz., hydropriming, nanopriming (25 ppm ZnO nanoparticles), hormonal priming (50 ppm Gibberellic acid, GA₃), halopriming (1% KNO₃) and ascorbic acid (100 ppm) priming in two rice cultivars under laboratory as well as dry direct seeded field conditions. All priming agents significantly improved seed germination and seedling growth, seedling length and seedling vigour under laboratory conditions and stand establishment of the crop in the field. Hormonal priming was found to be significantly superior to all the other treatments under laboratory conditions, but all the priming agents were equally effective under field conditions. The priming increased number of panicles/m² and ultimately grain yield of field crop significantly over the unprimed control, obviously due to improvement in crop stand (number of plants/m²).

Keywords: DSR, Germination, Hormonal Priming, Nano Priming, Plant Stand

¹Rice Research Station, Kaul (Kaithal), Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, 136021, India.

²Department of Botany and Plant Physiology, Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, 125004, India.

***Author for correspondence:**
sukham20@gmail.com

Received: 03/01/2026

Revised: 02/04/2026

Accepted: 08/04/2026

How to cite this article: Madaan S, M Ram, Lalita, S Saini, A Jain, R Kumar, M Bochalya, A Kumar, A Kumar, V Gandhi, C Singh, P Kumar, S Devi and A Kumari (2026) Effect of Various Seed Priming Agents on Germination, Seedling Vigour and Grain Yield of Dry Direct-Seeded Rice in the Northwestern Indo Gangetic Plains of India. *Indian J. Plant Genet. Resour.* 39(2):67-73.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.174774>

Introduction

Rice (*Oryza sativa* L.) is the most important staple food of the world and not only feeds more than half the global population but also serves as a cultural and economic cornerstone in many regions (Asma *et al.*, 2023). In India, it is a significant rainy season crop in the sub-tropical region of the northwestern Indo-Gangetic Plains (IGP), particularly in the states of Punjab and Haryana, and plays a vital role in the country's food security. The region is famous for growing fine-grain aromatic (basmati) rice in addition to high-yielding coarse-grain rice. Rice in this region is generally grown by the transplanting method of crop establishment, in which rice seedlings first grown in a nursery and then seedlings are transplanted (usually manually) into the puddled main field. But the transplanted rice (TPR) requires a large amount of water not only for field preparation (puddling) but also for maintenance of submergence for most part of the crop duration. Brar *et al.*, (2015) reported that TPR requires a very high quantity of irrigation water (1357-1666 mm) to obtain a higher yield. The water requirement of the rice crop is largely met through the use of groundwater, as it is the principal source of irrigation in

the region. This has resulted in the over-exploitation of groundwater resources in the region, resulting in a decline of the water table to the extent that pumping out of groundwater by tube wells is increasingly becoming a difficult and costly affair due to the high cost of installation and maintenance of large pump sets (deep tube wells) as well as high electricity consumption to operate them.

Therefore, it is the need of the hour to adopt water-saving rice establishment techniques as sustainable alternative to conventional water-intensive transplanting methods, aimed at maintaining rice production in the region while conserving valuable groundwater resources. Direct seeding of rice is an establishment technique in which rice seeds are sown directly in the main field rather than transplanting the seedlings from a nursery and is grown in aerobic conditions (irrigated at frequent intervals) as compared to TPR, which requires continuous flooding during most part of the growing period. The direct seeding offers a good alternative to transplanting as it saves the extra water required for puddling and maintenance of submerged conditions, throughout growing season, besides eliminating the need for nursery raising and manual labour required in conventional transplanting. The direct-seeded rice (DSR) has been reported to save up to 35% of irrigation water over the conventional TPR while saving up to 46% labour and 32% production cost (Saha *et al.*, 2022). In recent years, direct-seeded (aerobic) rice has gained momentum in irrigated lowland areas of northwestern IGP, where rainfall is inadequate and pumping groundwater from deep wells is highly expensive (Sandhu *et al.*, 2019).

Direct seeding of rice can be done either by sowing in moist soil following pre-sowing irrigation (locally known as *vattar* DSR) or by sowing in dry soil followed by irrigation (dry DSR). However, dry DSR is more common (conventional) practice in the area as where DSR is growing in Haryana and Punjab (Dhillon *et al.*, 2021). But there are several constraints to the widespread adoption of the direct seeding technique of rice cultivation. In addition to heavy weed infestation, poor germination and poor or non-uniform crop establishment are the major constraints to the adoption of DSR in northwestern IGP (Sandhu *et al.*, 2021; Dhillon *et al.*, 2021). Poor crop establishment of DSR can be caused by several factors that determine the soil moisture conditions during crop emergence (Alam *et al.*, 2020). In conventional DSR, the field inundation caused by unexpected rains during the crop germination period (before seedling emergence) creates temporary excess water stress. This may

adversely affect the seed germination and emergence of seedlings, resulting in poor crop establishment and ultimately poor crop yield (Dhillon *et al.*, 2021). Moreover, in many cases, the topsoil dries up quickly after the rains or irrigation, resulting in the formation of a hard-thin layer (crust) which can hinder the emergence of the seedlings, leading to poor crop stand. Rapid and uniform germination as well as early seedling vigour, are crucial to improve the performance of rice under direct-seeded situations (Mahender *et al.*, 2015). Therefore, sustainable and effective technologies are needed to improve germination, seedling emergence and early growth of DSR to obtain its higher yield.

Seed priming is a controlled hydration technique that involves soaking the seeds in a solution of priming agents to induce pre-germination metabolic processes, without actual germination. This technique improves germination and seedling vigour under adverse environments. The seeds are re-dried to a moisture content below the threshold required for triggering radicle emergence to complete the priming, and the primed seed is then transferred to a growing medium (such as soil) where it can reabsorb water until the radicle protrudes and grows. Priming reactivates metabolic processes crucial for germination, including respiration, protein synthesis and enzyme activation (Ali and Elozeiri, 2017). Seed priming has emerged as a practical and effective approach to enhance seed performance under various environmental stresses in crop plants, including rice (Dhillon *et al.*, 2021; Paul *et al.*, 2022). In addition to soaking of seeds in water (hydropriming), seed priming in rice can also be done by treating seeds with specific salts (halopriming), growth regulators (hormonal priming), nanoparticles (nanopriming) or other chemicals (Khalid *et al.*, 2019; Dhillon *et al.*, 2021; Adhikary *et al.*, 2022; Zhang *et al.*, 2024), but the chemical and its concentration need to be identified for the dry DSR.

However, there is a dearth of data on the effects of various priming agents in rice, particularly in the dry DSR system of the northwestern IGP. Despite the proven benefits of seed priming, challenges such as reduced seed longevity post-priming and variability in crop or varietal-specific responses remain (Rhaman *et al.*, 2020), necessitating tuning of variables such as priming agents for specific conditions (Pagano *et al.*, 2023). Moreover, many earlier studies have been limited to laboratory environments, concentrating mostly on emergence, germination and early development, with a very few evaluating the crop productivity (Zheng *et al.*, 2016; Dhillon *et al.*, 2021). The present investigation was, therefore, conducted to assess the efficacy of

various priming agents on the germination and growth of rice seedlings in the laboratory as well as on stand establishment and performance of the crop in the field under dry direct seeded aerobic conditions.

Materials and Methods

Laboratory and field experiments were conducted during rainy (*khari*) seasons of 2022 and 2023 at CCS Haryana Agricultural University, Rice Research Station, Kaul (Latitude 20°51' N and Longitude 76°39' E), Hisar, Haryana to study the effects of different seed priming treatments on germination and seedling vigour of rice under laboratory conditions and yield of rice crop grown in the field under dry direct seeded (aerobic) conditions. The treatments consisted of two rice varieties viz., HKR 128 (V_1), a coarse grain high yielding variety of rice, and Pusa Basmati 1121 (V_2), a fine grain variety of basmati (aromatic) rice, and six priming treatments viz., no priming or control (T_1), hydropriming (T_2), nanopriming with ZnO nanoparticles @ 25 ppm (T_3), hormonal priming with gibberellin (GA_3) @ 50 ppm (T_4), halopriming with KNO_3 @ 1% (T_5) and priming with ascorbic acid @ 100 ppm (T_6), resulting in twelve treatment combinations. The treatments were laid out in a factorial randomized block design with three replications for each combination. For priming, dry seeds (12-13% moisture) of the two rice varieties (500 g each) were soaked in solutions of various priming agents for 16 hours in an equal volume of either solution (for T_3 , T_4 , T_5 , and T_6) or water (for T_1). After priming, the seeds were redried (in shade) to their original moisture content. The primed seeds were put in polythene bags and kept in a refrigerator at $5\pm 2^\circ\text{C}$ until they were used.

One set of the primed seeds (300 seeds for each treatment combination) was used for the laboratory experiment and was divided into three replications (100 seeds/replication). In the laboratory experiment, the seeds of each replication were placed on moist germination papers and kept in a seed germinator under dark conditions at a temperature of $25\pm 2^\circ\text{C}$ and relative humidity of $90\pm 1\%$. These seeds were evaluated for laboratory observations on germination (%) and seedling vigour parameters viz., seedling length, seedling dry weight and vigour indices viz., vigour index-I and vigour index-II (ISTA, 2019) after 14 days. The number of germinated seeds in each replication was counted to determine germination percentage. For seedling length, the shoot length of ten randomly selected seedlings per replication was measured from the base of the seedling to the tip of the

longest leaf and averaged. After measuring the seedling length, these ten seedlings were oven-dried for 24 h at $80\pm 1^\circ\text{C}$ and subsequently weighed to determine the average seedling dry weight. The vigour indices were determined using the formula suggested by Abdul-Baki and Anderson (1973), according to which the vigour index-I was determined by multiplying the seed germination (%) by the average length of the seedlings, whereas the vigour index-II was determined as the product of the germination (%) and average dry weight of the seedlings.

The second set of primed seeds was used for a field experiment on the crop to be grown under conventional direct-seeded (aerobic) conditions. In the field experiment, the primed seeds for each treatment combination were sown in the field (at 3-4 cm depth) on a well prepared flat dry seed bed in plots of size 4.2 m $\times$ 3.0 m (for each replication) in rows 20 cm apart with a plant to plant (intra-row) spacing of 15 cm, sowing one seed per hill (place) in each row. The field was irrigated immediately after sowing as a standard agronomic practice for a dry direct-seeded crop. The recommended agronomic and plant protection practices, such as land preparation, weed management, irrigation, fertilization, and disease or insect-pest management, were followed uniformly across all plots. Observations on seedling establishment or plant stand (number of plants/ m^2), physiological indices, yield attributes (number of plants/ m^2 , number of grains/panicles, and 1000-grain weight), and grain yield of the DSR crop were recorded from the net plot (3.6 m $\times$ 2.6 m) of each treatment. The number of plants/ m^2 was determined at 15 days after sowing (DAS) by placing a quadrat (0.5 m $\times$ 0.5 m) randomly at three places along the rows in each plot and counting the total number of plants within the quadrat area and then converting the number on a per square metre basis. Physiological indices, viz. chlorophyll content (in terms of SPAD values) and chlorophyll fluorescence (CF), were recorded at flowering initiation and grain filling stages of the crop. The SPAD values were recorded by using a portable handheld chlorophyll meter (SPAD-502 Plus, Konica Minolta Optics, Japan), whereas chlorophyll fluorescence was measured with the help of a portable hand-held chlorophyll fluorometer (OS-30p+ Opti-Sciences, Inc.). The yield attributes and grain yield (kg/ha) of the crop were recorded at maturity. The number of panicles/ m^2 was obtained by counting all effective (panicle-bearing) tillers within a quadrat (0.5 m $\times$ 0.5 m) placed randomly at three different places along the rows in each plot and then converting the count on per square-meter basis. The number of

grains/panicles was estimated by harvesting a representative set of panicles (10 panicles) from each plot, threshing them separately, and counting their grains. The 1000-grain weight was determined by randomly taking a sample of grains from the harvested/threshed produce (grain) of each net plot, counting 1000-grains from the sample and recording their weight (g). Grain yield was measured as the total weight of grains obtained from each plot (after drying to 22% moisture) converted to kg/ha.

Data on the laboratory and field observations were statistically analysed by the analysis of variance (ANOVA) technique with the help of an online statistical tool (OPSTAT) accessible at the website of CCS HAU, Hisar (www.hau.ac.in). The means of each treatment were compared with the help of the critical difference (CD) determined at 5% level of significance.

Results and Discussion

Seed germination and seedling vigour

Under laboratory conditions, rice variety Pusa Basmati 1121 exhibited significantly higher germination percentage (82.9-87.4%) and seedling length (27.9-35.0 cm) as compared to HKR 128 (75.4-81.0% and 24.3-30.2 cm) during the two years of laboratory investigation. Pusa basmati 1121 also showed significantly higher seedling vigour Index-I (2298-3088) and vigour index-II (8.37-8.57) than HKR 128 (1842-2426 and 7.77-7.78). Rice varieties may differ in terms of activities of amylase

and other enzymes, as well as the contents of sugars and soluble proteins (Monajjem *et al.*, 2023), which might be responsible for the differences in their germination potential.

The data on laboratory studies presented in Table 1 showed that priming of rice seeds with all the agents significantly increased the germination and seedling growth in terms of seedling length, seedling vigour-I, and seedling vigour-II over the unprimed control during both years. Among the priming treatments, the highest mean (averaged over the varieties) germination (84.0-90.3%), seedling length (29.2-37.2 cm), seedling vigour-I (2460-3359) and seedling vigour-II (8.79-9.03) was achieved when seeds were primed with GA₃ @ 50 ppm (T₄) and this treatment was found significantly superior as compared to other priming treatments, including hydro-priming (T₂). Priming with GA₃ @ 50 ppm improved the germination, seedling length, vigour index-I and vigour index-II by 15.1-16.1, 22.0-26.0, 41.6-45.3 and 20.4-22.2 per cent, respectively, over the unprimed control during the two years of study. Seed priming with gibberellin (GA₃) improved the germination and seedling emergence, possibly due to breakage of seed dormancy and an increase in the activity of starch-metabolizing enzymes (hydrolases) such as α -amylase, which convert the stored starch in the endosperm into soluble sugars to provide energy for germination or growth of the embryo (Waqas *et al.*, 2019; Wang *et al.*, 2021). The better germination and seedling growth or vigour obtained with GA₃ priming

Table 1: Germination and growth of rice seedlings as affected by varieties and seed priming treatments under laboratory conditions

Treatment	Germination (%)		Seedling length (cm)		Seedling dry weight (g/seedling)		Vigour index-I		Vigour index-II	
	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023
Varieties										
V ₁	81.0	75.4	30.2	24.3	0.096	0.103	2426	1842	7.78	7.77
V ₂	87.4	82.9	35.0	27.9	0.098	0.101	3088	2298	8.57	8.37
CD 5%	5.5	3.4	2.1	1.9	NS	NS	255	120	0.92	0.53
Priming treatments										
T ₁	77.8	73.0	29.2	23.2	0.095	0.100	2271	1693	7.39,11.0	7.30,13.0
T ₂	83.8	80.0	32.3	26.3	0.097	0.103	2707	2117	8.13	8.23
T ₃	86.3	80.2	32.4	25.8	0.095	0.103	2796	2059	8.20	8.25
T ₄	90.3	84.0	37.2	29.2	0.100	0.105	3359	2460	9.03	8.79
T ₅	83.5	79.0	32.2	25.8	0.098	0.100	2689	2017	8.18	7.90
T ₆	83.5	78.7	32.4	26.3	0.097	0.101	2706	2074	8.10	7.95
CD 5%	3.2	1.9	1.3	1.2	NS	NS	147	69	0.53	0.41

V₁: HKR 128; V₂: Pusa Basmati 1121; T₁: No priming (control); T₂: hydropriming; T₃: priming with ZnO nanoparticles @ 25 ppm; T₄: priming with GA₃ @ 50 ppm; T₅: priming with KNO₃ @ 1%; T₆: priming with ascorbic acid @ 100 ppm; NS: non-significant.

might be due to the additional supply of growth-promoting substance (gibberellin). The remaining priming treatments (T_2 , T_3 , T_5 , and T_6) were, however, at par with each other, improving the germination by 7.3–10.3% besides improving the seedling length (10.3–13.4%), seedling vigour-I (18.4–25.0%) and seedling vigour-II (8.2–13.0%) over the control during the two years. It has already been reported that hydropriming increases germination rate and seedling growth in rice by increasing the activity of α -amylase, which in turn results in better mobilization of carbohydrate reserves in the endosperm (Kata et al., 2014), thus shortening the metabolic lag time for seed germination (Khalid et al., 2019). In nanopriming, nanoparticles can penetrate the seed coat to enhance seed germination by increasing the activity of enzymes (such as amylase), proteins, and lipids in seeds to produce more soluble sugars, besides regulating the balance between Absciscic acid (ABA) and GA (Jisha et al., 2013; Sharma et al., 2021). Seed priming with KNO_3 might have improved the germination and seedling growth through improvement in water absorption and activation of reserve enzymes such as amylases, dehydrogenases and xylanase (Ali et al., 2020). Priming with ascorbic acid is reported to induce a wide range of coordinated responses, including regulation of hormone synthesis and epigenetic control of gene expression, allowing primed seeds to germinate and grow under adverse environment (Terzjaghi and DeTullio, 2023).

Dry weight of the seedlings, however, was not affected significantly due to priming with any agent. Interaction between varieties and priming treatments

was not found to be significant in respect of germination or seedling growth, indicating that the effect of all the priming treatments on both the varieties was more or less similar magnitude. Increase in the germination and growth of seedlings due to priming with gibberellic acid (Zhang et al., 2024), hydropriming (Matsunami et al., 2021), priming with KNO_3 (Dhillon et al., 2021), priming with ZnO nano-particles (Li et al., 2021), and priming with ascorbic acid (Zhang et al., 2024) have also been reported earlier.

Physiological indices and yield of crop

The data on mean value (averaged over the priming treatments) of various physiological indices, yield attributes and grain yield of the DSR crop grown in field conditions (Table 2) revealed that Pusa Basmati 1121 showed significantly higher plant stand (25.9–27.6 plants/m²), number of panicles/m² (311–324) and 1000-grain weight (28.2–29.6 g) than HKR 128 (23.3–24.2, 289–304 and 23.6–24.6, respectively) but number of grains/panicle (95.6–106.8) and grain yield (5939–6373 kg/ha) of HKR 128 was significantly higher than that of Pusa Basmati 1121 (57.7–63.6 and 4545–4946 kg/ha, respectively) during the two years. The higher yield of HKR 128, despite its lesser panicles/m² and lower 1000-grain weight, is attributable to its much higher number of grains/panicles. The data on physiological indices, viz., chlorophyll content (SPAD) and Chlorophyll fluorescence (CF), showed that variety Pusa Basmati 1121 exhibited significantly higher values of SPAD (both at flowering initiation and grain filling stages) and CF (only at flowering initiation) as compared to HKR 128.

Table 2: Plant stand, physiological indices, yield attributes and grain yield of DSR as affected by varieties and priming treatments under field conditions

Treatment	Number of plants/m ²		SPAD at flowering initiation		SPAD at grain filling		CF at flowering initiation (%)		CF at grain filling (%)		Number of panicles/m ²		Number of grains/panicle		1000-grain weight (g)		Grain yield (kg/ha)	
	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023	2022	2023
Varities																		
V ₁	24.2	23.3	34.6	34.9	31.3	33.1	0.57	0.59	0.56	0.54	289	304	95.6	106.8	23.6	24.7	5939	6373
V ₂	27.6	25.9	36.1	37.5	33.8	35.3	0.61	0.63	0.57	0.56	311	324	57.7	63.6	28.2	29.6	4545	4946
CD 5%	1.1	1.6	1.3	1.4	1.1	1.3	0.03	0.02	NS	NS	10	12	3.1	5.7	2.03	1.4	307	222
Priming treatments																		
T ₁	22.8	21.6	34.7	34.4	32.1	33.3	0.58	0.58	0.55	0.53	278	289	76.0	84.6	23.6	26.8	4883	5214
T ₂	26.6	24.4	36.5	35.7	34.5	33.6	0.60	0.61	0.56	0.54	303	318	77.8	85.3	25.8	27.2	5304	5728
T ₃	26.8	25.6	36.4	37.3	32.1	34.6	0.61	0.62	0.53	0.54	303	317	77.0	85.1	25.4	27.2	5308	5715
T ₄	27.1	26.0	34.2	36.6	32.2	35.0	0.57	0.59	0.57	0.55	304	320	76.4	85.5	27.1	27.2	5322	5770
T ₅	25.7	25.4	34.1	36.9	31.4	34.2	0.60	0.62	0.60	0.59	307	319	76.1	85.4	27.2	27.0	5312	5753
T ₆	26.4	24.6	36.4	36.3	33.1	34.4	0.61	0.63	0.57	0.56	307	320	76.7	85.2	26.4	27.2	5323	5777
CD 5%	2.5	2.0	NS	NS	NS	NS	NS	NS	NS	NS	16	20	NS	NS	NS	NS	378	385

SPAD: Chlorophyll content; CF: Chlorophyll fluorescence.

These varietal differences in physiological indices and yield attributes might be due to differences in the genetic constitution of the varieties.

All the priming treatments significantly increased the mean (averaged over varieties) number of established seedlings or plant stand (number of plants/m²), number of panicles/m² and grain yield of the field crop over the unprimed control (T₁) under dry direct seeded (aerobic) conditions (Table 2). Where as, the number of grains and 1000-grain weight of the crop were not influenced significantly due to priming. A similar trend was recorded during both years. The increase in the number of plants/m² (stand establishment) over the unprimed control was to the tune of 12.8-15.8% during 2022 and 11.7-12.7% during 2023, whereas the increase in the number of panicles/m² was 11.3-12.8% during 2022 and 10.0-11.8% during 2023. The priming, irrespective of the priming agents, also increased grain yield of the DSR crop significantly (10.5-12.2% in 2022 and 9.8-11.1% in 2023) as compared to unprimed control (T₁). However, all the priming treatments, including the hormonal priming, were at par with each other in respect of their favourable effect on the stand establishment, panicle number/m² and grain yield. This indicated that the additional benefit of hormonal priming over the other priming agents was absent under the field conditions, probably due to prohibitive soil conditions. The results clearly indicate that the priming increased the grain yield of DSR primarily by improving its stand establishment (number of plants/m²) in the field, which resulted in more effective tillers or panicles/m² and ultimately higher yield. Halopriming with 2% potassium nitrate (KNO₃) and hormonal priming with 50 ppm GA₃ have also shown good potential in boosting crop establishment and yield of direct-seeded rice in soils of western IGP (Dhillon *et al.*, 2021). The increase in yield of DSR due to enhancement of germination, stand establishment and number of panicles/m² by hydropriming (Mahajan *et al.*, 2011), nanopriming with Zn nanoparticles (Adhikary *et al.*, 2022), and priming with ascorbic acid (Zhang *et al.*, 2024) has also been reported earlier. Physiological indices, viz., SPAD and CF both at flowering and maturity did not vary significantly due to the priming treatments. Mahajan *et al.*, (2011) also did not find any variation in SPAD values due to priming. There was no significant interaction between the varieties and priming treatments with respect to their effect on yield attributes and the yield of DSR crop in the field, indicating that the effect of priming in both varieties was of a similar nature.

Conclusion

Priming with all the agents significantly improved the germination, seedling growth, and seedling vigour under laboratory conditions and also improved the plant stand, number of panicles/m² and ultimately grain yield of dry DSR crop under field conditions over the unprimed control. Among the priming agents, priming with GA₃ @ 50 ppm showed the highest germination and seedling length under laboratory conditions. However, under field conditions all the priming agents were at par in respect of their favourable effect on plant stand, yield attributes, and yield of DSR crop.

Acknowledgments

The authors acknowledge the research facilities provided at Rice Research Station, Kaul (Kaithal) of Chaudhary Charan Singh, Haryana Agricultural University, Hisar, India. The contributions of several scientists in field and laboratory studies at the place of experimentation are also greatly acknowledged.

References

- Abdul-Baki AA and JD Anderson (1973) Vigour determination in soybean seed by multiple criteria. *Crop Sci.* 13: 630-633. <https://doi.org/10.2135/cropsci1973.0011183X001300060013x>
- Adhikary S, B Biswas, D Chakraborty, J Timsina, S Pal, JC Tarafdar, S Banerjee, A Hossain and S Roy (2022) Seed priming with selenium and zinc nanoparticles modifies germination, growth, and yield of direct-seeded rice (*Oryza sativa* L.). *Sci. Rep.* 12:7103. <https://doi.org/10.1038/s41598-022-11307-4>
- Alam K, RW Bell, M Hasanuzzaman, N Salatin, MH Rashid, N Akter, S Akhter, MS Islam, S Islam, S Naznin, MFA Anik, Md. MRB Apu, HB Saif, MJ Alam and MF Khatun (2020) Rice (*Oryza sativa* L.) establishment techniques and their implications for soil properties, global warming potential mitigation and crop yields. *Agronomy* 10: 888. <https://doi.org/10.3390/agronomy10060888>
- Ali AS and AA Elozeiri (2017) Metabolic processes during seed germination. In: JC Jimenez Lopez (ed.) *Advances in Seed Biology*. In Tech, London, UK. <https://doi.org/10.5772/intechopen.70653>
- Ali LG, R Nulit, MH Ibrahim and CYS Yien (2020) Enhancement of germination and early seedling growth of rice (*Oryza sativa* var. FARO44) by seed priming under normal and drought-stressed conditions. *J. Plant Nutr.* 43: 1579-1593. <https://doi.org/10.1080/01904167.2020.1739298>
- Asma J, D Subrahmanyam and D Krishnaveni (2023) The global lifeline: A staple crops sustaining two-thirds of the world's population. *Agric. Int. J.* 2: 15-18. <https://doi.org/10.51470/AGRI.2023.2.3.15>
- Brar AS, GS Buttar, D Jhanji, N Sharma, KK Vashist, SS Mahal, JS Deol and G Singh (2015) Water productivity, energy and economic analysis of transplanting methods with different

- irrigation regimes in Basmati rice (*Oryza sativa* L.) under north-western India. *Agric. Water Manag.* 158: 189-195. <https://doi.org/10.1016/j.agwat.2015.04.018>
- Dhillon BS, V Kaur, P Sagwal, N Kaur, GS Mangat and S Singh (2021) Seed priming with potassium nitrate and gibberellic acid enhances the performance of dry direct seeded rice (*Oryza sativa* L.) in north-western India. *Agron.* 11: 849. <https://doi.org/10.3390/agronomy11050849>
- ISTA (2019) *International Rules for Seed Testing for 2019*, Full issue-i-19-8 (300), International Seed Testing Association. Zürichstr. 50, CH-8303 Bassersdorf, Switzerland. Chapter 5, pp 5-53.
- Jisha KC, K Vijayakumari and JT Puthur (2013) Seed priming for abiotic stress tolerance: An overview. *Acta Physiol. Plant.* 35: 1381-1396. <https://doi.org/10.1007/s11738-012-1186-5>
- Kata LP, M Bhaskaran and U Ranganathan (2014) Influence of priming treatments on stress tolerance during seed germination of rice. *Int. J. Agric. Environ. Biotechnol.* 7: 225-232.
- Khalid MF, S Hussain, MA Anjum, S Ejaz, M Ahmad, M Jan, S Zafar, I Zakir, MA Ali, N Ahmad, MJ Rao and S Ahmad (2019) Hydropriming for plant growth and stress tolerance. In: M Hasanuzzaman and V Fotopoulos (eds) *Priming and Pretreatment of Seeds and Seedlings*. Springer, Singapore, pp 373-384
- Li Y, L Liang, W Liu, U Ashraf, L Ma, X Tang, S Pan, H Tian and Z Mo (2021) ZnO nanoparticle-based seed priming modulates early growth and enhances physio-biochemical and metabolic profiles of fragrant rice against cadmium toxicity. *J. Nanobiotechnology* 19: 75. <https://doi.org/10.1186/s12951-021-00820-9>
- Mahajan G, RS Sarlach, S Japinder and MS Gill (2011) Seed priming effects on germination, growth and yield of dry direct-seeded rice. *J. Crop Improv.* 25: 409-417. <https://doi.org/10.1080/15427528.2011.576381>
- Mahender A, A Anandan and SK Pradhan (2015) Early seedling vigour, an imperative trait for direct-seeded rice: An overview on physio-morphological parameters and molecular markers. *Planta* 241: 1027-1050. <https://doi.org/10.1007/s00425-015-2273-9>
- Matsunami M, H Hayashi, M Murai-Hatano and J Ishikawa-Sakurai (2021) Effect of hydropriming on germination and aquaporin gene expression in rice. *Plant Growth Regul.* 97: 263-270. <https://doi.org/10.1007/s10725-021-00725-5>
- Monajjem S, E Soltani, E Zainali, M Esfahani, F Ghaderi-Far, MH Chaleshtori and A Rezaei (2023) Seed priming improves enzymatic and biochemical performances of rice during seed germination under low and high temperatures. *Rice Sci.* 30: 335-347. <https://doi.org/10.1016/j.rsci.2023.03.012>
- Pagano A, A Macovei and A Balestrazzi (2023) Molecular dynamics of seed priming at the crossroads between basic and applied research. *Plant Cell Rep.* 42: 657-688. <https://doi.org/10.1007/s00299-023-02988-w>
- Paul S, S Dey and R Kundu (2022) Seed priming; an emerging tool towards sustainable agriculture. *Plant Growth Regul.* 97: 215-234. <https://doi.org/10.1007/s10725-021-00761-1>
- Rhaman MS, S Imran, F Rauf, M Khatun, CC Baskin, Y Murata and M Hasanuzzaman (2020) Seed priming with phytohormones: An effective approach for the mitigation of abiotic stress. *Plants* 10: 37. <https://doi.org/10.3390/plants10010037>
- Saha S, S Munda, A Mahapatra and BS Satapathy (2022) Prospects of direct-seeded rice under changing rice environment – present scenario and future needs. In: Bhattacharya et al., (eds) *Climate Resilient Technologies for Rice Based Production Systems in Eastern India*. ICAR-National Rice Research Institute, Cuttack, Odisha, pp 86-101.
- Sandhu N, S Yadav, V Singh and A Kumar (2021) Effective crop management and modern breeding strategies to ensure higher crop productivity under direct seeded rice cultivation system: A review. *Agronomy* 11: 1264. <https://doi.org/10.3390/agronomy11071264>
- Sandhu N, RB Yadaw, B Chaudhary, H Prasai, K Iftekharuddaula, C Venkateshwarlu, A Annamalai, P Xangsayasane, KR Battan, M Ram, MTS Cruz, P Pablico, PC Maturan, KA Raman, M Catolos and A Kumar (2019) Evaluating the performance of rice genotypes for improving yield and adaptability under direct seeded aerobic cultivation conditions. *Front. Plant Sci.* 10: 159. <https://doi.org/10.3389/fpls.2019.00159>
- Sharma D, S Afzal and NK Singh (2021) Nanopriming with phytosynthesized zinc oxidenanoparticles for promoting germination and starch metabolism in rice seeds. *J. Biotechnol.* 336: 64-75. <https://doi.org/10.1016/j.jbiotec.2021.06.014>
- Terzjaghi M and MC De Tullio (2023) Ascorbic acid in seeds, priming and beyond. *Seeds* 4: 421-435. <https://doi.org/10.3390/seeds2040032>
- Wang Y, Y Wang, R Yang, F Wang, J Fu, W Yang, T Bai, S Wang and H Yin (2021) Effects of gibberellin priming on seedling emergence and transcripts involved in mesocotylelongation in rice under deep direct-seeding conditions. *J. Zhejiang Uni. Sci. B.* 22:1002-1021. <https://doi.org/10.1631/jzus.b2100174>
- Waqas M, NE Korres, D Khan, A Nizami, F Deeba, I Ali and H Hussain (2019) Advances in the concept and methods of seed priming. In: MM Hasanuzzaman and V Fotopoulos V (eds) *Priming and Pretreatment of Seeds and Seedlings: Implication in Plant Stress Tolerance and Enhancing Productivity in Crop Plants*. Springer Nature Singapore Pvt. Ltd., Singapore, pp 11-39
- Zhang K, MN Khan, Z Khan, T Luo, B Zhang, J Bi, L Hu and L Luo (2024) Seed priming with ascorbic acid and spermidine regulated auxin biosynthesis to promote root growth of rice under drought stress. *Front. Plant Sci.* 15: 1482930. <https://doi.org/10.3389/fpls.2024.1482930>
- Zheng M, Y Tao, S Hussain, Q Jiang, S Peng, J Huang, K Cui and L Nie (2016) Seed priming in dry direct-seeded rice: Consequences for emergence, seedling growth and associated metabolic events under drought stress. *Plant Growth Regul.* 78: 167-178. <https://doi.org/10.1007/s10725-015-0083-5>

RESEARCH ARTICLE

Morphological Diversity Analysis in Brinjal (*Solanum melongena*) Germplasm

TH Singh^{1*}, Sushil Pandey², Chitra D Pandey² and Suresh R Yerasu³

Abstract

An experiment was conducted at the Division of Vegetable Crops, ICAR-Indian Institute of Horticultural Research, aimed to characterize 480 indigenous and exotic brinjal accessions received from the National Genebank, ICAR-NBPGR, New Delhi. A lot of variability was observed for agro-morphological traits recorded such as mean plant height (88.80 cm), plant spread (53.25 cm), number of primary branches 5.0, leaf length 19.26 cm, leaf width 13.03 cm, days to fifty per cent flowering 50.00, fruit length 11.18 cm, fruit breadth 4.83 cm, number of fruits per plant 14, average fruit weight 80.15g, yield per plant 1.13kg and 100 seed weight 0.5 g. The Principal Component Analysis (PCA) indicated 100% of the total variation based on 12 phenotypic traits. There was a significant positive relationship found among the fruit characters. The majority of the characters studied had high PCV, GCV, heritability, and GAM levels. The brinjal accessions, such as EC316273-X, EC-396492, EC-467275, IC-0090851, IC-0090891-Z, IC-0090858, IC-0090898, EC-384615, IC-0090762, IC-0090774 and IC-090922-X had shown high variability for quantitative characters.

Keywords: Brinjal, Correlation, Genetic advance, Genetic variation, Principal component analysis

¹ICAR-Indian Institute of Horticultural Research, Bengaluru, Karnataka, 560089, India.

²ICAR-National Bureau of Plant Genetic Resources, New Delhi, 110012, India.

³ICAR-Indian Institute of Vegetable Research, Varanasi, Uttar Pradesh, 221305, India.

***Author for correspondence:**

Singh.HT@icar.gov.in

Received: 13/11/2024

Revised: 19/12/2025

Accepted: 12/02/2026

How to cite this article: Singh TH, S Pandey, CD Pandey and SR Yerasu (2026) Morphological Diversity Analysis in Brinjal (*Solanum melongena*) Germplasm. *Indian J. Plant Genet. Resour.* 39(2): 74-82.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.176046>

Introduction

India has a strong basis for brinjal research in both public and private sector institutes involved in brinjal breeding. Plant Genetic Resources (PGR) exchange and its utilization in different crop improvement programs are continuous processes. To fulfil future demands caused by the introduction of new diseases, climate change, and increased demands for food and nutritional security, there is an ongoing quest for different new PGRs (De Jonge, 2009). The brinjal (*Solanum melongena* L., $2n = 24$) is a member of the Solanaceae family and is a highly valued vegetable crop cultivated in tropical and sub-tropical regions of the world. It originated in India (the Indo-Burma region is the primary centre of origin), and China is its secondary centre of origin. In India, brinjal is grown in an area of 676.70 thousand hectares, with production of 12.78 million metric tonnes (MT) and with a productivity of 18.90 MT/ha (India Statagri, 2023). In India, major brinjal-producing states are West Bengal, Odisha, Gujarat, Uttar Pradesh, Madhya Pradesh, Bihar, Chhattisgarh, Karnataka, Andhra Pradesh and Telangana. In Karnataka, it is cultivated in an area of 10,387 ha, producing 259,033 MT with a productivity of 24.94 MT/ha (Department of Horticulture, GOK, 2021-22). Brinjal ranked fourth among 25 different crops in terms of seed requests received by ICAR-NBPGR, New Delhi, from 2014 to 2019 (Tyagi *et al.*, 2021). ICAR-NBPGR collaborates with national

organizations for evaluation, multiplication, characterization, documentation, and conservation of PGR, which ensures their sustainable utilization in breeding programme.

The characterization of accessions involves recording and collecting data on crucial phenotypic traits, which facilitates grouping of accessions, formation of a core collection and identification of promising germplasm for crop breeding programs. Such studies enhance comprehension of the collection's composition and genetic diversity. Morphological traits play an important role in germplasm characterization, classification, and genotyping, which is useful in identifying elite genotypes for breeding purposes (Saikia et al., 2021). Understanding genetic variation is essential for effectively utilizing PGR in breeding programs. Yield is a complex trait influenced by various associated plant characteristics, character associations, assessing heritability and genetic progress. The heritability and genetic progress of different traits are pivotal factors in the selection processes. This study aimed to phenotypically characterize existing germplasms for yield and yield-related traits, and to evaluate their breeding potential. These insights will enhance our understanding of the brinjal germplasm potential and its suitability for future breeding programs.

Material and Methods

An experiment was conducted at the ICAR-Indian Institute of Horticultural Research, located in Bengaluru, Karnataka, positioned at latitude 13°8' 1.51"N and longitude 77°29' 27.32"E during the *kharif* 2022.

Plant materials and data collection

A group of 480 brinjal accessions from the National Genebank, ICAR-NBPGR, New Delhi, was used in the experiment (Table 1). Of the 480 brinjal accessions evaluated, one accession viz., *S. viarum*, was a wild relative, whereas the remaining 479 were cultivated *Solanum melongena* L. accessions. The study also included five standard checks, namely, IC628873 (BR-14), Kashi Prakash, Kashi Taru, Arka Kusumakar, and Arka Nidhi. Observations were recorded on the following 12 parameters viz., Plant height (cm), Plant spread (cm), Number of primary branches per plant, Leaf length (cm), Leaf width (cm), Days to fifty percent flowering, Fruit length (cm), Fruit breadth (cm), Number of fruits per plant, Yield per plant (kg), Average fruit weight (g), and 100-seed weight (g). Five plants of each

Table1: List and source of the brinjal accessions used in the study

S. No.	Accessions	S. No.	Accessions
1	IC-0099735	45	IC-136345
2	IC-0099737	46	IC-136347
3	IC-0099750	47	IC-136359
4	IC-0099755	48	IC-136364
5	IC-104088	49	IC-136367
6	IC-0111000	50	IC-136380
7	IC-111001	51	IC-136383
8	IC-0111003	52	IC-136367
9	IC-111004	53	IC-136445
10	IC-111005	54	IC-136551
11	IC-111008	55	IC-0099735
12	IC-111018	56	IC-0099737
13	IC-111027	57	IC-0099750
14	IC-111033	58	IC-136564
15	IC-111037	59	IC-137681
16	IC-111043	60	IC-137702
17	IC-111052	61	IC-137751
18	IC-111069	62	IC-144013
19	IC-111073	63	IC-144015
20	IC-111081	64	IC-144080
21	IC-111082	65	IC-169073
22	IC-111323	66	IC-169089
23	IC-111337	67	IC-201139
24	IC-1112347	68	EC-284270
25	IC-112732	69	EC-304994
26	IC-126640	70	EC-305006
27	IC-126649	71	IC-310883
28	IC-126721	72	IC-310884
29	IC-126829	73	IC-310883
30	IC-126860	74	IC-310884
31	IC-126916	75	IC-310885
32	IC-126860	76	EC-316230
33	IC-601513	77	EC-316242
34	IC-136142	78	EC-316258
35	IC-136206	79	EC-316283
36	IC-136226	80	EC-316315
37	IC-136228	81	EC-316317
38	IC-136226	82	EC-378654
39	IC-136281	83	EC-378820
40	IC-136287	84	EC-379246
41	IC-136311	85	EC-382577
42	IC-136312	86	EC-382579
43	IC-136321	87	EC-383345
44	IC-136312	88	EC-384619

S. No.	Accessions	S. No.	Accessions	S. No.	Accessions	S. No.	Accessions
89	EC-467271	135	IC-111302-X	181	IC-136248-1X	227	IC-345272
90	EC-467273	136	IC-111440-X	182	IC-0239393	228	IC-345275
91	EC-467277	137	IC-111448-X	183	IC-0603202	229	IC-345287
92	IC-510491	138	IC-112713	184	IC-136272	230	IC-345290
93	IC-265644	139	IC-112741	185	IC-136329-X	231	IC-345309
94	IC-136552	140	IC-112744	186	IC-136438-X	232	IC-345326
95	IC-136564	141	IC-112992-X	187	IC-136556	233	IC-345327
96	IC-137681	142	IC-113000-X	188	IC-136561	234	IC-345328
97	IC-320865	143	IC-126912-X	189	IC-137682	235	IC-345330
98	IC-362379	144	IC-135942-X	190	IC-137761	236	IC-345333
99	IC-362380	145	IC-136248-1	191	IC-144109-X	237	IC-345335
100	IC-362381	146	IC-136272-X	192	IC-201132	238	IC-345359
101	IC-362382	147	IC-136329	193	IC-285497	239	IC-345363
102	IC-362383	148	IC-136438	194	EC-305052	240	IC-345368
103	IC-362384	149	IC-136556-X	195	EC-316273-X	241	IC-345370
104	IC-362385	150	IC-136561-X	196	IC-321070-X	242	IC-345372
105	IC-362386	151	IC-137682-X	197	IC-422570	243	IC-345385
106	IC-362387	152	IC-137761-X	198	IC-422571	244	IC-361839
107	IC-362388	153	IC-144109	199	IC-422573	245	IC-361840
108	IC-362389	154	IC-201132-X	200	IC-422574	246	IC-361841
109	IC-362390	155	IC-256208-X	201	IC-422575	247	IC-361842
110	IC-362391	156	EC-305052-X	202	IC-422576	248	IC-369610
111	IC-362392	157	EC-316224-X	203	IC-422577	249	IC-373370
112	IC-362394	158	EC-316273-X2	204	IC-422578	250	IC-374777
113	IC-362395	159	EC-316285-X	205	IC-422579	251	IC-383085
114	IC-362396	160	IC-332498	206	IC-422580	252	IC-383099
115	IC-362397	161	IC-332501	207	IC-422581	253	IC-383101
116	IC-362399	162	IC-332502	208	IC-422582	254	IC-383102
117	IC-371739	163	IC-332503	209	IC-422583	255	IC-383103
118	IC-588090	164	IC-332504	210	IC-422584	256	IC-383119
119	IC-604614	165	IC-332508	211	IC-422585	257	IC-383137
120	IC-90870	166	IC-332510	212	IC-422586	258	IC-383158
121	IC-90893	167	IC-332513	213	IC-422587	259	IC-383170
122	IC-90955	168	IC-332514	214	IC-422588	260	IC-383175
123	IC-90967	169	IC-332515	215	IC-541227	261	IC-383190
124	IC-90980	170	IC-89815-X	216	IC-258028	262	IC-383195
125	IC-324556	171	IC-0603204	217	IC-260999	263	IC-383695-X
126	IC-324775	172	IC-0603970	218	IC-311233	264	IC-395714
127	IC-324870	173	IC-99747	219	IC-333519	265	IC-413668
128	IC-89900-X	174	IC-111089-X	220	IC-333523	266	IC-413717
129	IC-90975-X	175	IC-111308-X	221	IC-333527	267	IC-414993
130	IC-90979	176	IC-111431-X	222	IC-333582	268	IC-415033
131	IC-99630	177	IC-111440	223	IC-333607	269	IC-588104
132	IC-99702-X	178	IC-112741-X	224	IC-345255	270	IC-588106
133	IC-99747-X	179	IC-126912	225	IC-345270	271	IC-604481
134	IC-111089	180	IC-127070	226	IC-345271	272	EC-169075

S. No.	Accessions	S. No.	Accessions	S. No.	Accessions	S. No.	Accessions
273	EC-169080	319	IC-0089868	365	IC-0090118-X	411	IC-602927
274	EC-304976	320	IC-0089880	366	IC-0090119	412	IC-0090905-X
275	EC-304983	321	IC-0089885	367	IC-0090122-X	413	IC-0090907
276	EC-305027	322	IC-0089897	368	IC-0090124	414	IC-0090909
277	EC-305046	323	IC-0089898	369	IC-0090126	415	IC-0090913
278	EC-305129	324	IC-126630	370	IC-0090128	416	IC-090914
279	EC-316213	325	IC-126637	371	IC-0090130	417	IC-090922-X
280	EC-316223	326	IC-600305	372	IC-0090137	418	IC-0090923
281	EC-316234	327	IC-600317	373	IC-0090140	419	IC-0090927
282	EC-316237	328	IC-600330	374	IC-602915	420	IC-0090928
283	EC-316244	329	IC-600331	375	IC-602920	421	IC-0090931
284	EC-316246	330	IC-600788	376	IC-0090142	422	IC-0090932-X
285	EC-316305	331	IC-601079	377	IC-0090144	423	IC-0090935
286	EC-316309	332	IC-601091	378	IC-0090146	424	IC-0090936
287	EC-316310	333	IC-601098	379	IC-0090150	425	IC-0090938
288	EC-377624	334	IC-601154	380	IC-0090156	426	IC-0090939
289	EC-378655	335	IC-601155	381	IC-0090158	427	IC-0090940
290	EC-379244	336	IC-601415	382	IC-0090159-X	428	IC-0090942
291	EC-382582	337	IC-601474	383	IC-0090621	429	IC-0090946
292	EC-383343	338	IC-602528	384	IC-0090750	430	IC-0090951
293	EC-383348	339	IC-602632	385	IC-0090762	431	IC-0090952-X
294	EC-384550	340	IC-602912	386	IC-0090764	432	IC-0090962
295	EC-384556	341	IC-0089918-X	387	IC-0090774	433	IC-0090963-X
296	EC-384565	342	IC-0089928	388	IC-0090775	434	IC-0090965
297	EC-384606	343	IC-0089932	389	IC-0090783-X	435	IC-0090970
298	EC-384615	344	IC-0089935	390	IC-0090785	436	IC-0090972
299	EC-384618	345	IC-0089947	391	IC-0090790	437	IC-0090973
300	EC-384628	346	IC-0089957	392	IC-0090791	438	IC-0090974
301	EC-393239	347	IC-0089974-Z	393	IC-0090804	439	IC-90974
302	EC-396466	348	IC-0089998	394	IC-0090810	440	IC-90981
303	EC-396487	349	IC-0090030	395	IC-0090812-X	441	IC-90983-X
304	EC-396492	350	IC-0090031	396	IC-0090844-X	442	IC-090984
305	EC-467268	351	IC-0090032	397	IC-0090846-Z	443	IC-92719
306	EC-467272	352	IC-0090035	398	IC-0090851	444	IC-94209
307	EC-467274	353	IC-0090041	399	IC-0090858	445	IC-0099614
308	EC-467275	354	IC-0090043	400	IC-0090860-X	446	IC-99621
309	EC-467276	355	IC-0090064	401	IC-0090869	447	IC-99630-X
310	IC-0009081	356	IC-0090087-X	402	IC-0090871	448	IC-99635
311	IC-0074243-X	357	IC-0090100	403	IC-0090872	449	IC-0099640
312	IC-0089814-X	358	IC-0090101	404	IC-0090874	450	IC-0099643
313	IC-0089826	359	IC-0090102	405	IC-0090891-Z	451	IC-99645
314	IC-0089831	360	IC-0090110	406	IC-0090895	452	IC-99656-X
315	IC-0089832	361	IC-0090113-X	407	IC-0090898	453	IC-99658-X
316	IC-0089841	362	IC-0090115-X	408	IC-0090899	454	IC-99660
317	IC-0089845	363	IC-0090116	409	IC-0090901	455	IC-0099661
318	IC-0089846	364	IC-0090117	410	IC-0090902	456	IC-99670-X

S. No.	Accessions	S. No.	Accessions
457	IC-99675	472	IC-421189
458	IC-099686	473	IC-446754
459	IC-99688	474	IC-201136
460	IC-99695-X	475	IC-127074
461	IC-099699	476	IC-397558
462	IC-099703	477	IC-137701
463	IC-099708	478	IC-127115-X
464	IC-99736	479	IC-99703-X
465	IC-99740	480	IC-99758-X
466	IC-99743-X	C1	IC-628873 (BR-14)
467	IC-99744	C2	Kashi Prakash
468	IC-99748	C3	Kashi Taru
469	IC-446729	C4	Arka Kusumakar
470	IC-439263	C5	Arka Nidhi
471	IC-421199		

accession were tagged/selected for recording all the observations.

Experimental design

An augmented randomized complete block design (ABD) was employed as the experimental design to cultivate the accessions, wherein the checks were replicated in each block. A total of 480 accessions were grown in 6 blocks, with 80 accessions in each block. 35-day-old plants were transplanted to the main field with 100 cm × 60 cm space between rows and plants. Recommended agriculture practices were followed to raise a good crop.

Statistical analysis

The R package 'augmented RCBD' was employed to calculate adjusted averages for all considered traits across all genotypes (Aravind *et al.*, 2018). Genetic variability parameters, frequency distributions, and descriptive statistics were computed in R software. Phenotypic variance, genotypic variance, phenotypic coefficient of variation (PCV), genotypic coefficient of variation (GCV), broad-sense heritability (h^2) (Lush, 1940), and genetic advances (GA) (Johnson *et al.*, 1955) were estimated as:

$$\text{Genotypic variance } (\sigma_g^2) = \bar{P} - \bar{e}$$

where,

$\bar{P}$ Phenotypic variance = Sum of squares of test treatments (genotypes)

$\bar{e}$ Error Variance = Sum of squares of residuals (error)

$$\text{Phenotypic Coefficient of Variation (PCV)} = \frac{\text{Phenotypic Variance}}{\sqrt{\text{mean}}} \times 100$$

$$\text{Genotypic Coefficient of Variation (GCV)} = \frac{\text{Genotypic Variance}}{\sqrt{\text{mean}}} \times 100$$

$$\text{Broad sense heritability } (h^2) = \frac{\sigma_g^2}{\sigma_p^2}$$

$$\text{Genetic Advance (GA)} = k \times \sigma_g \times \frac{h^2}{100}$$

where k represents the selection intensity, and σ_g denotes the genotypic standard deviation.

Correlation analysis was conducted in the R package 'metan' (Olivoto and Lúcio, 2020), while PCA was done with the R package 'FactoMineR' (Le *et al.*, 2008).

Results and Discussion

Genetic parameters

Accurate estimation and reporting of genetic diversity within germplasm are crucial for its effective utilization (Islam *et al.*, 2004). Genetic diversity is often assessed using morphological traits and aid in cultivar development because of their simplicity and cost-effectiveness (Fufa *et al.*, 2005). In the present study, a significant range of variance was observed among the twelve characters analysed in the brinjal accessions. ANOVA results indicated that the variation for all characters was highly significant, highlighting inherent genetic differences. Each characteristic exhibited a wide range (Table 2), with the coefficient of variation (CV %) ranging from 0.23 for average fruit weight to 31.58 for fruit breadth. This study focused solely on morphological characteristics for diversity analysis. Previous research has utilized morphological features for diversification studies in brinjal (Savan kumar *et al.*, 2018; Khan *et al.*, 2023; Nagar *et al.*, 2024). The phenotypic coefficient of variance (PCV) for all traits exceeded the genotypic variance (GCV), suggesting environmental effect on these traits, except for days to 50% flowering, leaf width (cm), leaf length (cm), and the number of primary branches per plant. The remaining characters displayed high phenotypic and genotypic coefficients of variation (PCV and GCV). Most of the characters exhibited high broad-sense heritability, excluding days to 50% flowering and the number of primary branches per plant. Heritability, along with genetic advance, provides a more accurate estimate of selection value (Johnson *et al.*, 1955). All the characters exhibited a high genetic advance (GA) percentage except for the number of primary branches per plant (7.45) and days to 50% flowering (9.05). The minimum GCV and PCV observed for the number of primary branches per plant and days to 50% flowering indicate

Table 2: Genetic variability parameters of the brinjal germplasm

Trait	Mean	Range	CV %	GCV	GCV category	PCV	PCV category	ECV	hBS(h ²) (%)	hBS category	GA	GAM (%)	GAM category
PH	88.80	53-183	2.48	28.42	High	28.53	High	2.48	99.24	High	51.80	58.41	High
PS	53.25	40-132	4.94	24.94	High	25.42	High	4.91	96.27	High	27.06	50.49	High
NPB	5.0	3.0-7.0	16.87	8.22	Low	18.72	Medium	16.82	19.29	Low	0.37	7.45	Low
LL	19.26	2.2-28.2	19.24	15.16	Medium	15.93	Medium	4.88	90.63	High	5.73	29.78	High
LW	13.03	1.80-23.50	12.81	19.08	Medium	21.33	High	9.54	80.01	High	4.51	35.21	High
DFF	50.00	42.0-58.0	6.19	6.20	Low	8.76	Low	6.19	50.06	Medium	4.55	9.05	Low
FL	11.18	4.30-20.13	13.32	22.87	High	26.59	High	13.56	74.00	High	4.46	40.59	High
FB	4.83	1.80-10.10	31.58	26.23	High	27.36	High	31.90	68.42	High	3.25	38.86	High
FP	14.00	6-64	0.29	43.79	High	43.79	High	0.29	99.00	High	12.78	90.33	High
YPP	1.13	0.2-5.1	2.82	52.04	High	52.11	High	2.81	98.51	High	1.21	107.20	High
FW	80.15	11-188	0.23	33.22	High	33.22	High	0.23	97.12	High	55.21	68.52	High
100SW	0.5	0.2-0.5	12.94	25.42	High	28.51	High	12.91	79.50	High	0.22	46.76	High

PH = Plant height (cm), PS=Plant spread (cm), NPB = Number of primary branches, LL=Leaf length (cm), LW=Leaf width(cm), DFF = Days to fifty per cent flowering, FL=Fruit length (cm), FB=Fruit breadth (cm), FP= Number of fruits per plant, YPP = Yield per plant (kg), FW = Average fruit weight (g) and 100SW= 100 seed weight (g).

limited variation in these traits. This led to low heritability and low GA as a percentage of the mean for these two traits. In contrast, the remaining characteristics showed high GCV and broad-sense heritability, resulting in high GA as a percentage of the mean (Table 2). The variation found among the genotypes for the studied characters can be attributed to genetic diversity and differences in adaptability (Nagar *et al.*, 2024). Variability and genetic diversity among genotypes, combined with higher heritability, provide greater genetic gains in breeding programs (Balasubramaniyam *et al.*, 2021; Kumar *et al.*, 2021; Uddin *et al.*, 2021; Soumya *et al.*, 2023).

Yield frequency distribution

Yield plays a significant role in crop improvement programmes. The average yield per plant among the germplasm varied from 0.2 to 5.1 kg. Accessions such as EC-316273-X (2.53 kg), EC-396492 (2.55 kg), EC-467275(2.60 kg), IC-0090851(3.19 kg), IC-0090891-Z(3.19 kg), IC-0090858(3.20 kg), IC-0090898 (3.40 kg), EC-384615 (3.76 kg), IC-0090762 (4.10 kg), IC-0090774 (4.50 kg) and IC-090922-X (5.10 kg) had superior fruit yields similar findings were also reported by Soumya *et al.*, (2023) and Nagar *et al.*, (2024).

Correlation

Correlation analysis is a crucial aspect of any breeding program as it helps in understanding how associated characters influence selection and the genetic gain of complex traits like yield. Pearson correlation

coefficients were calculated to estimate the relationships among the characteristics (Fig. 1). Yield per plant has significant positive correlation with number of fruits per plant ($r = 0.72$, $P < 0.01$) & average fruit weight ($r = 0.61$, $P < 0.001$). Sonagara *et al.*, (2022) also reported a strong and significant relationship between fruit yield and the number of fruits per plant. Yield was negatively correlated with Leaf length ($r = -0.09$, $P < 0.05$, Plant spread ($r = -0.13$, $P < 0.01$) & Plant height ($r = -0.15$, $P < 0.001$). Plant height showed a




Fig. 1: Graphical depiction of correlation among the quantitative characteristics of the brinjal germplasm.

positively significant correlation with the plant spread ($r = 0.39$, $P < 0.001$), Leaf length ($r = 0.14$, $P < 0.01$) & 100 seed weight ($r = 0.09$, $P < 0.05$) and negatively correlated with average fruit weight ($r = -0.11$, $P < 0.05$). Plant spread exhibited a positive correlation with leaf length ($r = 0.10$, $P < 0.05$) these findings were in agreement with Balasubramaniyam *et al.*, (2021), Kumar *et al.* (2021), Sonagara *et al.*, (2022), and Khan *et al.*, (2023).

Principal component analysis

Principal component analysis (PCA) was employed to assess diversity on a multivariate scale. PCA simplifies the relationships among a large set of characters into a few components, retaining essential information from the original dataset. This method highlights the most influential characteristics contributing to the overall variation (Sanni *et al.*, 2008). In the present investigation, the initial five components with eigenvalues that were greater than 1.0 explained 56.90% of the variance among brinjal genotypes (Table 3). The first component

(PC1) accounted for 15.77% of the variation, primarily due to plant traits such as leaf length, leaf width, plant height, and the number of primary branches per plant. The second component (PC2) explained 12.50% of the variation, driven by plant spread, plant height, and leaf width. The third component (PC3) contributed 10.62% variation through the number of fruits per plant and leaf length (Fig. 2). The fourth component (PC4) showed 9.34% of the variation due to 100-seed weight and plant height, while the fifth component (PC5) contributes 8.66% of the variation mainly through average fruit weight. The first two PCs described 28.28% of the overall variation, with eigenvalues of 1.89 and 1.50, respectively. These components indicated that plant height, plant spread, the number of primary branches per plant, leaf length, and leaf width were the main discriminatory characteristics. Rahman *et al.*, (2014), Kumar *et al.*, (2016), and Savankumar *et al.*, (2018) found similar results, indicating that plant characteristics significantly contributed to the variation among brinjal germplasms.




Fig. 2: Two-dimensional graphical representations of component patterns along with different variables based on PC1 and PC2.

Table 3: Eigenvalues, proportion of the total variance represented by the first eight principal components, cumulative per cent variance and component loading of different characters in Brinjal

	PC1	PC2	PC3	PC4	PC5	PC6	PC7	PC8	PC9	PC10
Eigen Value	1.89	1.50	1.27	1.12	1.04	0.58	0.98	0.93	0.90	0.32%
variance Expressed	15.77	12.50	10.62	9.34	8.66	4.86	8.21	7.77	7.49	2.69
Cumulative variance expressed	15.77	28.28	38.90	48.24	56.90	97.31	65.11	72.87	80.37	100.00
PH	0.174	0.808	0.029	0.112	-0.063	-0.103	-0.051	-0.106	-0.201	0.037
PS	-0.041	0.837	-0.102	-0.059	-0.002	0.055	0.043	0.090	0.209	-0.120
NPB	0.167	0.015	0.004	0.024	-0.002	-0.012	-0.051	0.052	0.939	0.141
LL	0.910	0.017	0.109	0.035	0.052	0.050	-0.042	0.025	0.055	-0.043
LW	0.868	0.101	-0.161	-0.070	-0.071	-0.013	0.003	0.034	0.123	0.063
DF	-0.034	-0.005	-0.001	0.025	-0.003	0.001	0.992	0.039	-0.047	-0.087
FL	0.012	-0.074	-0.064	0.029	0.045	0.044	-0.090	0.006	0.137	0.972
FB	0.035	-0.034	0.020	-0.005	0.005	0.994	0.001	0.019	-0.009	0.042
FP	-0.035	-0.061	0.984	0.044	-0.016	0.020	-0.001	0.056	0.003	-0.061
YPP	0.053	-0.008	0.055	0.000	0.021	0.019	0.039	0.988	-0.047	0.005
FW	-0.012	-0.050	-0.015	0.030	0.995	0.005	-0.003	0.021	-0.002	0.043
100SW	-0.029	0.038	0.044	0.992	0.030	-0.005	0.025	0.000	0.021	0.028

PH = Plant height (cm), PS=Plant spread (cm), NPB = Number of primary branches, LL=Leaf length (cm), LW=Leaf width(cm), DF = Days to fifty per cent flowering, FL=Fruit length (cm), FB=Fruit breadth (cm), FP=Number of fruits per plant, YPP = Yield per plant (kg), FW = Average fruit weight (g) and 100SW= 100 seed weight (g).

Conclusion

Accessions such as IC-090922-X, IC-0090774, IC-0090762, EC-384615, IC-0090898, IC-0090858, IC-0090891-Z, IC-0090851, EC-467275, EC-396492, and EC-316273-X demonstrated strong contributions to yield and yield-related traits. These diverse germplasm accessions serve as reservoirs of important genes/alleles that help to get higher yield and can be utilized as foundational material in future breeding programs to develop new, improved varieties and advanced breeding lines in brinjal.

Acknowledgments

The authors are grateful to the Director of the ICAR-National Bureau of Plant Genetic Resources, New Delhi, for providing financial support through the CRP-Agro Biodiversity program, which enabled the ICAR-IIHR to acquire the germplasm for this study. The authors are also grateful to the Director and Head of the Division of Vegetable Crops, ICAR, Indian Institute of Horticultural Research, Bengaluru, for permitting the field experiments.

References

Aravind J, MS Sankar, PW Dhammaprakash and V Kaur (2018) AugmentedRCBD: Analysis of augmented randomized complete block designs. R package version 0.1.0. <https://aravind-j.github.io/augmentedRCBD>

Department of Horticulture, Government of Karnataka (2021-2022) Area and production of horticulture crops. <https://horticulturedir.karnataka.gov.in>

Balasubramaniam K, K HariPriya, TB Kumar and R Elangaimannan R (2021) Assessment of genetic variability, heritability and genetic advance in brinjal (*Solanum melongena* L.). *Plant Arch.* 21(1): 1784-1786.

De Jonge (2009) Plants, genes and justice: An enquiry into fair and equitable benefit-sharing. Unpublished PhD dissertation, Wageningen: Wageningen University. http://www.fao.org/agriculture/crops/thematic_sitemap/theme/seeds-pgr/conservation/en/.

Fufa H, PS Baenziger, BS Beecher, IDweikat, RA Graybosch and KM Eskridge (2005) Comparison of phenotypic and 1103 molecular marker-based classifications of hard red winter wheat cultivars. *Euphytica* 145: 133-146.

Islam FMA, S Beebe, M Munoz, J Tohme, RJ Redden and KE Basford (2004) Using molecular markers to assess the effect of introgression on quantitative attributes of common bean in the Andean gene pool. *Theor. Appl. Genet.* 108: 243-252.

Johnson HW, HF Robinson and RE Comstock (1955) Estimates of genetic and environmental variability in soybean. *Agron. J.* 47: 314-318.

Khan S, K Hussain, T Khan, LR Shah, G Ali, ZA Dar, A Gulzar, AA Malik and I Khan (2023) Genetic variability, correlation and path analysis for yield components in brinjal (*Solanum melongena* L.). *PharmaInnovation* 12(4): 773-781.

Kumar SR, T Arumugam and V Ulaganathan (2016) Genetic diversity study in eggplant germplasm by principal component analysis. *Sabao J. Breed. Genet.* 48(2): 162-171.

Le S, J Josse and F Husson (2008) FactoMineR: An R package for multivariate analysis. *J. Stat. Softw.* 25(1): 1-17.

Lush JL (1940) Intra-sire correlations or regressions of offspring on dam as a method of estimating heritability of

- characteristics. *Proc. Am. Soc. Anim. Nutr.* 1940: 293-301.
- Nagar KL, DK Rana, A Barela and S Rahangdale (2024) Study of genetic diversity and genetic advance in brinjal for morpho-economic traits (*Solanum melongena* L.). *Int. J. Environ. Clim. Change*. 14(3): 437-444.
- Olivoto T and AD Lúcio (2020) Metan: An R package for multi-environment trial analysis. *Methods Ecol. E.* 11: 783-789. <https://doi.org/10.1111/2041-210X.13384>.
- Rahman MO, MG Rabbani, R Yesmin and EJ Garvey (2014) Genetic diversity of brinjal (*Solanum melongena* L.) through multivariate analysis. *Int. J. Nat. Soc. Sci.* 1: 85-93.
- Rameshkumar D, RS Priya, BK Savitha, R Ravikesavan and NA Muthukrishnan (2021) Correlation and path analysis studies on yield and yield components in brinjal (*Solanum melongena* L.). *Electron. J. Plant Breed.* 12(1): 249-252.
- Saikia J, DB Phookan, NS Barua, P Das, L Saikia and J Saikia (2021) Morpho-biochemical characterization of brinjal (*Solanum melongena* L.) genotypes during kharif season. *Agric. Sci. Dig.* 41(3): 420-426.
- Sanni KA, I Fawole, RG Guei, DK Ojo and EA Somado (2008) Geographical patterns of phenotypic diversity in *Oryza sativa* landraces of Côte d'Ivoire. *Euphytica*, 160: 389-400.
- Savankumar N, Patel, C Raj, Popat, A Priya, Patel and RD Vekariya (2018) Genetic diversity analysis in brinjal (*Solanum melongena* L.) genotypes: A principal component analysis approach. *Int. J. Curr. Microbiol. App. Sci.* 7(1): 3296-3301.
- Sonagara MK, BN Patel, RR Acharya, A Parihar, R Patel and U Vaghela (2022) Assessment of genetic variability, heritability and genetic advance in brinjal (*Solanum melongena* L.). *Pharma innov.* 11(12): 1981-1983.
- Soumya BK, C Deo, S Singh, ND Bhutia, T Tayeng, BN Hazarika, A Debbarma, K Sakhamo and R Dutta (2023) Genetic variability studies in brinjal (*Solanum melongena* L.) for quantitative and qualitative characters. *Int. J. Environ. Clim. Change* 13 (10): 4463-4473.
- Statistical data agriculture (2023) <https://www.indiastat.com>
- Tyagi V, P Brahmi, SK Yadav, Pragya, SP Singh, S Singhand K Singh (2021) Germplasm access from ICAR-NBPGR and use within India. *Indian J. Plant Genet. Resour.* 34(2): 216-220.
- Uddin MS, M Billah, R Afroz, S Rahman, N Jahan and M Hossain (2021) Evaluation of 130 eggplant (*Solanum melongena* L.) genotypes for future breeding program based on qualitative and quantitative traits and various genetic parameters. *Horticulture* 7(376): 1-23.

RESEARCH ARTICLE

Preliminary Characterization of Wild *Actinidia Callosa* Var. *Strigillosa* as Potential Genetic Resources from North Sikkim, India

Abhilash Padhan^{1,2}, Dinesh Singh Thakur³, Vishal Singh Rana², RK Dogra², Narender Negi^{4*}, Akriti Chauhan³ and Pooja Sharma²

Abstract

A preliminary morpho-biochemical characterisation of fruit sampled from 81 wild kiwifruit (*Actinidia callosa* var. *strigillosa*) female vines distributed across the Lachung and Lachen territorial regions of North Sikkim district of Sikkim in Northeast India during 2021-2023 resulted in the identification of nine promising female accessions as potential genetic resources. Among these, SKNLC-18 and SKNLC-14 recorded the highest fruit weight, while SKNFF-05 and SKNLA-16 exhibited the highest vitamin C content, SKNLA-15 and SKNLC-16 resulted in the highest sugar: acid ratio, SKNZM-09 and SKNLC-10 produced the highest yield per vine, whereas SKNFF-07 exhibited greenish-yellow flesh colour. Sensory evaluation (9-point hedonic scale) of these nine superior accessions together revealed the highest overall acceptability in SKNLC-18 and SKNLA-16. The present preliminary assessment identified these nine accessions as promising plant genetic resources (PGRs) for breeding cold-hardy kiwifruit cultivars and expanding cultivation in the high-altitude Himalayan region.

Keywords: *Actinidia callosa* var. *strigillosa*, Cold-hardy, Genetic resources, North Sikkim, Vitamin C

¹Division of Genomic Resources, ICAR-National Bureau of Plant Genetic Resources, New Delhi, 110012, India.

²Department of Fruit Science, Dr YS Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh, 173230, India.

³Regional Horticultural Research and Training Station, Mashobra, Shimla, Himachal Pradesh, 171007, India.

⁴ICAR-National Bureau of Plant Genetic Resources, Regional Station, Phagli, Shimla, Himachal Pradesh, 171004, India.

***Author for correspondence:**

narender.nbpg@gmail.com

Received: 11/02/2026

Revised: 05/06/2026

Accepted: 09/06/2026

How to cite this article: Padhan A, DS Thakur, VS Rana, RK Dogra, N Negi, A Chauhan and P Sharma (2026) Preliminary Characterization of Wild *Actinidia callosa* var. *strigillosa* as Potential Genetic Resources from North Sikkim, India. *Indian J. Plant Genet. Resour.* 39(2): 83-99.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.176004>

Introduction

Kiwifruit (*Actinidia* spp.), a deciduous vine of the family *Actinidiaceae* and genus *Actinidia*, is predominantly cultivated in major kiwifruit-producing regions such as New Zealand, Chile, and China (Lee *et al.*, 2020). It is well known for its high vitamin C content and natural laxative properties, making it a nutritious and effective fruit for relieving constipation in people of all ages (Nishiyama *et al.*, 2004; Kim *et al.*, 2011). It is an excellent natural source of dietary folate (Vitamin B9), containing approximately 31-38 µg per 100 g of fresh fruit (Richardson *et al.*, 2018). Before the late 1990s, kiwifruit cultivation was largely focused on the New Zealand origin 'Hayward' cultivar (*A. deliciosa*), known for its green flesh and commercial reliability (Kwack *et al.*, 2010). The kiwifruit cultivars 'Hayward' and 'Allison' (*A. deliciosa*), introduced to India in the early 1960s, perform well between 300 and 2000 m above mean sea level (AMSL) in the Indian Himalayan region. However, their subtropical origin limits cultivation above 2000 m AMSL, where harsh winter conditions lead to leaf wilting, flower and fruit drop, delayed phenology, reduced fruit quality, and frequent vine mortality (Joshi, 1998; Jhalegar *et al.*, 2011). In addition, the high acidity of 'Hayward' and difficulties associated with uniform ripening under household conditions

have reduced its consumer acceptability in recent years (Wi *et al.*, 2019; Kwack *et al.*, 2022). Of the 56 recognized species within the genus *Actinidia*, 52 are native to China (Henare, 2016), while India harbours three species, viz. *A. callosa*, *A. strigosa* (Mao & Dash, 2020) and *A. indica*, a recent documentation from Arunachal Pradesh in Northeast India (Maity *et al.*, 2026). Among these, *A. callosa* is particularly noteworthy due to its strong frost tolerance and cold-hardy traits (Padhan *et al.*, 2023), making it a valuable genetic resource for the introgression of cold tolerance into *A. deliciosa* through interspecific hybridization (Arora, 2014). Recent explorations in the forested regions of North Sikkim, India, have for the first time in the country, documented the natural occurrence of *A. callosa* var. *strigillosa* C.F. Liang, with female wild kiwifruit selections thriving at elevations of 2441–2843 m AMSL, underscoring their value as cold-hardy genetic resources for expanding kiwifruit cultivation and breeding in high-altitude Himalayan environments (Padhan *et al.*, 2023; Padhan & Thakur, 2025a).

Since the early 20th century, kiwifruit breeding programmes have primarily focused on improving fruit weight, vitamin C content, sweetness, yield potential, and flesh colour. With the development of numerous new cultivars, consumer preferences, particularly among younger populations, have shifted toward hairless, sweeter fruits with diverse flesh colours (Latocha & Krupa, 2007). Nowadays, growers too are increasingly adopting fruit cultivars with intense skin and flesh colour to receive greater consumer preference and higher market value (Kumar *et al.*, 2026). The cultivars, viz. 'Tsechelidis', 'Hayward', and 'Jinyan' (JY) are among the kiwifruit varieties routinely selected for their consistently larger fruit size in commercial production and breeding initiatives (Zhong *et al.*, 2011). Considering vitamin C as a key selection criterion, *A. kolomikta* ranks as the species with the greatest ascorbic acid accumulation, while among widely cultivated germplasm, the *A. chinensis* cultivar 'Zespri Sun Gold' is distinguished for its comparatively high vitamin C concentration (Latocha, 2017). Evidence from multiple research reports demonstrates that yellow-fleshed and certain red-fleshed *A. chinensis* var. *chinensis* genotypes consistently show higher TSS and overall sugar accumulation compared with traditional green-fleshed *A. deliciosa* cultivars such as 'Hayward' (Zhang *et al.*, 2020). For enhanced yield potential, the principal parental resources comprise the commercially dominant cultivar 'Hayward', advanced high-yielding germplasm such as 'Zesy002' (Zespri Sun Gold), and diverse wild *Actinidia* taxa (Feng & Huang, 2025).

Traditionally, the green-fleshed cultivar *A. deliciosa* 'Hayward' has served as the predominant female parent in kiwifruit breeding programmes. In contrast, the development of yellow-fleshed cultivars has primarily utilised elite *A. chinensis* var. *chinensis* germplasms, viz. 'Hort-16A' (Zespri Gold), 'Gold3' (Zespri Sun Gold), 'Jintao' (Jin gold), 'Jinyan', and 'Jinshi 1' as key female progenitors. For the breeding of red-fleshed kiwifruit, cultivars including 'Hongyang', 'Donghong' (Oriental Red), and 'Zespri Ruby Red' have been widely adopted as principal parental lines due to their stable red pigmentation, high anthocyanin concentration, and favourable horticultural performance (Liu *et al.*, 2017). By strategically crossing these parental lines, breeders can introgress complementary traits, including yield potential, flavour quality, disease resistance, and distinct flesh-colour phenotypes to generate novel, commercially attractive kiwifruit cultivars (Hanley, 2018).

Under prolonged evolutionary pressures and natural selection, *Actinidia* species have developed extensive phenotypic and genetic diversity both within and among species, enabling them to adapt and survive across diverse environments. Leveraging this extensive genetic diversity, kiwifruit breeding programs worldwide have developed a wide range of improved cultivars (Lee *et al.*, 2020). The wild kiwifruit (*A. callosa* var. *strigillosa*) populations of North Sikkim exhibit substantial genetic diversity. Based on key fruit-quality attributes for table purposes, viz. TSS and Total sugars, six superior female accessions have been identified (Padhan & Thakur, 2025a), along with two promising male accessions selected for their superior pollinating traits, viz. higher average stamens count and longer flowering duration (Padhan & Thakur, 2025b).

Although the mid-hill regions (500–2000 m AMSL) of the Indian Himalayan states offer favourable agroclimatic conditions for the cultivation of kiwifruit (*A. deliciosa*), there remains a considerable scope for expanding its cultivation into higher altitude areas (2000–3000 m AMSL). Diversification in existing varieties by bringing wild and underutilized species into commercial cultivation may help in increasing the total kiwifruit production to meet the growing consumer demand and reduce imports. These naturally evolved populations possess several desirable traits, including variation in fruit size, flesh colour, sugar-acid balance, ascorbic acid content and may serve as potential rootstocks to improve cold and drought tolerance in commercial *Actinidia* spp. Despite their potential importance as genetic resources and rootstocks, limited efforts have been made to

systematically explore and characterize the superior wild kiwifruit accessions.

Therefore, the present investigation was undertaken to assess the nature and magnitude of morpho-biochemical variability among the wild kiwifruit populations growing in their natural habitats and to select and conserve superior accessions possessing horticulturally desirable traits, viz. fruit size, flesh colour, sugar-acid balance, ascorbic acid content and vine yield, which may be utilized as parental breeding lines for future crossing and hybridisation programmes.

Materials and Methods

Study location

The present study was carried out during 2021-2023 in the Lachung and Lachen valleys of the North Sikkim district of the Northeastern Himalayan state of Sikkim, India. According to the Koppen-Geiger climatic classification, the study area comprises a warm temperate climate (Cfb) in Lachen, gradually

transitioning to a sub-alpine climate (ET) in Lachung. A preliminary survey identified 81 female vines of wild kiwifruit (*Actinidia callosa* var. *strigillosa*) distributed across 15 locations. Of these, eight sub-locations in the Lachung region, viz. Katao Road (LA1), Sarchowk (LA2), Shanti Nagar (LA3), New Cutting Road (LA4), Lanka (LA5), Fakha Farey (FF), Bitchu (BC), and Teeling (TL) and seven sub-locations in the Lachen region, viz. Check Post (LC1), Hotel Panghen Metok (LC2), Puchi (LC3), Near Chaten (LC4), Zemu Chu Bridge (LC5), Chaten (CH), and Zeema (ZM), were included for morpho-biochemical evaluation (Fig. 1). Based on field observations and preliminary assessment, nine superior accessions were selected from five sub-locations, viz. Shanti Nagar (LA3) and Fakha Farey (FF) in the Lachung region, and Puchi (LC3), Zemu Chu Bridge (LC5), and Zeema (ZM) in the Lachen region (Table 1; Fig. 1). The geographical distribution of these superior accessions ranged from 27°41'41.244" N to 27°45'33.840" N latitude and 88°32'18.636" E to 88°45'00.972" E longitude, at elevations between 2632 and 2829 m AMSL (Table 1;




Fig. 1: Location details of selected wild-kiwifruit female accessions from various sub-locations under Lachen and Lachung territorial regions in North Sikkim district of Sikkim.

Out of 15 locations, 8 sub locations under Lachung area, i.e. LA1 – Katao Road, LA2 – Sarchok, LA3 – Shanti Nagar, LA4 – New Cutting Road, LA5 – Lanka, FF – Fakha Farey, BC – Bitchu, TL – Teeling, 7 sub locations under Lachen area, i.e. LC1 – Check Post, LC2 – Hotel Panghen Metok, LC3 – Puchi, LC4 – Near Chaten, LC5 – Zemu Chu Bridge, CH – Chaten, ZM – Zeema.

Fig. 1). Field experiments were approved by the Department of Forest, Environment and Wildlife Management, Government of Sikkim (F.No:78/GOS/FEWMD/BD-R-2015/191).

Plant material

At the beginning of 2021, a preliminary survey was conducted in the North Sikkim district to document wild kiwifruit accessions in their natural habitats. The survey led to the identification of 81 healthy and regular fruit-bearing female vines across 15 different sub-locations in the Lachung and Lachen valleys of North Sikkim (Supplementary Excel Sheet 1). Subsequently, during the following year, fruit samples were systematically collected from the selected vines during the peak fruiting period in September 2022 (Table 1). Scion wood from the nine promising female selections was collected from 5 different sub-locations under the North Sikkim district as described in Table 1, during the first week of March 2023 and tongue-grafted onto seedling rootstocks of kiwifruit (*A. deliciosa*) for field conservation and subsequent utilization.

Morphological observations

Twenty fruits were randomly sampled from each vine and analysed at the Centre for Agricultural Engineering and Post-Harvest Technology (CAEPHT), Gangtok,

Sikkim. Average fruit yield per vine was estimated by multiplying the mean fruit weight of 20 fruits collected from random branches by the total number of fruits per vine recorded during the field survey. The average value of 10 randomly selected fruits from each vine was taken into calculations for measuring fruit weight, fruit size and stalk length. Fruit weight was recorded using an electronic weighing balance (Model BT-2245), while fruit length, diameter, and stalk length were measured using Vernier Calipers (Model CD-6' CS, Mitutoyo Corp., Japan). Fruit size was calculated as the product of fruit length and diameter, and the length-to-diameter ratio was also determined.

Fruit's flesh colour was assessed visually following the DUS test guidelines of UPOV (2019) using Royal Horticultural Society (RHS), London colour charts. Fruit's skin and flesh colour traits, including colour of the outer pericarp, locule, and core, were recorded in accordance with the DUS guidelines for *Actinidia* (Kiwifruit) as prescribed by UPOV (2019) and compared with the RHS colour charts. Sensory evaluation of only selected superior nine accessions was conducted by a panel of four judges, who assessed fruit samples for flavour, texture, aroma, taste, and overall acceptability using a nine-point hedonic scale (Lawless & Heymann, 2010). Harvest maturity was determined by monitoring TSS levels, with fruits harvested upon reaching the

Table 1: Location details and morphological non-metric characters of selected wild- kiwifruit female accessions from North Sikkim district of Sikkim

Sl.	Accessions	Sub locations & Major Locations under North Sikkim District	Denoted in the Map (refer Figure 1)	Latitude	Longitude	Altitude (metres above mean sea level)	Estimated Age	Fruit shape	Conspicuousness of lenticels on fruit's skin	Hairiness of fruit's skin	Fruit skin colour	Colour of outer pericarp	Fruit's flesh colour Colour of locule	Colour of core	Width of core relative to fruit	General shape of core in cross section	Date of full bloom	Anticipated date of harvest	Harvesting period (days)
1	SKNLC-14	Puchi, Lachen	LC3	27° 44' 44.448" N	88° 32' 40.812" E	2710	35	Oblong	Medium	Absent	Medium green	Dark green	Dark green	White	Small	Circular	27-06-2022	12-09-2022	78
2	SKNLC-18	Zemu Chu Bridge, Lachen	LC5	27° 45' 20.268" N	88° 32' 18.636" E	2734	43	Obovate	Medium	Absent	Light green	Dark green	Light green	White	Medium	Oblate	25-06-2022	28-09-2022	96
3	SKNLA-16	Shanti Nagar, Lachung	LA3	27° 41' 41.244" N	88° 45' 0.972" E	2638	48	Obovate	Strong	Absent	Medium green	Medium green	Light green	White	Medium	Transverse elliptic	25-06-2022	15-09-2022	83
4	SKNFF-05	Faka Farey, Lachung	FF	27° 42' 45.864" N	88° 44' 59.208" E	2807	29	Oblong	Medium	Absent	Light green	Medium green	Light green	White	Small	Oblate	03-07-2022	20-09-2022	80
5	SKNLA-15	Shanti Nagar, Lachung	LA3	27° 41' 41.424" N	88° 45' 0" E	2632	32	Oblong	Medium	Absent	Medium green	Medium green	Medium green	White	Medium	Oblate	30-06-2022	14-09-2022	77
6	SKNLC-16	Puchi, Lachen	LC3	27° 44' 44.808" N	88° 32' 40.488" E	2713	63	Obovate	Medium	Absent	Light green	Medium green	Light green	White	Small to medium	Circular	27-06-2022	10-09-2022	76
7	SKNZM-09	Zeema, Lachen	ZM	27° 45' 33.84" N	88° 32' 19.5" E	2781	58	Oblong	Strong	Absent	Medium green	Dark green	Medium green	White	Medium	Oblate	05-07-2022	26-09-2022	84
8	SKNLC-10	Puchi, Lachen	LC3	27° 44' 49.56" N	88° 32' 39.048" E	2740	41	Oblong	Medium	Absent	Light green	Dark green	Dark green	White	Small to medium	Transverse elliptic	04-07-2022	13-09-2022	72
9	SKNFF-07	Fakha Farey, Lachung	FF	27° 42' 48.78" N	88° 44' 56.94" E	2829	17	Oblong	Medium	Absent	Medium green	Greenish yellow	Greenish yellow	Yellow white	Medium	Transverse elliptic	08-07-2022	24-09-2022	89

[i.e. Comparison with RHS Colour Charts :Colour of outer pericarp (Dark green: Yellow Green group 144 A, Medium green: Yellow Green group 144 B, Greenish yellow: Yellow Green group 154 B), Colour of locule (Greenish Yellow: Yellow Green group 154 B, Dark green: Green Group 143 A, Medium Green: Green Group 143 B, Light Green: Green Group 143 C), Colour of core (White: Green-White group 157 D, Yellow white: Yellow- White group 158 A), Fruit skin colour: (Light green: Yellow Green group 151 A); (Medium green: Yellow Green group 152 B)]

optimum stage (TSS >6.5 °Brix) as suggested by Beever and Hopkirk (1990). The harvesting period for each selected accession was calculated as the duration from full bloom to the anticipated harvest date.

Biochemical observations

TSS: Total soluble solids (TSS) were measured using a digital Erma hand refractometer (0-32 °Brix). The instrument was calibrated with distilled water before analysis. A few drops of fruit juice were placed on the prism surface, and the readings were recorded after applying temperature correction whenever the temperature deviated from 20°C (A.O.A.C., 1970). The TSS content was expressed as °Brix. For each accession, observations were recorded from five randomly collected fruits per vine, and the mean value was considered for analysis.

Titrateable acidity (%): The titrateable acidity (%) was obtained by using the procedure given by Ranganna (2009). For the estimation of titrateable acidity, 25 g of fruit pulp from each sample was crushed, and the volume was made up to 250 ml with distilled water in a volumetric flask. From this extract, a 50 ml aliquot was used for acidity estimation, while the remaining extract was utilized for the determination of total and reducing sugars. An aliquot of 10 ml juice extract was titrated against N/10 NaOH solution using phenolphthalein as an indicator until the appearance of a faint pink endpoint. The titrateable acidity was expressed as per cent malic acid, considering that 1 ml of N/10 NaOH is equivalent to 0.0067 g of anhydrous malic acid. The acidity percentage was calculated using the following formula:

$$\text{Titrateable acidity (\%)} = \frac{\text{Titre value} \times \text{Normality of Alkali} \times \text{Volume-madeup} \times \text{Equivalent weight of acid} \times 100}{\text{Volume of sample taken for estimation} \times \text{Volume of aliquot taken} \times 1000}$$

Total sugars (%): Total sugars were estimated by the Lane and Eynon volumetric method (A.O.A.C., 1970). The remaining 200 ml of juice extract obtained after acidity estimation was clarified by adding 10 ml of saturated lead acetate solution. The contents were shaken thoroughly and filtered into a flask containing 10 ml of potassium oxalate solution to remove excess lead. After shaking and allowing the precipitate to settle, the solution was filtered again. An aliquot of 50 ml filtrate was transferred to a 250 ml volumetric flask, mixed with 5 ml concentrated HCl, and kept overnight at room temperature for hydrolysis of non-reducing

sugars. The excess acid was neutralized using a saturated NaOH solution. The hydrolysed extract was then titrated against a boiling mixture of 5 ml each of Fehling's solution A and B using methylene blue as an indicator. The endpoint was marked by the disappearance of blue colour and the formation of a brick-red precipitate. The total sugar content was expressed as a percentage on a fresh pulp weight basis as:

$$\text{Total sugars (\%)} = \frac{\text{Factor} \times \text{Dilution}}{\text{Titre value} \times \text{Weight or volume of sample taken}} \times 100$$

*Factor=0.05

Reducing sugars (%): The reducing sugars (%) were obtained by using the procedure given by Ranganna (2009). For the estimation of reducing sugars, 5 ml each of Fehling's solution A and Fehling's solution B were mixed with 50 ml of distilled water in a beaker and boiled for about 10 minutes. Subsequently, 2-3 drops of methylene blue indicator were added, and the solution was titrated against the prepared filtrate. The contents were continuously shaken during titration, and the endpoint was indicated by the appearance of a brick-red precipitate. The reducing sugar content was expressed as a percentage on a fresh pulp weight basis and calculated using the following formula:

$$\text{Reducing sugars (\%)} = \frac{\text{Factor} \times \text{Dilution}}{\text{Titre value} \times \text{Weight or volume of sample}} \times 100$$

*Factor=0.05

Non-reducing sugars (%): The non-reducing sugars (%) were obtained by using the procedure given by Ranganna (2009). The non-reducing sugar content was calculated by subtracting the reducing sugars from the total sugars and multiplying the difference by the standard factor 0.95. The non-reducing sugars were expressed as a percentage on a fresh pulp weight basis using the following formula:

$$\text{Non-reducing sugars} = (\text{Total sugars} - \text{Reducing sugars}) \times 0.95$$

Sugar/acid ratio: It was calculated by dividing the percentage of total sugars by the percentage of titrateable acidity.

$$\text{Sugar/acid ratio} = \frac{\text{Total sugars (\%)}}{\text{Acidity (\%)}}$$

Total phenols (mg GAE/100 g): Total phenol content was determined by the Folin-Ciocalteu method described by Singleton & Rossi (1965), and absorbance was recorded at 650 nm using a spectrophotometer. In alkaline medium, phenolic compounds react with phosphomolybdic acid present in the Folin-Ciocalteu reagent to form a dark blue coloured complex (molybdenum blue). The intensity of the developed colour was measured at 650 nm. A standard calibration curve was prepared using different concentrations of gallic acid, and the total phenol content of the samples was estimated from the standard curve and expressed as mg/g of sample.

Ascorbic acid (mg/100 g): Quantitative estimation of ascorbic acid was carried out following the A.O.A.C. (1980) method using 2,6-dichlorophenol indophenol dye. Five grams of the fruit sample were homogenised in 3% metaphosphoric acid (extraction solution), and the volume was made up to 50 ml. An aliquot of the extract was then titrated against the dye solution until the appearance of a light pink colour endpoint. The ascorbic acid content was calculated using the following formula:

$$\text{Ascorbic acid (mg/100 g)} = \frac{\text{Dye factor} \times \text{Titration reading}}{\text{Weight of fruit taken}} \times \frac{\text{Volume made}}{\text{Volume taken for estimation}} \times 100$$

Statistical analysis

The data generated from the evaluation of nine promising wild kiwifruit accessions for various morphological, biochemical, and sensory traits were subjected to statistical analysis using Analysis of Variance (ANOVA) in Microsoft Excel. The significance of differences among accessions was tested using the F-test, and treatment means were compared using the Critical Difference (CD) test at the 5% level of significance. Standard Error of Mean (SEm ($\pm$)), coefficient of variation (CV%), overall mean, and range were computed for each morphological and

biochemical metric character to assess the extent of variability among the accessions. The statistical parameters, including mean, range, SEm ($\pm$), CD (0.05), CV (%), and F-test significance levels, are presented for all evaluated traits to facilitate comparison of accession performance and identification of superior accessions.

Results and Discussion

In 2022, fruit samples obtained from 81 female wild kiwifruit vines were subjected to detailed morphological and biochemical evaluations (Supplementary Excel Sheet 1). Based on overall performance and their potential value as genetic resources, nine accessions, viz. SKNLC-14, SKNLC-18, SKNLA-16, SKNFF-05, SKNLA-15, SKNLC-16, SKNZM-09, SKNLC-10, and SKNFF-07, were identified as superior. These accessions exhibited superior performance for key economically and nutritionally important traits, including higher fruit weight, elevated ascorbic acid content, superior sugar-acid ratio, greater yield per vine, and a desirable greenish-yellow flesh colour. The morphological characteristics of these selected accessions are illustrated in Tables 1-4 & Fig. 1-10. The selection criteria emphasized traits critical for genetic improvement and utilization in breeding programmes. Consequently, these nine accessions represent valuable genetic resources with significant potential for incorporation into kiwifruit breeding efforts aimed at developing improved cultivars that meet market requirements and consumer preferences.

Morphological and biochemical metric characters as selection criteria

Fruit weight: Among the 81 wild kiwifruit accessions, SKNLC-18 exhibited the highest mean fruit weight (9.16 g), followed by SKNLC-14 (8.07 g), whereas the fruit weight ranged between 2.94 and 9.16 g among the selected 9 accessions (Table 2). These values exceed or are comparable to those reported in earlier studies,

Table 2: Morphological metric characters of selected wild kiwifruit female accessions from North Sikkim district of Sikkim

Accessions	Fruit weight (g)	Fruit length (mm)	Fruit diameter (mm)	Fruit size (Length *Diameter) (mm ²)	Fruit (length to diameter ratio)	Length of fruit stalk (mm)	Yield (kg/vine)
SKNLC-14	8.07	23.26	19.94	463.80	1.17	15.93	9.00
SKNLC-18	9.16	32.21	22.42	722.15	1.44	17.33	9.00
SKNLA-16	5.18	25.34	19.04	482.47	1.34	16.32	4.50
SKNFF-05	2.98	23.62	19.11	451.38	1.24	17.14	18.50
SKNLA-15	4.62	22.85	19.26	440.09	1.20	17.85	5.50

SKNLC-16	4.14	23.38	16.66	389.51	1.40	16.43	10.50
SKNZM-09	4.25	18.86	17.14	323.26	1.10	11.94	28.50
SKNLC-10	3.83	24.82	16.08	399.11	1.56	11.18	26.50
SKNFF-07	2.94	18.34	16.32	299.31	1.13	15.14	2.50
Range	2.94-9.16	18.34-32.21	16.08-22.42	299.31-722.15	1.10-1.56	11.18-17.85	2.50-28.50
Mean	5.02	23.63	18.44	441.23	1.29	15.47	12.72
SEm ($\pm$)	0.09	0.44	0.26	8.88	0.02	0.27	0.13
CD (P = 0.05)	0.27	1.30	0.77	26.63	0.05	0.80	0.39
CV (%)	3.13	3.19	2.41	3.49	2.28	3.00	1.78
F -test	**	**	**	**	**	**	**

I.e.: SEm ($\pm$) = Standard Error of Mean; CD = Critical Difference; CV = Coefficient of Variation; ** = Significant at $P \leq 0.01$

where average fruit weights of 6.9 g were recorded in the hardy kiwifruit cultivar 'Ananasnaya' (*Actinidia arguta*) grown in Oregon, USA (Pescie & Strik, 2004), 8.1 g average weight in Bingo, a F_1 hybrid between *A. purpurea* 'Purpurna Sadova' and *A. arguta* (Latocha, 2012), 4.8 g in *Actinidia callosa* var. *strigillosa* from China

(Hongwen, 2014), 6.71 g in *A. melanandra* (4x), and 4.85 g in *A. arguta* var. *arguta* (4x) from Northwest China (Zhang *et al.*, 2017). The comparatively higher fruit weight observed in SKNLC-18 (Fig 3a, b, c) and SKNLC-14 (Fig 2a, b, c) underscores substantial intra-specific variability within wild germplasm and highlights the



SKNLC-14

Fig. 2 (SKNLC-14): Depicting wild kiwifruit (*A. callosa* var. *strigillosa*). (2a) fruit's flesh with outer pericarp, locule and inner core; (2b) length and diameter of fruit with measuring scale; (2c) vine during the fruiting period in North Sikkim, India.



SKNLC-18

Fig. 3 (SKNLC-18): Depicting wild kiwifruit (*A. callosa* var. *strigillosa*). (3a) fruit's flesh with outer pericarp, locule and inner core; (3b) length and diameter of fruit with measuring scale; (3c) vine during the fruiting period in North Sikkim, India.

potential of these accessions as valuable genetic resources for breeding programmes aimed at fruit size improvement.

Vitamin C level: The accessions SKNFF-05 (106.60 mg/100g) and SKNLA-16 (91.00 mg/100 g) exhibited the highest vitamin C among the 81 wild kiwifruit female accessions, with a range of 28.60-106.60 mg/100 g Vit. C among the selected 9 accessions (Table 3), reflecting substantial variability for ascorbic acid accumulation in Himalayan *Actinidia* resources. These values are comparable to those reported for the green kiwifruit cultivar 'Hayward' (80-120 mg/100 g fresh weight (Beever & Hopkirk, 1990), though lower than gold

kiwifruit cultivars such as *A. chinensis* cv. 'Zespri Sun Gold', which contains approximately 161.3 mg/100 g (USDA, 2016; Richardson *et al.*, 2018). Wild and semi-domesticated *Actinidia* species from the Himalayan region are recognised for their superior nutraceutical traits, with *A. kolomikta* reported to possess the highest ascorbic acid concentration among the genus (Latocha, 2017). The comparatively high vitamin C levels observed in SKNFF-05 (Fig 5a, b, c) and SKNLA-16 (Fig 4a, b, c) highlight its potential as a promising donor accession for biofortification-oriented breeding programmes and for broadening the nutritional base of cultivated kiwifruit under temperate Himalayan conditions.

Table 3: Biochemical characters of wild kiwifruit female accessions from North Sikkim district of Sikkim

Accessions	TSS (°B)	Titrateable acidity (%)	Total sugars (%)	Reducing sugars (%)	Non-reducing sugars (%)	Sugar: acid ratio	Vit. C (mg/100 g)	Total Phenols (mg GAE/100g)
SKNLC-14	16.40	0.77	8.57	6.67	1.80	11.13	54.60	39.30
SKNLC-18	15.46	1.05	7.55	5.55	1.90	7.19	46.80	57.15
SKNLA-16	16.76	0.77	8.33	5.88	2.32	10.82	91.00	86.11
SKNFF-05	14.40	0.56	6.81	5.55	1.19	12.16	106.60	75.94
SKNLA-15	15.36	0.49	7.22	5.68	1.46	14.73	65.00	30.92
SKNLC-16	13.80	0.49	6.81	5.26	1.47	13.90	33.80	34.67
SKNZM-09	12.35	0.56	5.26	3.90	1.29	9.39	28.60	71.17
SKNLC-10	12.70	0.70	5.35	4.16	1.13	7.64	33.80	81.20
SKNFF-07	15.60	0.6	37.50	5.88	1.53	11.90	36.40	27.60
Range	12.35-16.76	0.49-1.05	5.26-8.57	3.90-6.67	1.13-2.32	7.19-14.73	28.60-106.60	27.60-86.11
Mean	14.76	0.67	7.04	5.39	1.56	10.98	55.18	56.01
SEm (±)	0.26	0.01	0.12	0.08	0.13	0.15	0.95	1.22
CD (P = 0.05)	0.79	0.03	0.36	0.23	0.02	0.44	2.84	3.67
CV (%)	3.08	2.59	2.98	2.49	2.54	2.31	2.97	3.79
F-test	**	**	**	**	**	**	**	**

I.e.: SEm (±) = Standard Error of Mean; CD = Critical Difference; CV = Coefficient of Variation; ** = Significant at $P \leq 0.01$



SKNLA-16

Fig. 4 (SKNLA-16): Depicting wild kiwifruit (*A. callosa* var. *strigillosa*). (4a) fruit's flesh with outer pericarp, locule and inner core; (4b) length and diameter of fruit with measuring scale; (4c) vine during the fruiting period in North Sikkim, India.



SKNFF-05

Fig. 5 (SKNFF-05): Depicting wild kiwifruit (*A. callosa* var. *strigillosa*). (5a) fruit's flesh with outer pericarp, locule and inner core; (5b) length and diameter of fruit with measuring scale; (5c) vine during the fruiting period in North Sikkim, India.



SKNLA-15

Fig. 6 (SKNLA-15): Depicting wild kiwifruit (*A. callosa* var. *strigillosa*). (6a) fruit's flesh with outer pericarp, locule and inner core; (6b) length and diameter of fruit with measuring scale; (6c) vine during the fruiting period in North Sikkim, India.



SKNLC-16

Fig. 7 (SKNLC-16): Depicting wild kiwifruit (*A. callosa* var. *strigillosa*). (7a) fruit's flesh with outer pericarp, locule and inner core; (7b) length and diameter of fruit with measuring scale; (7c) vine during the fruiting period in North Sikkim, India.

Highest sugar: acid ratio: The accessions SKNLA-15 (14.73) and SKNLC-16 (13.90) recorded the highest sugar: acid ratio among the 81 accessions, whereas it ranged between 7.19-14.73 among the selected 9

accessions under present investigation (Table 3), indicating superior sweetness perception and overall fruit palatability. Earlier studies have reported comparatively lower sugar-acid ratios across different

**SKNZM-09**

Fig. 8 (SKNZM-09): Depicting wild kiwifruit (*A. callosa* var. *strigillosa*). (8a) fruit's flesh with outer pericarp, locule and inner core; (8b) length and diameter of fruit with measuring scale; (8c) vine during the fruiting period in North Sikkim, India.

**SKNLC-10**

Fig. 9 (SKNLC-10): Depicting wild kiwifruit (*A. callosa* var. *strigillosa*). (9a) fruit's flesh with outer pericarp, locule and inner core; (9b) length and diameter of fruit with measuring scale; (9c) vine during the fruiting period in North Sikkim, India.

**SKNFF-07**

Fig. 10 (SKNFF-07): Depicting wild kiwifruit (*A. callosa* var. *strigillosa*). (10a) fruit's flesh (Greenish yellow) with outer pericarp, locule and inner core; (10b) length and diameter of fruit with measuring scale; (10c) vine during the fruiting period in North Sikkim, India.

Actinidia species. Wang *et al.*, (2018) observed ratios ranging from 5.33 to 8.05 in *A. kolomikta*, *A. arguta*, and *A. chinensis* cultivars, with Hayward (*A. chinensis* var. *deliciosa*) exhibiting a ratio of 6.50. Similarly, Liang *et al.*, (2021) reported sugar: acid ratios between 5.91 and 10.39 in 15 cultivars of *A. chinensis* and *A. chinensis* var.

deliciosa, with only a few cultivars (Hongyang, Yate, Jinyan, Cuiyu) exceeding a ratio of 10.0. These values fall at the upper end of the range (5.39-14.17) reported by Wen *et al.*, (2023) for hardy kiwifruit (*A. arguta*) cultivars at the soft-ripe stage, suggesting that the present accessions, viz. SKNLA-15 (Fig. 6a, b, c) and SKNLC-16

(Fig. 7a, b, c) possess a favourable biochemical balance comparable to advanced commercial cultivars, indicating their potential as suitable donor parents for enhancing fruit sweetness in kiwifruit breeding programmes in the Himalayan region.

Fruit yield: Among the 81 evaluated accessions, SKNZM-09 (28.50 kg/vine) and SKNLC-10 (26.50 kg/vine) recorded the highest yield per vine, with a range of 2.50-28.50 kg/vine fruit yield among the 9 selected accessions (Table 2), demonstrating outstanding yield potential. These values are substantially higher than those reported for several hardy kiwifruit cultivars, where average yields typically range from 4.2 to 6.2 kg /vine, as documented for 'HORTGEM TORU' and 'HORTGEM WHA' in New Zealand (McNeillage *et al.*, 2003a, b, c) and 'Vip Green' and 'Vip Red' in Romania (Stanica *et al.*, 2022). While higher cumulative yields have been reported in certain European cultivars under long-term cultivation (Latocha *et al.*, 2023), the superior per-vine productivity observed in SKNZM-09 (Fig. 8a, b, c) and SKNLC-10 (Fig. 9a, b, c) underscores the wider yield variability inherent in wild hardy kiwifruit germplasm. These accessions therefore represent promising elite selections for yield improvement and domestication-oriented breeding programmes in Himalayan regions.

Morphological non-metric character as a selection criterion

Greenish yellow flesh: The single accessions 'SKNFF-07' (Fig. 10a, b, c) exhibited a distinct greenish-yellow flesh colour, a trait of significant commercial and breeding relevance in kiwifruit among the evaluated 81 accessions. In this single accession, the outer pericarp and locular tissues were characterized by a greenish-yellow hue (Yellow-Green Group 154B), while the core region displayed a yellow-white colouration (Yellow-White Group 158A) (Table 1). Yellow-fleshed kiwifruit gained global prominence following the release of *Actinidia chinensis* 'Hort-16A', the first cultivar developed through a systematic breeding programme to be successfully commercialized internationally (Asadi *et al.*, 2024). Since then, substantial progress has been made worldwide, with yellow-fleshed cultivars developed both through direct selection from wild germplasm in China and through structured breeding programmes in other kiwifruit-growing regions. Notable examples include 'Dori' and 'Soreli' from Italy (Testolin & Cipriani, 2008; Costa *et al.*, 2014), 'Goldrush', 'Haegeum', 'Halla Gold', and 'Jecy Gold' from Korea (Kim *et al.*, 2007; Cho *et al.*, 2011; Kwack *et al.*, 2011; Kim *et al.*,

2012), 'Sanuki Gold' from Japan (Fukuda *et al.*, 2007), and 'Y368' from New Zealand (US Plant Patent, 2010). The occurrence of greenish-yellow flesh in the present accession highlights the potential of indigenous germplasm as a valuable genetic resource for developing novel yellow-fleshed kiwifruit cultivars adapted to Himalayan agro-climatic conditions, thereby reducing dependence on imported cultivars and broadening the genetic base of kiwifruit improvement programmes.

Other morphological and biochemical metric characters of selected accessions

In the present study, significant variation was observed among the nine accessions with respect to fruit size and biochemical attributes. Fruit length ranged from 18.34 mm in SKNFF-07 (Fig. 10b) to 32.21 mm in SKNLC-18 (Fig. 3b), while fruit diameter varied from 16.08 mm in SKNLC-10 (Fig. 9 b) to 19.94 mm in SKNLC-14 (Fig. 2b) (Table 2). The superior fruit length recorded in SKNLC-18 exceeded the values reported by Zhang *et al.*, (2017) for kiwi berry species in Northwest China, where fruit length ranged from 27.44 mm in *A. arguta* var. *arguta* (4x) to 29.49 mm in *A. arguta* var. *girdalii* (8x and 10x). Similarly, fruit diameter in the present study was comparable to that reported for *A. melanandra* (4x), which ranged from 19.66 to 21.36 mm, but higher than values reported for *A. arguta* var. *girdalii*.

Total soluble solids (TSS) content among the accessions varied from 12.35 °Brix in SKNZM-09 to 16.76 °Brix in SKNLA-16, with a mean value of 14.76 °Brix in the present study (Table 3). The higher TSS observed in SKNLA-16 (16.76 °B) was comparable to the post-ripening TSS values reported by Jung *et al.*, (2016) in *A. arguta* cultivars 'Mansu' (16.5 °B) and 'Chiak' (16.9 °B), indicating favourable sweetness levels. In contrast, TSS values reported at harvest by Jung *et al.*, (2016) and by Kishor *et al.*, (2017) in *A. chinensis* var. *deliciosa* cultivars were notably lower, suggesting species- and genotype-specific differences in sugar accumulation.

Total sugar content varied significantly among the accessions, ranging from 5.26 % in SKNZM-09 to 8.57% in SKNLC-14 (Table 3). The higher sugar content observed in SKNLC-14 (8.5%) was comparable to that reported by Kishor *et al.*, (2017), who recorded maximum total sugars of 8.31% in cv. Monty, followed by Allison (6.88%) and other commercial cultivars. These findings highlight the promising biochemical attributes of certain wild kiwifruit accessions, particularly SKNLC-14 and SKNLA-16, which exhibited fruit quality traits comparable to or exceeding those of cultivated kiwifruit (*A. chinensis*) varieties.

Other morphological non-metric characters of selected accessions

Considerable variation was observed in non-metric fruit characters among the selected accessions. Fruit shape ranged from oblong in six accessions (SKNLC-14 (Fig. 2b), SKNFF-05 (Fig. 5b), SKNLA-15 (Fig. 6b), SKNZM-09 (Fig. 8b), SKNLC-10 (Fig. 9b), and SKNFF-07 (Fig. 10b) to obovate in three accessions (SKNLC-18 (Fig. 3b), SKNLA-16 (Fig. 4b) and SKNLC-16 (Fig. 7b)), indicating substantial morphological diversity in the present study (Table 1). Similar cultivar-dependent variation in fruit shape has been reported earlier, with ovate fruits in hardy kiwifruit cvs. Mansu and Chiak (*A. arguta*) (Jung *et al.*, 2016) and oblong, elliptical and ovoid fruits in *A. chinensis* cvs. Goldone, Hayward, Goldrush and Hort16A (Kwack *et al.*, 2017). The conspicuousness of lenticels on the fruit skin varied from medium in seven accessions to strong in two accessions viz. SKNLA-16 (Fig. 4b) and SKNZM-09 (Fig. 8b) (Table 1). In contrast, Rodriguez (2019) reported a high density of conspicuous lenticels with the absence of skin hairs in cv. CRISTAL (*A. chinensis* var. *deliciosa*), suggesting genotype-specific expression of this trait. Fruit skin colour also differed among accessions, ranging from light green (Yellow Green group 151 A) in four accessions to medium green (Yellow Green group 152 B) in five accessions in the present investigation (Table 1), whereas a greenish-brown skin colour was reported in cv. Hongyang (*A. chinensis*) under China agroclimatic conditions

(Mingzhang & Mingzhong, 2008).

The harvesting period among the nine accessions ranged from 72 to 96 days, reflecting differences in maturity behaviour (Table 1). Comparable variation in harvest maturity has been documented in hardy kiwifruit, with harvesting at 81-84 days after flowering in cv. Issai (Lei *et al.*, 2022) and 100-110 days after flowering in cv. Ananasnaya (Strik & Hummer, 2006). Overall, the observed variability in non-metric traits underscores the genetic diversity of the evaluated germplasm and its potential relevance for selection and breeding programmes.

Sensory analysis

Consumer acceptance and purchase decisions for new kiwifruit cultivars are strongly influenced by the fruit's aroma and flavour. Cultivars that fail to meet consumer sensory expectations are unlikely to achieve sustained market success (Latocha & Krupa, 2007). Sensory evaluation of fruit samples from selected wild kiwifruit female accessions was carried out using a nine-point hedonic rating scale (Table 4). The attributes assessed included flavour, texture, aroma, taste, and overall acceptability. Scores were assigned to identify the most consumer-preferred accessions based on sensory perception. Hedonic scales have a well-established and robust history in consumer research and are widely used to effectively assess consumer preferences and satisfaction (Stone & Sidel, 2004). The

Table 4: Sensory evaluation of selected wild- kiwifruit female accessions from North Sikkim district of Sikkim

Genotypes	Flavour	Texture	Aroma	Taste	Overall acceptability
SKNLC-14	7.00	7.00	6.50	7.00	6.88
SKNLC-18	8.00	9.00	8.00	8.00	8.25
SKNLA-16	8.00	8.00	7.50	8.00	7.88
SKNFF-05	7.00	6.50	7.00	6.50	6.75
SKNLA-15	6.00	5.00	5.00	7.00	5.75
SKNLC-16	6.00	6.00	7.00	6.00	6.25
SKNZM-09	7.00	7.00	6.00	7.00	6.75
SKNLC-10	5.00	5.00	6.00	6.00	5.50
SKNFF-07	5.00	6.50	5.00	4.00	5.13
Range	5-8	5-9	5-8	4-8	5.13-8.25
Mean	6.55	6.67	6.44	6.61	6.57
SEm ($\pm$)	0.11	0.15	0.11	0.10	0.07
CD (P = 0.05)	0.34	0.44	0.31	0.31	0.22
CV (%)	2.96	3.83	2.82	2.69	1.95
F -test	**	**	**	**	**

I.e.: SEm ($\pm$) = Standard Error of Mean; CD = Critical Difference; CV = Coefficient of Variation; ** = Significant at $P \leq 0.01$

nine-point hedonic scale, with ratings ranging from 1 to 9, is widely employed to capture consumer preferences. The scale assumes that participants' sensory responses lie along a continuum of liking and disliking, where 1 represents 'dislike extremely', 2 'dislike very much', 3 'dislike moderately', 4 'dislike slightly', 5 'neither like nor dislike', 6 'like slightly', 7 'like moderately', 8 'like very much', and 9 'like extremely'. These ratings collectively reflect graded variations in consumer liking or disliking (Lawless & Heymann, 2010).

The flavour was rated excellent in SKNLC-18 and SKNLA-16, both receiving an 8-point hedonic scale, while the remaining selections were assessed as good. Pulp texture was outstanding in SKNLC-18 (9), followed by SKNLA-16 (8), whereas the other accessions exhibited acceptable to fair texture. Aroma was exceptional in SKNLC-18 (8) and SKNLA-16 (7.5); the remaining selections were generally rated as fair, except SKNFF-07, which recorded a comparatively low score (5.0). Taste was rated excellent in SKNLC-18 and SKNLA-16 (8), while SKNFF-07 was evaluated as poor. Overall, based on overall acceptability, SKNLC-18 and SKNLA-16 clearly outperformed all other accessions (Table 4).

Sensory attributes such as flavour, texture, aroma, and taste are key determinants of consumer acceptability in wild kiwifruit, with overall flavour perception representing an integrated response of taste sensations and aroma-active volatiles. Considerable variation in sensory quality has been reported among *Actinidia* species. Hongwen (2014) described *A. callosa* var. *strigillosa* as slightly acidic, astringent, and tart, with low juiciness and a mildly crispy texture, indicating limited sensory appeal. In contrast, Stanica *et al.*, (2022) reported a fine, distinctive, and predominantly sweet flavour in *A. arguta* cultivars 'Vip Green' and 'Vip Red'. In the present study, SKNLC-18 and SKNLA-16 exhibited superior sensory profiles, characterised by high scores for flavour, texture, aroma, and taste (Table 4), closely resembling the favourable attributes reported for sweet-tasting *A. arguta* cultivars. These results underscore the potential of SKNLC-18 and SKNLA-16 as promising accessions for consumer acceptance and gene source for future varietal development.

Conclusions and future prospective

Among the 81 evaluated wild kiwifruit vines, nine elite accessions based on superior fruit weight, vitamin C content, sugar: acid ratio, yield potential, and yellow flesh colour were identified, of which SKNLC-18 and SKNLA-16 emerged as outstanding accessions with superior fruit size, while SKNFF-07 was distinguished by

its greenish-yellow flesh colour. The occurrence of greenish-yellow flesh in wild Himalayan kiwifruit is of particular significance, as it represents a rare and valuable trait with high commercial and breeding relevance. The present study confirms a significant level of variation among the nine superior accessions that harbour immense untapped genetic potential and constitute an important reservoir of traits, especially for cold as well as drought tolerance, yield improvement, nutritional quality and consumer-preferred fruit attributes. These accessions also hold potential as parental lines for breeding and development of climate-resilient cold-hardy cultivars suited for the high-altitude areas of the Himalayan region.

These nine superior accessions have been established in the field gene bank of the researcher's institute. Their stability, adaptability, and horticultural performance will be validated through multi-location evaluations across high-altitude environments (2000–3000 m AMSL) in the Indian Himalayan region under homogenous conditions. In addition, comparative transcriptomic analyses involving cultivated kiwifruit (*Actinidia* spp.) groups will be undertaken to identify cold-responsive genes and gain insights into the genetic basis of cold tolerance, thereby facilitating their potential utilization in kiwifruit breeding programmes and expanding cultivation into the higher altitude regions of the Himalayan Mountains.

Acknowledgments

The authors are grateful to the Forest Department, Government of Sikkim, India, for allotting permission to carry out the exploration; to Dr. Rajiv Gogoi, Scientist-E/HOO, Botanical Survey of India, Gangtok, for providing preliminary information to carry out the exploration and to the Centre for Agricultural Engineering and Post Harvest Technology (CAEPHT), Gangtok, Sikkim, for providing lab facilities to conduct biochemical experiments.

References

- AOAC (1970) Official methods of analysis, Benjamin Franklin Statistics, Washington, D.C.
- AOAC (1980) Official methods of analysis of the Association of Official Analytical Chemists. A.O.A.C. Benjamin Franklin Station, Washington, D.C.
- Arora RK (2014) Diversity in underutilized plant species; an Asia Pacific perspective. Biodiversity International, New Delhi, India, 203 p.
- Asadi M, M Ghasemnezhad, A Bakhshipour, J Olfati and A Atak (2024) Breeding of new kiwifruit (*Actinidia chinensis*) cultivars with yellow (golden) fleshed and superior characteristics. *BMC Plant Biol.* 24: 1045.

- Beever DJ and G Hopkirk (1990) Fruit development and fruit physiology. In: Warrington IJ, Weston GC (eds) *Kiwifruit: Science and Management*. Auckland: Ray Richards, pp 97-126.
- Cho YS, HS Cho and MY Park (2011) A selection of Korean Gold Kiwifruit 'Haegeum' and its performances. *Acta Hortic.* 913: 249-253.
- Costa G, G Fiori, I Donati, F Spinelli, L Piccinini, G Cipriani and R Testolin (2014) Dori, nuova varietà di *A. chinensis*. *Rivista di frutticoltura e di ortofloricoltura*. 76(1): 16-21.
- Dal-Pane M (2021) Kiwi Plant named 'Red Queen' (US Patent No. USPP32895P3).
- Deka BC, A Thirugnanavel, RK Patel, A Nath and N Deshmukh (2012) Horticultural diversity in North-East India and its improvement for value addition. *Indian J. Genet. Plant Breed.* 72: 157-167.
- Dichio B, AC Tuzio, C Xiloyannis, G Rigo, R Lovato, G Comuzzo, R Frezza, D Macor, G Cipriani, R Testolin and S Micolini (2015) The new yellow-fleshed kiwifruit (*Actinidia chinensis* Pl.) 'Soreli'; conclusions from six years of cultivation in different climatic areas. *Proceedings of XIII IS on Kiwifruit*. In: Hongwen Huang and Qiong Zhang (eds). *Acta Hortic.* 1096: 149-154.
- Feng XZ and DD Huang (2025) Recent advances in hybrid breeding of kiwifruit; strategies and outcomes. *Tree Genet. Mol. Breed.* 15(2): 70-79.
- Fukuda T, K Suezawa and T Katagiri (2007) New kiwifruit cultivar 'Sanuki Gold'. *Acta Hortic.* 753: 243-246.
- Govaerts R, EN Lughadha, N Black, R Turner and A Paton (2021) The world checklist of vascular plants, a continuously updated resource for exploring global plant diversity. *Sci. Data* 8(1): 1-10.
- Hanley Z (2018) Kiwifruit (*Actinidia* spp.) breeding. In: *Advances in Plant Breeding Strategies: Fruits*, Vol. 3. Springer; Cham, Switzerland, pp 377-401.
- Henare SJ (2016) Chapter 15-The nutritional composition of kiwifruit (*Actinidia* spp.). In: Simmonds MSJ and Preedy VR (eds) *Nutritional Composition of Fruit Cultivars*. Academic Press, San Diego, pp 337-370.
- Hongwen H (2014) Species. In: *Kiwifruit: The Genus Actinidia*. Science press, Beijing, pp 49-57.
- Huang H, S Wang, R Huang, Z Jiang and Z Zhang (2002) 'Jintao', a novel, hairless, yellow-fleshed kiwifruit. *Hortic. Sci.* 37(7): 1135-1136.
- Huang H, Y Wang, Z Zhang, Z Jiang and S Wang (2004) *Actinidia* germplasm resources and kiwifruit industry in China. *Hortic. Sci.* 39: 1165-1172.
- Jhalegar MJ, RR Sharma, RK Pal, A Arora and A Dahuja (2011) Analysis of physiological and biochemical changes in kiwifruit (*Actinidia deliciosa* cv. Allison) after the postharvest treatment with 1-Methylcyclopropene. *J. Plant Biochem. Biotechnol.* 20: 205-210.
- Joshi BD (1998) Indigenous horticultural crops of the Indian Himalayas. In: Partap T and Sthapit B (eds) *Managing Agrobiodiversity: Farmers' Changing Perspectives and Institutional Responses in HKH region*. International Centre for Integrated Mountain Development, Kathmandu, Nepal, pp 205-233.
- Jung BJ, HS Cho, MY Park and YS Cho (2016) 'Mansu', a hardy kiwifruit (*Actinidia arguta* Planch. et Miq.) cultivar with improved storage life. *Korean J. of Hortic. Sci. Tech.* 34(5): 755-760.
- Kim CH, SC Kim, KC Jang, EY Song, M Kim, DY Moon, KC Seong, JS Lee, HD Suh and KJ Song (2007) A new kiwifruit cultivar, 'Jecy Gold' with yellow flesh. *Korean J. Breed. Sci.* 39: 258-259.
- Kim DG, YG Jin, JY Jin, S Kim, SC Kim, CH Han and YJ Lee (2011) Effects of the *Actinidia chinensis* on loperamide-induced constipation in rat. *Korean J. Plant Res.* 24(1): 61-68.
- Kim SC, EY Song and CH Kim (2012) A new kiwifruit variety, 'Halla Gold' with high soluble solids content and early harvesting. *Hortic. Sci. Technol.* 30(3): 334-337.
- Kishor A, SK Verma, M Brijwal, R Narayan, A Kumar, S Debnath and MS Mer (2017) Yield and Physico-chemical performance of different kiwifruit cultivars in Kumaon hills of Uttarakhand. *Res. Crops.* 18(2): 256-259.
- Kumar A, DP Sharma, P Kumar and P Verma (2026) Chimeric apple; a study of morphological and biochemical variations. *Indian J. Plant Genet. Resour.* 39(1): 56-61.
- Kwack YB, HS Choi, WB Chae and MI Jeong (2010) 'Skinny Green', a novel hairless green-fleshed baby kiwifruit. *Korean J. Hortic. Sci. Technol.* 28: 708-710.
- Kwack YB, HL Kim, WB Chae, DW Lee, HS Choi and KH Kang (2011) 'Goldrush', a newly bred Kiwifruit Cultivar in Korea. *Acta Hortic.* 913: 181-183.
- Kwack YB, HL Kim, JH Lee, KH Chung and WB Chae (2017) 'Goldone', a yellow-fleshed kiwifruit cultivar with large fruit size. *Hortic. Sci. Technol.* 35(1): 142-146.
- Kwack YB, MH Lee, HL Kim, WB Chae, SH Kim and SK Kang (2022) Green-fleshed sweet kiwifruit 'Garmrok'. *J. Korean Soc. Int. Agric.* 34: 43-48.
- Latocha P (2012) Some morphological and biological features of 'Bingo'; a new hardy kiwifruit cultivar from Warsaw University of Life Sciences (WULS) in Poland. *Rocznik Polskiego Towarzystwa Dendrologicznego* 60: 61-67.
- Latocha P (2017) The Nutritional and Health Benefits of Kiwiberry (*Actinidia arguta*); a Review. *Plant Foods Hum. Nutr.* 72: 325-334.
- Latocha P and T Krupa (2007) Morphological, chemical and sensory analyses of promising genotypes of hardy kiwifruit (*Actinidia* Lindl.) obtained in the breeding programme at WULS. *Annals of Warsaw University of Life Sciences- SGGW Horticulture and Landscape Architecture*, 28: 111-119.
- Latocha P, R Wolosiak, E Worobiej and T Krupa (2013) Clonal differences in antioxidant activity and bioactive constituents of hardy kiwifruit (*Actinidia arguta*) and its year-to-year variability. *J. Sci. Food Agric.* 93: 1412-1419.
- Lawless HT and H Heymann (2010) Sensory evaluation of food. *Principles and Practices*, 2nd edn. Springer, New York, pp 326-329.
- Lee MH, HL Kim, HC Rhee, YB Kwack, HMPC Kumarihami and JG Kim (2020) Evaluation of the genetic resources of kiwifruit with multivariate analysis. *Hortic. Sci. Technol.* 38(4): 569-581.
- Lei D, Y Lin, Q Chen, B Zhao, H Tang, Y Zhang, Q Chen, Y Wang, M Li, W He, Y Luo, X Wang, H Tang and Y Zhang (2022) Transcriptomic analysis and the effect of maturity stage on fruit quality reveal the importance of the L-galactose pathway in the ascorbate biosynthesis of hardy kiwifruit (*Actinidia arguta*). *Int. J. Mol. Sci.* 23(12): 6816.
- Li HL (1952) A taxonomic review of the genus *Actinidia*. *J. Arnold Arbor.* 33: 1-61.
- Liang J, Y Ren, Y Wang, M Han, T Yue, Z Wang and Z Gao (2021) Physicochemical, nutritional, and bioactive properties of

- pulp and peel from 15 kiwifruit cultivars. *Food Biosci.* 42: 101157.
- Liu Y, B Zhou, Y Qi, X Chen, C Liu, Z Liu and X Ren (2017) Expression differences of pigment structural genes and transcription factors explain flesh coloration in three contrasting kiwifruit cultivars. *Front. Plant Sci.* 8: 1507.
- Lowe RG (2010) Kiwifruit variety named ZESY002 (U.S. Patent no. US 20100333243P1).
- Maity R, UKL Tiwari and SS Dash (2026) A new species of wild-kiwi from India. *Feddes Repertorium*. 137: 1-6.
- Mao AA and SS Dash (2020) Flowering plants of India: an annotated check I. *BSI, Kolkata*. 1: 161.
- Mc-Neillage MA, RA Beatson and EA Mac-Rae (2003a) Kiwi plant named 'HORTGEM TORU' (U.S. Plant Patent No. US PP14, 213 P3). U.S. Patent and Trademark Office.
- Mc-Neillage MA, RA Beatson and EA Mac-Rae (2003b) Kiwi plant named 'HORTGEM TAHI' (U.S. Plant Patent No. U.S. PP13, 815 P3). U.S. Patent and Trademark Office.
- Ming X, GH Jiang, XX Chen and XZ Gao (1992) Selection and breeding of kiwifruit in China. *Acta Hort.* 297: 57-60.
- Nishiyama I, Y Yamashita, M Yamanaka, A Shimohashi, T Fukuda and T Oota (2004) Varietal difference in vitamin C content in the fruit of kiwifruit and other *Actinidia* species. *J. Agric. Food Chem.* 52(17): 5472-5475. <https://doi.org/10.1021/jf049398z>
- Padhan A, A Kumar, R Pathirana, DP Sharma, DS Thakur, VS Rana, P Kumar and A Chauhan (2023) Potential of wild, underutilized *Actinidia callosa* and *Actinidia strigosa* from Northeast India for kiwifruit breeding. *Genet. Resour. Crop Evol.* 71: 39-52.
- Padhan A and DS Thakur (2025a) Morphological, biochemical and sensory analysis of selected *Actinidia callosa* var. *strigillosa* female accessions from the existing wild populations in North Sikkim, India. *N. Z. J. Crop Hortic. Sci.* 53(4): 1150-1167.
- Padhan A and DS Thakur (2025b) Preliminary selection of wild male kiwifruit (*Actinidia callosa* var. *strigillosa* C. F. Liang) accessions in North Sikkim, India, for future breeding programmes. *N. Z. J. Bot.* 63(4): 759-770. DOI: 10.1080/0028825X.2024.2333834
- Padhan A, DS Thakur and Ambrish K (2024) An endangered wild kiwi *Actinidia callosa* var. *strigillosa* C. F. Liang (*Actinidiaceae*) from India. *Curr. Sci.* 126: 1-3.
- Pescie MA and BC Strik (2004) Thinning before bloom affects fruit size and yield of hardy kiwifruit. *Hortic. Sci.* 39(6): 1243-1245.
- Raja WH, M Noomphy, G Bam, SUN Nabi, OC Sharma, JI Mir, MA Shah and GS Madhu (2021) Potential and prospects of kiwifruit production in the state of Arunachal Pradesh. *PharmaInnov.* 10(12): 1749-1751.
- Ranganna S (2009) *Handbook of Analysis and Quality Control for Fruit and Vegetable Products*. Tata McGraw Hill, New Delhi, 1112 p.
- Richardson DP, J Ansell and LN Drummond (2018) The nutritional and health attributes of kiwifruit; a review. *Eur. J. of Nutr.* 57(8): 2659-2676.
- Singleton VL and JA Rossi (1965) Colorimetry of total phenolics with phosphomolybdic phosphotungstic acid reagents. *Am. J. Enol. Vitic.* 16: 144-58.
- Skelton DA (2010) Kiwifruit plant named 'Y374' (US Patent No. US00PP21174P3).
- Stanica F, LM Iliescu and G Zuccherelli (2022) 'Vip Green' and 'Vip Red'-two new registered kiwi berry (*Actinidia arguta*) cultivars. *Proceedings of X International Symposium on Kiwifruit. Acta Hort.* 1332.
- Stone H and JL Sidel (2004) *Sensory Evaluation Practices*, 3rd edn. Elsevier Academic Press, London, pp 87-90.
- Strik BC and KE Hummer (2006) 'Ananasnaya' hardy kiwifruit. *J. Am. Pomol. Soc.* 60: 106-112.
- Testolin R and G Cipriani (2008) Soreli: una nuova varietà di Kiwi a polpa gialla. Ottenuta e diffusa dall'Università Di Udine. *Rivista Di Frutticoltura E Di Ortofloricoltura*, 70(7-8): 54-56.
- UPOV (2019) Kiwifruit (*Actinidia* Lindl.). International Union for Protection of New Varieties of Plants, Geneva, pp 48.
- US Department of Agriculture (2016) Green Kiwifruit. USDA National Nutrient Database for Standard Reference, Release 28 (slightly revised). Version: May 2016. US Department of Agriculture (USDA), Agricultural Research Service (ARS), Nutrient Data Laboratory, Beltsville (MD).
- US Plant Patent (2010) United States Plant Patent, Patent No: US PP20,721 P3, 3 p.
- Wang Y, CI Zhao, JY Lia, YJ Liang, RQ Yang, JY Liu, Z Maa and L Wu (2018) Evaluation of biochemical components and antioxidant capacity of different kiwifruit (*Actinidia* spp.) genotypes grown in China. *Biotechnol. Biotechnol. Equip.* 32(3): 558-565.
- Wen J, Y Wang, Y He, N Shu, W Cao, Y Sun, P Yuan, B Sun, Y Yan, H Qin, S Fan and W Lu (2023) Flavour quality analysis of ten *Actinidia arguta* fruits based on high-performance liquid chromatography and headspace gas chromatography-ion mobility spectrometry. *Molecules* 28: 7559.
- Wi TS, GS Lee, MH Lee, YM Son and SC Go (2019) Consumer and distribution trends of kiwifruit. *Proceedings of the symposium*, Rural Development Administration, Wanju-si, Republic of Korea, pp 9-25.
- Williams MH, LM Boyd, MA Mc-Neillage, EA Mac-Rae, AR Ferguson, RA Beatson and PJ Martin (2003) Development and commercialization of 'Baby Kiwi' (*Actinidia arguta* Planch.). *Proceedings of International Symposium on Kiwifruit. Acta Hort.* 610: 81-86.
- Zhang H, Q Zhao, T Lan, T Geng, C Gao, Q Yuan, Q Zhang, P Xu, X Sun, X Liu and T Ma (2020) Comparative analysis of physicochemical characteristics, nutritional and functional components and antioxidant capacity of fifteen kiwifruit (*Actinidia*) cultivars- comparative analysis of fifteen kiwifruit (*Actinidia*) cultivars. *Foods* 9(9): 1267.
- Zhang Y, C Zhong, Y Liu, Q Zhang, X Sun and D Li (2017) Agronomic trait variations and ploidy differentiation of kiwi berries in northwest China; implication for breeding. *Front. Plant Sci.* 8: 711.
- Zhong C, S Wang, H Huang and Z Jiang (2011) 'Jinyan'; a superior yellow-fleshed kiwifruit cultivar with excellent storage quality. *Acta Hort.* 913: 135-143.

Mean values of traits for selection of nine superior accessions

Serial Number	Accession name	Fruit weight (g)	Titratable acidity (%)	Total sugars (%)	Sugar: acid ratio	Ascorbic acid/ Vitamin C (mg/100g)	Fruit yield (kg/vine)	Colour of Outer Pericarp	Colour of Locule	Colour of Core
1	SKNLA-01	7.55	0.77	8.57	11.13	39.00	3.00	Dark green	Light green	White
2	SKNLA-02	7.23	0.70	6.60	9.43	31.20	2.50	Dark green	Light green	White
3	SKNLA-03	7.71	1.05	7.89	7.51	65.00	5.50	Dark green	Light green	White
4	SKNLA-04	7.13	0.91	7.69	8.45	33.80	4.50	Dark green	Light green	White
5	SKNLA-05	7.84	0.98	8.82	9.00	36.40	3.50	Dark green	Medium green	Yellow white
6	SKNLA-06	4.19	0.63	7.79	12.37	31.20	6.50	Dark green	Light green	White
7	SKNLA-07	3.81	0.84	7.61	9.06	41.60	8.00	Dark green	Light green	White
8	SKNLA-08	3.93	0.63	6.25	10.42	44.20	2.50	Dark green	Dark green	White
9	SKNLA-09	3.12	0.49	5.88	12.00	36.40	1.80	Light green	Light green	White
10	SKNLA-10	4.63	0.91	6.86	7.54	39.00	6.50	Dark green	Light green	White
11	SKNLA-11	3.69	1.05	10.52	10.02	41.60	8.50	Dark green	Dark green	White
12	SKNLA-12	3.51	0.77	8.10	10.52	44.20	10.50	Dark green	Light green	White
13	SKNLA-15	4.62	0.49	7.22	14.73	65.00	5.50	Medium green	Medium green	White
14	SKNLA-16	5.18	0.77	8.33	10.82	91.00	4.50	Medium green	Light green	White
15	SKNLA-17	5.43	0.63	6.66	10.57	59.80	7.50	Dark green	Light green	White
16	SKNLA-18	3.74	1.05	8.21	7.82	33.80	11.00	Medium green	Light green	White
17	SKNLA-19	4.85	0.91	7.05	7.75	46.80	3.50	Dark green	Light green	White
18	SKNLA-20	5.48	0.84	6.97	8.30	31.20	2.00	Dark green	Medium green	White
19	SKNLA-21	6.34	0.56	6.52	11.64	62.40	2.50	Dark green	Light green	White
20	SKNLA-22	5.85	0.77	8.47	11.00	70.20	4.50	Dark green	Light green	White
21	SKNLA-23	5.32	0.63	5.45	8.65	85.80	5.50	Dark green	Light green	White
22	SKNLA-24	3.27	0.84	6.25	7.44	46.80	6.50	Dark green	Light green	White
23	SKNLA-25	4.75	0.98	6.67	6.81	57.20	9.50	Dark green	Medium green	White
24	SKNLA-26	4.66	0.63	7.50	11.90	41.60	8.50	Dark green	Light green	White
25	SKNLA-27	4.08	0.70	7.69	10.99	39.00	5.50	Dark green	Light green	White
26	SKNLA-28	5.34	0.56	5.26	9.39	62.40	6.50	Dark green	Light green	White
27	SKNLA-29	6.31	0.49	5.76	11.76	75.40	8.17	Dark green	Light green	White
28	SKNLA-30	6.92	0.56	6.74	12.04	33.80	2.50	Medium green	Light green	White
29	SKNLA-31	5.95	0.98	7.89	8.05	46.80	11.00	Dark green	Light green	White
30	SKNLA-32	5.97	0.49	4.68	9.55	44.20	19.00	Dark green	Light green	White
31	SKNLA-33	4.37	0.63	6.38	10.13	31.20	14.50	Light green	Light green	White
32	SKNLA-34	4.86	0.77	6.74	8.75	44.20	12.50	Dark green	Light green	White
33	SKNLA-35	5.15	0.91	7.31	8.03	52.00	15.50	Dark green	Light green	White
34	SKNLA-36	5.31	1.05	10.34	9.85	41.60	14.50	Light green	Light green	White
35	SKNFF-01	5.28	0.56	6.25	11.16	33.80	2.00	Light green	Light green	White
36	SKNFF-03	3.06	0.49	6.52	13.31	31.20	9.00	Light green	Light green	White
37	SKNFF-05	2.98	0.56	6.81	12.16	106.60	18.50	Medium green	Light green	White
38	SKNFF-07	2.94	0.63	7.50	11.90	36.40	2.50	Greenish yellow	Greenish yellow	Yellow white

39	SKNFF-08	3.46	0.56	5.17	9.23	39.00	2.50	Dark green	Light green	White
40	SKNBC-01	6.33	0.91	9.83	10.80	31.20	3.50	Light green	Light green	White
41	SKNBC-02	3.92	1.05	9.90	9.43	33.80	5.50	Dark green	Dark green	White
42	SKNBC-03	6.56	0.91	6.67	7.33	31.20	2.00	Dark green	Light green	White
43	SKNBC-04	4.07	0.84	6.38	7.60	54.60	6.50	Medium green	Light green	White
44	SKNBC-05	8.05	0.56	4.61	8.23	41.60	20.50	Light green	Light green	White
45	SKNTL-01	3.52	0.70	6.60	9.43	67.60	2.50	Dark green	Light green	White
46	SKNLC-01	2.94	0.56	6.89	12.30	39.00	7.50	Dark green	Dark green	White
47	SKNLC-02	3.71	0.84	9.52	11.33	33.80	2.50	Dark green	Dark green	White
48	SKNLC-03	2.92	0.70	6.38	9.11	44.20	5.50	Dark green	Medium green	White
49	SKNLC-04	2.69	0.63	6.25	9.92	36.40	22.50	Dark green	Light green	White
50	SKNLC-05	1.76	0.91	8.57	9.42	44.20	4.50	Dark green	Light green	White
51	SKNLC-06	1.57	0.84	8.33	9.92	41.60	22.50	Dark green	Light green	White
52	SKNLC-07	5.28	1.19	9.83	8.26	31.20	3.50	Medium green	Light green	Greenish white
53	SKNLC-08	4.36	0.49	6.67	13.61	26.00	4.50	Light green	Light green	White
54	SKNLC-09	2.64	0.77	8.57	11.13	59.80	7.60	Dark green	Light green	White
55	SKNLC-10	3.83	0.70	5.35	7.64	33.80	26.50	Dark green	Light green	White
56	SKNLC-11	6.28	0.56	4.68	8.36	49.40	7.50	Medium green	Light green	White
57	SKNLC-12	6.02	0.63	6.25	9.92	36.40	9.50	Dark green	Medium green	White
58	SKNLC-14	8.07	0.77	8.57	11.13	54.60	9.00	Dark green	Dark green	White
59	SKNLC-15	6.37	0.63	8.45	13.41	31.20	4.50	Dark green	Dark green	Yellow white
60	SKNLC-16	4.14	0.49	6.81	13.90	33.80	10.50	Medium green	Light green	White
61	SKNLC-17	2.96	1.26	9.37	7.44	41.60	8.50	Dark green	Light green	White
62	SKNLC-18	9.16	1.05	7.55	7.19	46.80	9.00	Dark green	Light green	White
63	SKNLC-19	3.49	0.56	5.76	10.29	36.40	6.50	Medium green	Light green	White
64	SKNLC-20	5.22	0.63	6.38	10.13	44.20	14.50	Medium green	Light green	White
65	SKNCH-01	4.71	0.49	6.59	13.45	70.20	9.50	Dark green	Light green	White
66	SKNCH-02	5.62	0.77	6.81	8.84	52.00	3.50	Medium green	Light green	White
67	SKNCH-03	6.68	0.84	7.14	8.50	80.60	4.50	Dark green	Dark green	Greenish white
68	SKNCH-04	6.51	0.91	8.33	9.15	36.40	5.50	Dark green	Light green	White
69	SKNZM-01	6.34	0.77	8.04	10.44	59.80	4.50	Dark green	Light green	White
70	SKNZM-02	5.82	1.05	10.60	10.10	31.20	7.50	Dark green	Light green	White
71	SKNZM-03	4.32	0.77	7.63	9.91	44.20	4.50	Dark green	Medium green	White
72	SKNZM-04	4.71	0.56	6.67	11.24	31.20	5.50	Light green	Dark green	White
73	SKNZM-05	2.24	0.49	5.26	10.73	46.80	3.50	Dark green	Light green	White
74	SKNZM-06	6.25	0.56	6.25	11.16	41.60	1.80	Dark green	Dark green	White
75	SKNZM-07	6.77	0.42	5.17	12.31	33.80	3.00	Medium green	Light green	White
76	SKNZM-08	5.29	0.77	6.25	8.12	39.00	14.50	Dark green	Medium green	Yellow white
77	SKNZM-09	4.25	0.56	5.26	9.39	28.60	28.50	Dark green	Medium green	White
78	SKNZM-10	5.32	0.91	6.81	7.48	46.80	16.00	Dark green	Light green	White
79	SKNZM-11	5.76	0.84	5.45	6.49	33.80	12.50	Dark green	Light green	White
80	SKNZM-12	4.49	0.98	7.50	7.65	39.00	4.50	Medium green	Dark green	White
81	SKNZM-13	6.63	0.49	5.45	11.12	33.80	14.60	Medium green	Light green	Yellow white

RESEARCH ARTICLE

Present status of *Pubistylus andamanensis* Thoth. in Andaman Islands and its IUCN Red List Assessment

CS Purohit¹, Vivek CP^{2*}, BC Dey³ and LJ Singh⁴

Abstract

The endemic genus *Pubistylus* Thoth. was first described from the Interview Island in the North Andamans. The genus is represented solely by the species, *Pubistylus andamanensis* Thoth. It was previously known by a few collections from the North Andaman and Little Andaman Islands. The present authors have recently recorded the species from the Middle Andaman Island, showing its range extension in the Andaman group of Islands up to the Middle Andamans. Considering the limited distribution in the new locality and the decline in previously recorded areas, the species has been assessed as Endangered by applying the IUCN Red List Category and Criteria. As part of the ongoing activities on *ex-situ* conservation of endemic plants, the species has been introduced in the Dhanikhari Botanical Garden cum Arboretum of Botanical Survey of India, Andaman and Nicobar Islands.

Keywords: Conservation, Endemic Genus, Middle Andaman Island, *Pubistylus*, Range extension

¹Botanical Survey of India, Arid Zone Regional Centre, Jodhpur, Rajasthan, 342008, India.

²Botanical Survey of India, Headquarter, Kolkata, 700064, India.

³Botanical Survey of India, Acharya Jagadish Chandra Bose Indian Botanic Garden, Howrah, West Bengal, 711103, India.

⁴Botanical Survey of India, Andaman & Nicobar Regional Centre, Port Blair, 744102, India.

***Author for correspondence:**
vvkcpoulose@gmail.com

Received: 11/01/2025

Revised: 15/04/2026

Accepted: 22/05/2026

How to cite this article: Purohit CS, CP Vivek, BC Dey and LJ Singh (2026) Present status of *Pubistylus andamanensis* Thoth. in Andaman Islands and its IUCN Red List Assessment *Indian J. Plant Genet. Resour.* 39(2): 100-104.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.178050>

Introduction

The endemic genus *Pubistylus* Thoth. is one of the two monotypic genera in the tribe Albertaeae of family Rubiaceae. *Octotropis* Bedd. is the other member that has been described from Travancore Hills in 1874. The type species, *Pubistylus andamanensis* Thoth. was described in 1966 from Interview Island in North Andamans. The species was subsequently collected from different localities in Little Andaman Islands (Hut Bay & Dugong Creek by N Bhargava; Quarryroad & 4 km north from Hut Bay by L Rasingam) and North Andaman (Interview Island by K Thothathri, East Coast by Lakshminarasimhan, West Coast and cave forest, by GS Lakra & VM Radhakrishnan). There was no additional information regarding its presence in the Andaman and Nicobar Islands until the present authors identified approximately 25 mature individuals, along with a few seedlings, collectively from Chitragud hill forests and Adajig forest in the Middle Andamans. Considering the restricted distribution and vulnerability of its habitat, the species has been assessed for the IUCN Red List Category.

Materials and methods

Survey and identification

Extended distribution of *Pubistylus andamanensis* was recorded during a floristic survey in 2021 at Chitragud hills and




Fig. 1: Map showing the distribution of *Pubistylus andamanensis* Thoth. in Andaman & Nicobar Islands.

Adajig forest in the Middle Andamans (Fig.1). The voucher specimens were prepared and deposited in the herbarium of Botanical Survey of India, Andaman and Nicobar Regional Centre (PBL). The identity and the present distributional status of the species were determined with the help of relevant literatures (Thothathari, 1966; Gangopadhyay & Chakrabarty, 1988, 1993; Bhakat, 1990; Ali & Robbrecht, 1991; Deb & Rout, 1993; Hajra *et al.*, 1999; Sinha, 1999; Pandey & Diwakar, 2008; Selvan, 2014; Murugan *et al.*, 2016; Mao&Dash, 2020; Kamble, 2020; Rasingam & Diwakar, 2021; Singh *et al.*, 2021; Purohit & Vivek, 2022; Purohit *et al.*, 2023) and herbarium specimens deposited at PBL, CAL, K, and BLAT.

Assessment with Geo CAT (geospatial conservation assessment tool)

Current and past distribution maps of the species were prepared with the help of GPS coordinates of collection localities of earlier reports and the present survey (Fig.1) and its population data were used to assess the IUCN threat category (IUCN, 2013). Online software GeoCAT (<http://geocat.kew.org/>) was used to calculate the Extent Of Occurrence (EOO) and Area Of Occupancy (AOO) by keeping a grid size 4×4 km (Fig. 2 & 3).




Fig. 2: Map showing EOO & AOO of *Pubistylus andamanensis* Thoth. in Andaman & Nicobar Islands.




Fig. 3: Convex hull of occurrence points for analysis of EOO and AOO using GeoCAT.

Results and discussion

The species has been dealt with in the paper with a detailed description, distribution, ecology, notes, threat status, distribution maps, AOO & EOO maps and photographs.

Pubistylus andamanensis Thoth., Reinwardtia 7: 283. 1966. Bhakat, *J Econ.Taxon.Bot.* 14 (1): 215. 1990; Ali & Ribbrecht, *Blumea* 35(2): 305. 1991. *Diplospora parkinsonii* Gang. et T Chakrab., *J Econ. Taxon. Bot.* 12 (2): 497. 1988; Gangopadhyay & Chakrab., *J Econ. Taxon. Bot.* 17 (2): 439. 1993. *Diplospora andamanensis* (Thoth.) Gang & T Chakrab., *J Econ. Taxon. Bot.* 18 (2): 397. 1994;

AA Mao & SS Dash, *Fl. Pl. India Annot. Checkl.* 1:698. 2020.

Type: Interview Island, 12.02.1959, *K Thothathri* 9204A (holo CAL-551295). Local name: *Haddi-Balli*.

Shrub or small tree, 2–4 m high, glabrous; branches more or less drooping, glabrous; stem angular. Leaves elliptic to lanceolate, 8–9 × 2–3.5 cm, entire, membranous, caudate-acuminate at apex, cuneate at base, glabrous; petiole 5.5–8 mm long; stipule broadly ovate, acute. Inflorescence panicked cyme, 3.5–5.5 cm long, axillary or rarely terminal. Flowers white, 3.2–5.2 mm long; pedicels slender, 4.5–8 mm long; bract small, ovate, ciliate; bracteole just below the middle of the pedicel. Calyx tube turbinate, 1–2 mm long, slightly lobed. Corolla campanulate, 2.5–3.5 mm long, 5-lobed; lobes ovate, twisted, villous within. Stamens 5; filaments short; anthers 1–1.5 mm long, linear-oblong, 2-celled. Ovary 2-celled; ovule single in each cell; style clavate, 2–2.4 mm long, pubescent; stigma bifid. Fruit globular to sub-globular, deep green, glabrous, c. 1 cm in diameter (Fig. 4).

Flowering and Fruiting



Fig. 4: *Pubistylus andamanensis* Thoth.: (A-D) Natural Habitats & Habits; (E-F) Twigs showing leaf morphology; (G-I) Inflorescence & Close-up of flower; (J-M) Fruiting branches & Close-up of fruits.

August–December (All the previous records indicate that the flowering and fruiting of the species occur from August to September. However, the present study has documented flowering and fruiting in December, indicating the extension of the reproductive period.

Habitat

The species is growing in the Inland evergreen forests between 20–150 m elevations, in association with *Mesua manii* (King) Kosterm., *Cassia fistula* L., *Knema andamanica* (Warb.) WJde Wilde, *Ardisia oxyphylla* Wall. ex A.DC., *Blachia andamanica* (Kurz) Hookf, *Dillenia andamanica* CE Parkinson, *Grewia calophylla* Kurz ex Mast, *Canarium euphyllum* Kurz, *Capparis floribunda* Wight, *Drypetes longifolia* (Blume) Pax & K Hoffm, *Ficus racemosa* L., *Garcinia xanthochymus* Hookf, *Myristica andamanica* Hookf, *Terminalia procera* Roxb., *Vitex diversifolia* Kurz ex CB Clarke, etc.

Distribution

INDIA: Andaman and Nicobar Islands, North Andamans (Interview Island: East & West coast, cave forest); Little Andaman (Hut Bay, Dugong Creek, Quarry road, 4 km); Middle Andaman (Chitragud hill, Adajig forest). Endemic.

Threat

The population of *Pubistylus andamanensis* is found in fragments in its habitats with a limited number of individual plants. The stem is hard and straight. Local people prefer the stem for crafting polls and sticks. In addition, unsustainable forest clearing poses a threat to its survival.

Voucher specimens

Andaman Islands, Middle Andaman, Chitragud hills, forest trekking path, N12°32'48.79", E 92°58'21.65", 12.12.2021, CS Purohit & Vivek CP 34201 (PBL); Adajig forest, N 12°18'21.10", E 92°46'44.55", 13.12.2021, CS Purohit & Vivek CP 34212 (PBL).

Others specimen examined

Little Andaman, Way to Forest Nursery, Hut Bay, 14.08.1976, *N Bhargava* 4118 (PBL-9838, 9839); Dugong Creek, 25.11.1977, *N Bhargava* 6561 (PBL-14586, 14587); Quarry road, 22.04.2006, *L Rasingam* 20950 (PBL-24760, 24761); 4 km North from Hut Bay, 27.05.2004, *L Rasingam* 19876 (PBL-25118, 25119). North Andaman, Interview Island: East coast, 22.02.1991, *P Lakshminarasimhan* 15665 (PBL-19759, 19760); Interview Island: way to West coast, 23.09.2010, *GS Lakra & VM Radhakrishnan* 28725 (PBL-27971, 27972);

Interview Island: way to cave forest, 25.02.2011, *GS Lakra* 28855 (PBL-28317, 28318); Interview Island: way to cave forest, 26.02.2011, *GS Lakra* 28871 (PBL-28347, 28348); Interview Island, 12.02.1959, *KThothathri* 9204A (CAL-551295).

Notes

Gangopadhyay & Chakrabarty (1988) synonymized the genus *Pubistylus* Thoth. under *Diplospora* DC., highlighting the morphological similarities between the two genera. Subsequently, Bhakat (1990) identified distinct morphological characteristics of *Pubistylus*, including acute stipules, more branched cymes, lax panicle, 5-merous flowers, pubescent and clavate style, and 2-celled pendulous ovary with single ovule per cell. Ali & Robbrecht (1991), while working on the Tropical Asian and Australian species of Rubiaceae included in *Diplospora* and *Tricalysia* A.Rich. ex DC., favoured the reinstatement of *Pubistylus*. Deb & Rout (1993) supported Bhakat's argument for the reinstatement by providing additional characteristic features and clarifying the taxonomic position. Gangopadhyay & Chakrabarty (1993) recognized *Diplospora* and *Pubistylus* as distinct sections based on the comparative length of inflorescence, suggesting that *Diplospora andamanensis* (Thoth.) Gang & T Chakrab. and *Diplospora parkinsonii* Gang & T Chakrab. should be treated conspecific with *Pubistylus andamanensis*. Govaert (2003) adopted the treatment proposed by Gangopadhyay & Chakrabarty (1993). In a more recent study, Mao & Dash (2020) also treated *Pubistylus* as an established endemic genus in the Andaman Islands.

Red list assessment as per IUCN criteria

For Red List assessment of *Pubistylus andamanensis*, we used the guidelines of IUCN Red List criteria (IUCN, 2013). Criteria A has not been followed due to non-availability of sufficient quantified data on population reduction over specified time. The EOO of the species has been calculated using the GeoCAT online tool as 4,186 km² (Fig. 3), which is between the threshold of 100 km² and 5000 km², favoured its placement under B1 Endangered Category. The AOO has been calculated as 144 km² (Fig. 3) which is between the threshold of 10 km² and 500 km², favoured its placement under B2 Endangered Category.

The species had previously been documented in five sites in the Andaman Islands, of which it could be relocated at one site from Little Andaman. This indicates a decline in its population within its natural habitat. Further, the population was found to have fragmented into the newly identified 2 sites. Over the past 4 years, a

gradual decrease in the number of mature individuals has been noted, suggesting its placement in B1+2ab under the Endangered category. The consistent reduction in its AOO, EOO, and overall population justifies its placement in B1+2ab(i,ii,iv,v) under the Endangered category. Due to insufficient data or a lack of studies, Criteria C has not been applied.

The species has been distributed in small pockets in its natural habitats and the total number of mature individuals has been counted as about 103, which is between the threshold of 50 and 250 for placing it under D1 Endangered category. No sufficient data was available for applying the Criteria E. Hence, its final assessment using IUCN criteria best fitted under the category Endangered [criteria - EN/B1+2ab(i,ii,iv,v); D1].

Conclusion

The present study identifies the Chitragud hills and Adajig forest in Middle Andaman as new distribution sites for the endemic tree species, *Pubistylus andamanensis*, where they are found restricted in numbers. Approximately 25 individual plants have been located collectively in the newly recognized sites. This species was previously documented in five different localities from the Little Andaman Islands and North Andaman Islands (collection from five different localities in the Interview Islands has been considered as a single location). Of these five localities, it has been able to relocate from only one site in the Little Andamans. The EOO and AOO have exhibited a gradual decrease, attributed to their limited distribution, potential habitat loss, and declining population. It has been noted that several of the areas where the species occurs are currently under threat from various biotic pressures. Following the guidelines of the IUCN Red List, this endemic tree species has been assessed globally as Endangered [EN/B1+2ab(i,ii,iv,v); D1].

The species has been conserved *ex-situ* in the Dhanikhari Experimental Gardencum Arboretum of Botanical Survey of India (BSI), Andaman and Nicobar Islands, with a view to multiplying and reintroducing the species in its natural habitats. Furthermore, BSI educates the local settlers about the importance of conserving the biodiversity of the Islands.

Acknowledgements

The authors are grateful to the Director, Botanical Survey of India, Kolkata, for facilities and encouragement. Thanks also extend to Dr MC Naik for valuable input. Authors are also grateful to the Forest Department, Andaman & Nicobar Islands, for provide facility during the field survey.

References

- Ali SJ and E Robbrecht (1991) Remarks on the tropical Asian and Australian taxa included in *Diplospora* or *Tricalysia* (Rubiaceae-Ixoroideae-Gardenieae). *Blumea* 35: 279-305.
- Sinha BK (1999) Flora of Great Nicobar Island. Botanical Survey of India, Kolkata.
- Bhakat RK (1990) Taxonomic notes on the status of monotypic genus *Pubistylus* Thoth. (Rubiaceae). *J. Econ. Taxon. Bot.* 14: 215-216.
- Deb DB and RC Rout (1993) The Indian genus *Pubistylus* (Rubiaceae): An investigation into the characteristics and taxonomic position. *Opera Bot. Belg.* 6: 93-100.
- Gangopadhyay M and T Chakrabarty (1988) *Pubistylus* Thoth. (Rubiaceae) reduced to *Diplospora* DC. *J. Econ. Taxon. Bot.* 12: 495-497.
- Gangopadhyay M and T Chakrabarty (1993) Additional notes on the status of *Pubistylus* Thoth. (Rubiaceae). *J. Econ. Taxon. Bot.* 17: 439-440.
- GeoCAT Geospatial Conservation Assessment Tool <<https://geocat.iucnredlist.org/>>
- Hajra PK, PSN Rao and V Mudgal (1999) Flora of Andaman & Nicobar Islands, Vol1: Ranunculaceae to Combretaceae. Botanical Survey of India, Kolkata.
- IUCN (2013) The IUCN Red List of Threatened Species. Version 2013.2. <<http://www.iucnredlist.org>>
- Kamble MY (2020) Flora of North Andaman Islands (in reference to the six wildlife sanctuaries). Botanical Survey of India, Kolkata.
- Mao AA and SS Dash (2020) Flowering plants of India: An Annotated Checklist (Dicotyledons). Botanical Survey of India.
- Murugan C, S Prabhu, R Sathiyaseelan and RP Pandey (2016) A Checklist of Plants of Andaman and Nicobar Islands. ENVIS Centre on Floral Diversity. Botanical Survey of India, Kolkata. http://bsienvis.nic.in/Database/Checklist-of-Andaman-Nicobar-Islands_24427.aspx.
- Pandey RP and PG Diwakar (2008) An integrated checklist of plants in Andaman & Nicobar Islands, India. *J. Econ. Taxon. Bot.* 32: 403-500.
- Purohit CS and CP Vivek (2022) Project Report on "Curatorial work of Botanic Garden (Multiplication and nursery development of Bamboos, Palms, Zingibers, Endemic tree species) and raise nursery", submitted to Director, BSI, Kolkata.
- Purohit CS, LJ Singh, CP Vivek and BC Dey (2023) Project Report on "Conservation Assessment, ENM studies including GIS mapping of Endemic trees of Andaman & Nicobar Islands", submitted to the Director, BSI, Kolkata.
- Rasingam L and PG Diwakar (2021) Flora of Little Andaman Islands, India. Botanical Survey of India, Kolkata.
- Selvan T (2014) Vegetation pattern of commercially important trees in evergreen forests of Baratang Island, Andamans. *J. Tree Sc.* 33(1): 12-21.
- Singh LJ, V Ranjan, BK Sinha, S Mishra, CS Purohit, CP Vivek, MC Naik and GA Ekka (2021) An Overview of Phytodiversity of the Andaman and Nicobar Islands, India. In: Singh LJ and V Ranjan (eds) *New Vistas in Indian Flora*. Bishen Singh and Mahendra Pal Singh, Dehradun, India 2: 381-399.
- Thothathari K (1966) *Pubistylus* Thoth. - An interesting new genus of Rubiaceae from Andaman Islands. *Reinwardtia* 7(3): 283-286.

RESEARCH ARTICLE

Correlation and Path Analysis for Seed Yield and Components Traits in Mothbean [*Vigna aconitifolia* (Jacq) Marechal] under Arid Environment

Savita Meena^{1,2}, NK Sharma¹, Pinky Yadav^{3*}, Rukoo Chawla¹ and Ajit Kumar Meena⁴

Abstract

Among pulses, the mothbean is an important pulse crop of hot arid regions of India and adapts to extreme ecological niches, particularly extreme drought and hot climatic conditions. Being a pulse crop, it is a cheap source of vegetable protein for balancing the nutritional deficiency, most commonly cultivated on less productive soils, on which, financially less equipped, a great portion of people have been depending on their livelihood. A field experiment was conducted with 36 genotypes evaluated in a randomized block design, three replications during *kharif* 2020 at Swami Keshwanand Rajasthan Agricultural University, Bikaner, to study the correlation and path coefficient analysis for seed yield and its component traits. Each genotype was sown in double rows of 3-meter length and plant-to-plant distance at 12-15 cm. For each of the 26 traits studied, genotypes differed significantly. Seed yield per plant was positively correlated with agro-morphological characters such as number of pods per plant and harvest index, as well as physiological and seed characteristics such as chlorophyll-a, chlorophyll-b, total chlorophyll content, relative water content, bulk density, and particle/true density. Days to maturity, pod length, biological yield per plant, harvest index, bulk density, particle density, seedling vigour index at 40°C, and seedling dry weight at 40°C had a positive direct effect on seed yield in mothbean. This suggests a strong hereditary link between seed yield and these traits, indicating that selection for these features may improve mothbean yield.

Keywords: Arid environment, Correlation, Mothbean, Path analysis, Seed yield

¹College of Agriculture, Swami Keshwanand Rajasthan Agricultural University, Bikaner, Rajasthan, 334006, India.

²Genetics and Plant Breeding, Rajasthan College of Agriculture, Maharana Pratap University of Agriculture and Technology, Udaipur, Rajasthan, 313001, India.

³Department of Education in Science and Mathematics, Regional Institute of Education, Mysore, NCERT, Karnataka, 570006, India.

⁴ICAR-National Bureau of Soil Survey and Land Use Planning, Regional Centre, Udaipur, Rajasthan, 31300, India.

*Author for correspondence:

yadavpinky@riemysore.ac.in

Received: 10/01/2026

Revised: 05/05/2026

Accepted: 20/05/2026

How to cite this article: Meena S, NK Sharma, P Yadav, R Chawla and AK Meena (2026) Correlation and Path Analysis for Seed Yield and Components Traits in Mothbean [*Vigna aconitifolia* (Jacq) Marechal] under Arid Environment. *Indian J. Plant Genet. Resour.* 39(2): 105-113.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.175071>

Introduction

Pulses are an important part of the diet because they are rich in protein (20-25%) and can help alleviate health issues such as obesity, diabetes, and malnutrition. The mothbean is a popular pulse crop in India's hot, dry regions, and it responds to a variety of ecological niches, including low rainfall (250-450 mm) and hot weather. Mat bean, Matki, Turkish gram, and Dew bean are all names for the same thing (Yaqoob *et al.*, 2015). It belongs to the family Leguminaceae/Fabaceae, sub-family Papilionaceae, and is a self-pollinated diploid crop with $2n=2x=22$ chromosomes (Verdcourt, 1970). Dal (Mogar) and sprouts are the most common forms of consumption. It's the main ingredient in Bikaneri Bhujia, a popular spicy snack. It's also used in the making of various foods, including papad, namkeen, mangori, and dal vada, among others. Agri-industries in Bikaner consume almost 80% of Rajasthan's mothbean production (Sharma, 2016). The cultivation of mothbean is integral to arid agro-ecosystems, as it significantly contributes to crop diversification and long-term sustainability of agricultural production (Sharma, 2016).

Materials and Methods

Plant material and experimental details

The current study was undertaken at the experimental farm of Swami Keshwanand Rajasthan Agricultural University in Bikaner, with sowing done during the month of July of *kharif* 2020. Under normal soil and water conditions, the experimental material, which consisted of 36 genotypes, was examined in a randomised block design with three replications, accommodating a 3-metre length, two rows in each replication, at 30 cm spacing and plant-to-plant distance of 12-15 cm.

Data recording

Observations were recorded for 26 characters *viz.*, days to 50 percent flowering, days to maturity, plant height, number of pods per plant, number of seeds per pod, pod length, 100-seed weight, biological yield per plant, harvest index, seed yield per plant, chlorophyll-a, chlorophyll-b, total chlorophyll content (estimated by the method of Hiscox and Israelstom, 1979), membrane stability index (estimated by the method of Premchand *et al.*, (1990) and modified by Sairam *et al.*, (1997) , relative water content, seed volume (liquid displacement technique – Shepherd and Bhardwaj, 1986), bulk density (method reported by Singh and Goswami, 1996 and Gupta and Das, 1997), particle/true density (Toluene displacement method, Mohsenin, 1986 and Singh and Goswami, 1996), porosity (formula used by Singh and Goswami, 1996), water absorption capacity (recorded by formula, Mohsenin, 1986), water absorption index recorded by formula Williamsa *et al.*, 1993), and seedling vigour index at 40°C (determined by the multiplying the seedling length with concerned germination percentage by the following formula, Iqbal and Rahmati, 1992).

Statistical analysis

The Dewey and Lu (1959) approach was used to calculate phenotypic and genotypic correlations and path coefficients of variation.

Results and Discussion

The analysis of variance demonstrated significant variation in seed yield and yield-contributing factors across 36 mothbean genotypes (Table 1) for all 26 characters studied, indicating that there is scope for change in seed yield and yield-contributing characters through selection (Table 2).

Table 1: List of 36 genotypes and source of mothbean genotypes included in the present study

S. NO.	Name of germplasm	Source of procurement
1	IC 983	NBPGR, Regional Station, Jodhpur
2	IC 120966	NBPGR, Regional Station, Jodhpur
3	IC 120968	NBPGR, Regional Station, Jodhpur
4	IC 121005	NBPGR, Regional Station, Jodhpur
5	IC 121064	NBPGR, Regional Station, Jodhpur
6	IC 121065	NBPGR, Regional Station, Jodhpur
7	IC 140616	NBPGR, Regional Station, Jodhpur
8	IC 140663	NBPGR, Regional Station, Jodhpur
9	IC 251877	NBPGR, Regional Station, Jodhpur
10	IC 310670	NBPGR, Regional Station, Jodhpur
11	IC 311396	NBPGR, Regional Station, Jodhpur
12	IC 311399	NBPGR, Regional Station, Jodhpur
13	IC 324034	NBPGR, Regional Station, Jodhpur
14	IC 329040	NBPGR, Regional Station, Jodhpur
15	IC 370469	NBPGR, Regional Station, Jodhpur
16	IC 415104	NBPGR, Regional Station, Jodhpur
17	IC 415116	NBPGR, Regional Station, Jodhpur
18	IC 415138	NBPGR, Regional Station, Jodhpur
19	IC 415167	NBPGR, Regional Station, Jodhpur
20	GP- 47	Agricultural Research Station, Bikaner
21	GP-387	Agricultural Research Station, Bikaner
22	GP-722	Agricultural Research Station, Bikaner
23	GP-940	Agricultural Research Station, Bikaner
24	PLMO- 16	NBPGR, Regional Station, Jodhpur
25	MOTH- 259	NBPGR, Regional Station, Jodhpur
26	Jadiya	NBPGR, Regional Station, Jodhpur
27	Jwala	NBPGR, Regional Station, Jodhpur
28	RMO- 40	NBPGR, Regional Station, Jodhpur
29	RMO- 225	Agricultural Research Station, Bikaner
30	RMO-2251	Agricultural Research Station, Bikaner
31	RMO- 257	Agricultural Research Station, Bikaner
32	RMO- 423	NSP, SKRAU, Bikaner
33	RMO- 435	Agricultural Research Station, Bikaner
34	RMB- 25	NSP, SKRAU, Bikaner
35	FMM-12-6-134	NBPGR, Regional Station, Jodhpur
36	CAZARI-MOTH-2	NSP, SKRAU, Bikaner

Table 2: Analysis of variance (mean sum of squares) for various characters in mothbean genotypes

S.No.	Characters	Replications	Sources of variation Genotypes	Error
	D.F.	2	35	70
1	Days to 50 percent flowering	44.6203	388.6052**	0.9727
2	Days to maturity	34.4814	195.5013**	7.3862
3	Plant height (cm)	71.76584	103.9436**	26.3914
4	Number of pods per plant	62.9242	486.7185**	26.8773
5	Number of seeds per pod	0.0708	0.6557**	0.2731
6	Pod length (cm)	0.0605	0.1753**	0.084
7	100-seed weight (g)	0.0338	0.2785**	0.0916
8	Biological yield per plant (g)	49.5988	696.4172**	16.6979
9	Harvest index (%)	0.0078	193.9523**	2.7607
10	Seed yield per plant (g)	1.2872	9.5413**	0.427
11	Chlorophyll-a (mg/g)	0.1101	0.1465**	0.0353
12	Chlorophyll-b (mg/g)	0.0023	0.0013**	0.0006
13	Total chlorophyll content (mg/g)	0.0616	0.154**	0.0198
14	Membrane stability index	13.919	329.4652**	27.7779
15	Relative water content (%)	204.1605	143.7202**	70.9955
16	Seed volume (µl/seed)	15.2358	25.6626**	5.9301
17	Bulk density (g/cm ³)	0.0013	0.0092**	0.0029
18	Particle/ true density (g/cm ³)	0.0028	0.0503**	0.0153
19	Porosity (%)	8.7161	36.8021**	11.4868
20	Water absorption capacity (mg/seed)	18.1545	145.8899**	21.3937
21	Water absorption index (WAI)	0.0007	0.1024**	0.0254
22	Germination percentage at 40°C	137.037	228.2275**	103.7037
23	Seedling length (cm) at 40°C	10.0051	8.48101**	3.2079
24	Seedling fresh weight (mg) at 40°C	0.0065	0.0163**	0.0008
25	Seedling dry weight (mg) at 40°C	0.0005	0.0004**	0.0001
26	Seedling vigour index at 40°C	50395.8867	105176.5874**	16838.2189

*Significant at P = 0.05, ** Significant at P = 0.01.

Correlation analysis

Correlation analysis provided a measure of genetic association among traits and normally used in selection, while environmental as well as genetic architecture of a genotype plays a great role in achieving higher yield combined with better quality (Chawla *et al.*, 2024). At phenotypic level, Days to 50 percent flowering exhibited a positive and significant correlation with days to maturity (0.9677**) and plant height (0.7049**). Number of pods per plant (-0.6130**), harvest index (-0.5360**) and seed yield per plant (-0.5889**) depicted negative significant correlation with days to 50 percent flowering only at phenotypic level. Days to maturity had a significant and negative relationship with the number of pods per plant (-0.6149**), harvest index (-0.5136**) and seed yield per plant (-0.5690**). The traits, namely

harvest index (0.6971**) and seed yield per plant (0.9018**) showed significant and positive correlation with number of pods per plant at phenotypic level. Pod length possessed significant positive relationship with 100 seed weight (0.4377**) at phenotypic level. Pod length (0.4377**) showed a significant and positive correlation with 100-seed weight at phenotypic level. Harvest index showed significant positive correlation with number of pods per plant (0.6971**) and seed yield per plant (0.8451**); while significant negative correlation exhibited by days to 50 percent flowering (-0.5360**), days to maturity (-0.5136**), plant height (-0.6811**) and biological yield per plant (-0.5780**) at phenotypic level, whereas it had significant and negative correlation with days to 50 percent flowering (-0.5889**), days to maturity (-0.5690**) and plant height

Table 3: Estimates of phenotypic and genotypic correlation coefficient for agro-morphological characters of mothbean genotypes

Characters		Days to 50 percent flowering	Days to maturity	Plant height	No. of pods per plant	No. of seeds per pod	Pod length	100-seed weight	Biological yield per plant	Harvest index
Days to 50 percent flowering	P	1								
	G	1								
Days to maturity	P	0.9677**	1							
	G	0.9889	1							
Plant height	P	0.7049**	0.6449**	1						
	G	0.8151	0.7634	1						
No. of pods per plant	P	-0.613**	-0.615**	-0.69**	1					
	G	-0.6311	-0.642	-0.86	1					
No. of seeds per pod	P	0.0736	0.1104	0.0441	-0.081	1				
	G	0.0956	0.1465	0.0377	-0.12	1				
Pod length	P	-0.0062	-0.028	-0.013	0.0325	0.2024	1			
	G	-0.007	-0.035	-0.084	0.043	0.0976	1			
100-seed weight	P	0.059	-0.02	0.1296	-0.171	-0.111	0.4377**	1		
	G	0.067	0.0134	0.3001	-0.205	-0.098	0.7988	1		
Biological yield per plant	P	0.1456	0.0906	0.2068	0.0084	-0.197	0.0609	0.08	1	
	G	0.1485	0.0942	0.2259	0.001	-0.245	0.089	0.1136	1	
Harvest index	P	-0.536**	-0.514**	-0.68**	0.6971**	0.2267	0.0305	-0.124	-0.578**	1
	G	-0.541	-0.524	-0.808	0.7023	0.2965	0.0471	-0.145	-0.588	1
Seed yield per plant	P	-0.5889**	-0.569**	-0.75**	0.9018**	0.0592	0.0339	-0.187	-0.143	0.8451**
	G	-0.6029	-0.591	-0.937	0.909	0.0841	0.0484	-0.213	-0.157	0.85

*Significant at $P = 0.05$, ** Significant at $P = 0.01$.

(-0.7536**) at also phenotypic level (Table 3).

A wide number of other component traits contribute to yield, a complicated polygenic trait. Based on phenotypic correlation coefficients among agro-morphological characters, the number of pods per plant and the harvest index showed positive and significant relationships with seed yield per plant. Kohakade *et al.*, (2017³), Ramakrishnan *et al.*, (2018), Singh *et al.*, (2018), Asari *et al.*, (2019), and B.B. Dhunde *et al.*, (2021) previously reported a significant positive correlation in seed yield per plant and number of pods per plant. Garg *et al.*, (2017), Sahoo *et al.*, (2018), Choudhary and Lal (2019) and Saran *et al.*, (2020) previously reported a significantly positive correlation between seed yield per plant and harvest index. The phenotypic correlation coefficient of physiological and seed parameters demonstrated a positive and significant relationship between seed yield per plant and chlorophyll a, chlorophyll b, total chlorophyll content, relative water content, bulk density, and particle/true density (Table 4). Therefore, these

characters may be emphasized for the mothbean breeding programme.

The phenotypic correlation coefficients of days to 50% flowering, days to maturity, plant height, seedling fresh weight at 40°C, and seed yield per plant were found to be significant and negative, suggesting that they might be used in breeding programmes for earliness and dwarfism (Table 5). Raturi *et al.*, (2015), Kanavi *et al.*, (2019) and Choudhary *et al.*, (2019) showed similar findings at 50 percent flowering. Choudhary *et al.*, (2021) and BB Dhunde *et al.*, (2021) found a significant negative correlation between seed yield per plant and days to maturity, and Kakani *et al.*, (2003) found a significant negative correlation between seed yield per plant and plant height.

Path coefficient analysis

Path coefficient analysis was used to determine the impact of different characters on seed yield by dividing the correlation coefficient into direct and indirect effects using seed yield as the dependent variable. For

Table 4: Estimates of phenotypic and genotypic correlation coefficients for physiological and seed characters of mothbean

Characters		CHL- a	CHL-b	Total CHL	MSI	RWC	Seed volume	Bulk density	Particle density	porosity
CHL- a	P	1								
	G	1								
CHL-b	P	0.796**	1							
	G	0.9638	1							
Total CHL	P	0.9773**	0.8894**	1						
	G	1.0063	1.0204	1						
MSI	P	-0.227	-0.254	-0.224	1					
	G	-0.242	-0.358	-0.235	1					
RWC	P	-0.15	0.2494	0.0215	0.1581	1				
	G	-0.274	0.8172	0.0901	0.2498	1				
Seed volume	P	-0.253	-0.221	-0.224	0.1453	0.3271*	1			
	G	-0.325	-0.371	-0.276	0.1944	0.4536	1			
Bulk density	P	0.5403**	0.7203**	0.6463**	-0.146	0.5034**	0.0357	1		
	G	0.7531	1.1255	0.8254	-0.185	0.9306	-0.004	1		
Particle density	P	0.4994**	0.6101**	0.5803**	0.0276	0.3793*	0.0097	0.8257**	1	
	G	0.6434	0.941	0.712	0.0384	0.7001	0.0225	1.0295	1	
Porosity	P	0.3443*	0.3335*	0.3753*	0.3266*	0.0166	-0.255	0.2528	0.5563**	1
	G	0.4951	0.6311	0.5044	0.4198	0.0682	-0.27	0.3417	0.5697	1
WAC	P	-0.086	-0.114	-0.117	0.2716	-0.19	0.1341	-0.207	-0.108	-0.017
	G	-0.088	-0.111	-0.115	0.3264	-0.334	0.1745	-0.309	-0.133	0.023
WAI	P	-0.054	-0.107	-0.094	0.2795	-0.213	0.0919	-0.069	0.0832	0.169
	G	-0.053	-0.125	-0.098	0.367*	-0.379	0.123	-0.161	0.1302	0.3053
Yield	P	0.3942*	0.5988**	0.513**	-0.268	0.4876**	-0.105	0.681**	0.5311**	0.1286
	G	0.4434	0.8114	0.5469	-0.28	0.6971	-0.112	0.8241	0.6699	0.1726

*Significant at P = 0.05, ** Significant at P = 0.01.

CHL-a; Chlorophyll-a, CHL-b; Chlorophyll-b, Total CHL; Total chlorophyll content, MSI; Membrane stability index, RWC; Relative water content.

Table 5: Estimates of phenotypic and genotypic correlation coefficient for seedling characters of mothbean at 40°C temperature

Characters		Germination percentage	Seedling length	Seedling fresh weight	Seedling dry weight	Seedling vigour index	Yield per plant
Germination percentage	P	1					
	G	1					
Seedling length	P	0.0413	1				
	G	0.1705	1				
Seedling fresh weight	P	0.1421	0.1606	1			
	G	0.458	0.089	1			
Seedling dry weight	P	0.2494	0.3897*	0.5622**	1		
	G	0.4433	0.5092	0.5361	1		
Seedling vigour index	P	0.734**	0.6529**	0.2109	0.3572*	1	
	G	0.9239	0.6748	0.2636	0.4354	1	
Yield per plant	P	0.204	-0.103	-0.342*	-0.05	0.0857	1
	G	0.2969	-0.13	-0.45	-0.042	0.0966	1

*Significant at P = 0.05, ** Significant at P = 0.01.

agro-morphological traits, at phenotypic level the highest positive direct effect on seed yield per plant was found for harvest index (0.7764) followed by biological yield per plant (0.3210), number of pods per plant (0.3192), days to maturity (0.2434) and pod length (0.0138); while the highest negative direct effect was found for days to 50 percent flowering (-0.1941), plant height (-0.0825), 100-seed weight (-0.0457) and number of seeds per pod (-0.0447). At genotypic level, the highest positive direct effect on seed yield per plant was found for days to maturity (1.0212), harvest index (0.9237), biological yield per plant (0.4273), number of pods per plant (0.2091) and pod length (0.0005); while the highest negative direct effect on seed yield per plant was found for days to 50 percent flowering (-0.9439), number of seeds per pod (-0.1167), plant height (-0.1087) and 100-seed weight (-0.0143) (Table 6) (Fig.1).

For physiological and seed traits, at phenotypic level the highest positive direct effect on seed yield per plant was reported for total chlorophyll content (3.5986), bulk density (0.1672), water absorption index (0.1420), relative water content (0.1374) and




Fig. 1: At phenotypic level, path diagram of mothbean genotypes based on agro-morphological characters for seed yield.

1, Days to 50 percent flowering; 2, Days to maturity; 3, Plant height; 4, Number of pods per plant; 5, Number of seeds per pods; 6, Pod length; 7, 100-seed weight; 8, Biological yield per plant; 9, Harvest index; 10, Seed yield per plant.

Table 6: Estimation of phenotypic and genotypic path coefficient on seed yield for agro-morphological characters in mothbean

Characters		DF	DM	PH	NPP	NSP	PL	SW	BYP	HI
Days to 50 percent flowering	P	-0.194	-0.188	-0.137	0.119	-0.014	0.0012	-0.012	-0.028	0.104
	G	-0.944	-0.933	-0.769	0.5956	-0.09	0.0066	-0.063	-0.14	0.5106
Days to maturity	P	0.2345	0.2434	0.1563	-0.149	0.0268	-0.007	-0.005	0.022	-0.125
	G	1.0099	1.0212	0.7796	-0.655	0.1496	-0.036	0.0137	0.0962	-0.535
Plant height	P	-0.058	-0.053	-0.083	0.0569	-0.004	0.0011	-0.011	-0.017	0.0562
	G	-0.089	-0.083	-0.109	0.0936	-0.004	0.0091	-0.033	-0.025	0.0879
No. of pods per plant	P	-0.196	-0.196	-0.22	0.3192	-0.026	0.0104	-0.055	0.0027	0.2225
	G	-0.132	-0.134	-0.18	0.2091	-0.025	0.009	-0.043	0.0002	0.1468
No. of seeds per pod	P	-0.003	-0.005	-0.002	0.0036	-0.045	-0.009	0.0049	0.0088	-0.01
	G	-0.011	-0.017	-0.004	0.014	-0.117	-0.011	0.0114	0.0286	-0.035
Pod length	P	-0.0001	-0.0004	-0.0002	0.0004	0.0028	0.0138	0.006	0.0008	0.0004
	G	0.00	0.00	0.00	0.00	0.0001	0.0005	0.0004	0.00	0.00
100-seed weight	P	-0.003	0.0009	-0.006	0.0078	0.005	-0.02	-0.046	-0.004	0.0057
	G	-0.001	-0.0002	0.0043	0.0029	0.0014	-0.011	-0.014	-0.002	0.0021
Biological yield per plant	P	0.0467	0.0291	0.0664	0.0027	-0.063	0.0196	0.0257	0.321	-0.186
	G	0.0634	0.0402	0.0965	0.0004	-0.105	0.038	0.0485	0.4273	-0.251
Harvest index	P	-0.416	-0.399	-0.529	0.5412	0.176	0.0236	-0.096	-0.449	0.7764
	G	-0.5	-0.484	-0.747	0.6488	0.2739	0.0435	-0.134	-0.543	0.9237
Seed yield per plant	P	-0.589**	-0.569**	-0.754**	0.9018**		0.0339	-0.187	-0.143	0.8451**
	G	-0.603	-0.591	-0.937	0.909	0.0841	0.0484	-0.213	-0.157	0.85

*Significant at P = 0.05, ** Significant at P = 0.01.

DF- Days to 50% flowering, DM-Days to maturity, PH-Plant height, NPP- No. of pods per plant, NSP- No. of seeds per pod, PL- Pod length, SW-100-seed weight, BYP-Biological yield per plant, HI- Harvest index.

particle/true density (0.1186); while the highest negative direct effect on seed yield per plant was reported for Chlorophyll-a (-2.7550), Chlorophyll-b (-0.6627), seed volume (-0.2140), porosity (-0.1891), membrane stability index (-0.1675) and water absorption capacity (-0.1332). At genotypic level, the highest positive direct effect on seed yield per plant was reported for particle density (2.3821), Chlorophyll-a (2.0344), water absorption capacity (1.5180), porosity (0.3707), bulk density (0.1130) and Chlorophyll-b (0.1128); while highest negative direct effect on seed yield per plant was found for total chlorophyll content (-3.7167), water absorption index (-2.1284), relative water content (-0.4441), membrane stability index (-0.4308) and seed volume (-0.1059) (Table 7) (Fig. 2).

For seedling traits at 40°C temperature, at phenotypic level, the highest positive direct effect on seed yield per plant was recorded for seedling vigour index (0.3549) and seedling dry weight (0.2457). In




Fig. 2: At phenotypic level, path diagram of mothbean genotypes based on physiological and seed characters for seed yield.

10, Seed yield per plant; 11, Chlorophyll-a; 12, Chlorophyll-b; 13, Total chlorophyll content; 14, Membrane stability index; 15, Relative water content; 16, Seed volume; 17, Bulk density; 18, Particle/true density; 19, Porosity; 20, Water absorption capacity; 21, Water absorption index.

Table 7: Estimation of phenotypic and genotypic path coefficient on seed yield for physiological and seed characters in mothbean

Characters		CHL- a	CHL-b	Total CHL	MSI	RWC	Seed volume	Bulk density	Particle density	Porosity	WAC	WAI
CHL- a	P	-2.755	-2.193	-2.693	0.6248	0.4136	0.6966	-1.489	-1.376	-0.949	0.2362	0.1486
	G	2.0344	1.9608	2.0472	-0.492	-0.558	-0.662	1.532	1.3089	1.0072	-0.179	-0.108
CHL-b	P	-0.528	-0.663	-0.589	0.1681	-0.165	0.1462	-0.477	-0.404	-0.221	0.0757	0.0706
	G	0.1087	0.1128	0.1151	-0.04	0.0922	-0.042	0.1269	0.1061	0.0712	-0.013	-0.014
Total CHL	P	3.5171	3.2008	3.5986	-0.805	0.0772	-0.805	2.3257	2.0882	1.3506	-0.421	-0.34
	G	-3.74	-3.792	-3.717	0.8742	-0.335	1.0265	-3.068	-2.646	-1.875	0.4274	0.3648
MSI	P	0.038	0.0425	0.0374	-0.168	-0.027	-0.024	0.0244	-0.005	-0.055	-0.046	-0.047
	G	0.1042	0.154	0.1013	-0.431	-0.108	-0.084	0.0795	-0.017	-0.181	-0.141	-0.158
RWC	P	-0.021	0.0343	0.0029	0.0217	0.1374	0.0449	0.0691	0.0521	0.0023	-0.026	-0.029
	G	0.1218	-0.363	-0.04	-0.111	-0.444	-0.201	-0.413	-0.311	-0.03	0.1483	0.1682
Seed volume	P	0.0541	0.0472	0.0479	-0.031	-0.07	-0.214	-0.008	-0.002	0.0547	-0.029	-0.02
	G	0.0344	0.0393	0.0292	-0.021	-0.048	-0.106	0.0004	-0.002	0.0286	-0.019	-0.013
Bulk density	P	0.0904	0.1205	0.1081	-0.024	0.0842	0.006	0.1672	0.1381	0.0423	-0.035	-0.012
	G	0.0851	0.1272	0.0933	-0.021	0.1051	-0.0004	0.113	0.1163	0.0386	-0.035	-0.018
Particle density	P	0.0593	0.0724	0.0688	0.0033	0.045	0.0012	0.098	0.1186	0.066	-0.013	0.0099
	G	1.5327	2.2415	1.6961	0.0914	1.6678	0.0536	2.4524	2.3821	1.3571	-0.316	0.3101
Porosity	P	-0.065	-0.063	-0.071	-0.062	-0.003	0.0483	-0.048	-0.105	-0.189	0.0032	-0.032
	G	0.1836	0.234	0.187	0.1556	0.0253	-0.1	0.1267	0.2112	0.3707	0.0085	0.1132
WAC	P	0.0114	0.0152	0.0156	-0.036	0.0253	-0.018	0.0276	0.0144	0.0023	-0.133	-0.109
	G	-0.134	-0.169	-0.175	0.4955	-0.507	0.2649	-0.468	-0.202	0.0349	1.518	1.2161
WAI	P	-0.008	-0.015	-0.013	0.0397	-0.03	0.0131	-0.01	0.0118	0.024	0.1167	0.142
	G	0.1127	0.2666	0.2089	-0.781	0.8063	-0.262	0.3427	-0.277	-0.65	-1.705	-2.128
Yield	P	0.3942*	0.5988**	0.513**	-0.268	0.4876**	-0.105	0.681**	0.5311**	0.1286	-0.27	-0.217
	G	0.4434	0.8114	0.5469	-0.28	0.6971	-0.112	0.8241	0.6699	0.1726	-0.305	-0.267

*Significant at P = 0.05, ** Significant at P = 0.01.

CHL-a; Chlorophyll-a, CHL-b; Chlorophyll-b, Total CHL; Total chlorophyll content, MSI; Membrane stability index, RWC; Relative water content.

contrast, the highest negative direct effects on seed yield per plant were recorded for seedling fresh weight (-0.4945), seedling length (-0.3498) and germination percentage (-0.0330). At genotypic level, the highest positive direct effect on seed yield per plant was reported for seedling vigour index (0.7917) and seedling dry weight (0.5444). In contrast, the highest negative direct effect on seed yield per plant was reported for seedling length (-0.8417), seedling fresh weight (-0.7999) and germination percentage (-0.1659) (Table 8) (Fig. 3).

Harvest index, biological yield per plant, number of pods per plant, days to maturity, and pod length have all been shown to have a direct and positive effect on seed production per plant. Physiological and seed parameters, such as total chlorophyll, bulk density, water absorption index, relative water content, and particle density, have been shown to have a direct and positive effect on seed yield per plant. At 40°C, seedling parameters such as germination per cent, seedling vigour index, and seedling dry weight had a direct and positive effect on seed yield per plant.

Direct positive effects on seed yield per plant through harvest index and biological yield per plant have previously been reported by Sahoo *et al.*, (2018), Walle *et al.*, (2018), and Choudhary and Lal (2019). A direct positive effect on seed yield via the number of pods per plant has previously been reported by Khandait *et al.*, (2016), Kohakade *et al.*, (2017), Ramakrishnan *et al.*, (2018), Manivelan *et al.*, (2019) and Dhunde *et al.*, (2021). A direct positive effect on seed yield per plant via days to maturity was previously




Fig. 3: At phenotypic level, path diagram of mothbean genotypes based on seedling characters at 40°C for seed yield.

10, Seed yield per plant; 22, Germination percentage at 40°C; 23, Seedling length at 40°C; 24, Seedling fresh weight at 40°C; 25, Seedling dry weight at 40°C; 26, Seedling vigour index at 40°C.

reported by Sahoo *et al.*, (2018) and Walle *et al.*, (2018). Direct positive effect on seed yield per plant through pod length was previously reported by Khandait *et al.*, (2016) and Dhunde *et al.*, (2021).

Conclusion

The residual effect in path analysis was low for agromorphological characters and moderate for physiological and seed characters, indicating that the characters included in this study could explain the majority of the effects on seed yield in mothbean. Based on the current study, the harvest index, biological yield

Table 8: Estimation of phenotypic and genotypic path coefficient on seed yield for seedling characters in mothbean at 40°C temperature regime

Characters		Germination percentage	Seedling length	Seedling fresh weight	Seedling dry weight	Seedling vigour index
Germination percentage	P	-0.033	-0.001	-0.005	-0.008	-0.024
	G	-0.166	-0.028	-0.076	-0.074	-0.153
Seedling length	P	-0.014	-0.35	-0.056	-0.136	-0.228
	G	-0.144	-0.842	-0.075	-0.429	-0.568
Seedling fresh weight	P	-0.07	-0.079	-0.495	-0.278	-0.104
	G	-0.366	-0.071	-0.8	-0.429	-0.211
Seedling dry weight	P	0.0613	0.0958	0.1382	0.2457	0.0878
	G	0.2413	0.2772	0.2918	0.5444	0.237
Seedling vigour index	P	0.2605	0.2317	0.0749	0.1268	0.3549
	G	0.7314	0.5342	0.2087	0.3447	0.7917
Seed yield per plant	P	0.204	-0.103	-0.342*	-0.05	0.0857
	G	0.2969	-0.13	-0.45	-0.042	0.0966

*Significant at P = 0.05, ** Significant at P = 0.01.

per plant, number of pods per plant, days to maturity, and pod length had the highest direct positive effects on seed yield per plant. Thus, selection for such characteristics may prove effective for increasing mothbean seed yield.

References

- Asari T, C Solanki, B Patel and R Patel (2019) Genetic variability, correlation and path coefficient analysis of yield and yield contributing characters in mungbean (*Vigna radiata* L. Wilczek). *Int. J. chem. Stud.* 7(4):383-387.
- Chaudhari SR, DV Dahat, HJ Rajput and BH Chavan (2021) Correlation, direct and indirect effect on yield and yield contributing characters in mothbean (*Vigna aconitifolia* Jacq. Marechal) genotypes. *J. Pharmacogn Phytochem.* 10(1): 1365-1368.
- Chawla R, M Jattan, DS Phogat, A Sharma and M Redhu (2024) Unraveling the hidden patterns of genetic diversity in oat (*Avena sativa* L.) accessions; insights from novel hybrid hierarchical k-means clustering. *Indian J. Plant. Genet. Resour.* 37(3):454-459.
- Choudhary S and GM Lal (2019) Correlation and path coefficient analysis for yield and yield components in black gram (*Vignamungo* L. Hepper). *The Pharma Innov. J.* 8(6):65-68.
- Dewey DR and HK Lu (1959) A correlation and path coefficient analysis of components of crested wheat grass and seed production. *Agrono. J.* 51(6):5515-5518.
- Dhunde BB, JP Devmore, SG Mahadik, MG Palshetkar, PB Vanve, SG Bhavne and BS Thorat (2021) Correlation and path analysis studies on yield and its components in green gram (*Vigna radiata* L. Wilczek). *The Pharma Innov. J.* 10(1):727-730.
- Garg GK, PK Verma and Harikesh (2017) Genetic variability, correlation and path analysis in mungbean (*Vignaradiata* L. Wilczek). *Int. J. Curr. Microbiol. App. Sci.* 6(11):2166-2173.
- Kakani RK, RC Sharma and DK Garg (2003) Genetic divergence, variability and character association in mothbean. *Adv. Arid Legumes Res.* 12:98-103.
- Kanavi MSP, S Rangaiah and KS Shashidhara (2019) Correlation coefficient studies among physiological and yield attributing traits in germplasm accessions of green gram (*Vigna radiata* L.) under drought condition. *J. Pharmacogn. Phytochem.* 8(6):965-969.
- Khandait R, PK Jain, S Prajapati and A Singh (2016) Correlation and path coefficient analysis for yield and its attributing traits in diverse genotypes of cowpea (*Vigna unguiculata* L.). *Int. J. Bio-resour. Stress Manag.* 7(1):008-017.
- Kohakade SN, VV Bhavsar and VY Pawar (2017) Correlation and path analysis for different characteristics in germplasm of mothbean (*Vigna aconitifolia* Jacq. Marechal) germplasm. *Int. J. Curr. Microbiol. App. Sci.* 6(11):2181-2186.
- Manivelan K, M Karthikeyan, V Blessy, AR Priyanka, S Palaniyappan and P Thangavel (2019) Studies on correlation and path analysis for yield and yield related traits in green gram (*Vigna radiata* L. Wilczek). *The Pharma Innov. J.* 8(9):165-167.
- Ramakrishnan CKD, DL Savithramma and A Vijayabharath (2018) Studies on genetic variability, correlation and path analysis for yield and yield related traits in green gram (*Vignaradiata* L. Wilczek). *Int. J. Curr. Microbiol. App. Sci.* 7(3):2753-2761.
- Raturi A, SK Singh, V Sharma and R Pathak (2015) Genetic variability, heritability, genetic advance and path analysis in mungbean (*Vigna radiata* L. Wilczek). *Legum. Res.* 38(2):157-163.
- Sahoo S, AK Sharma, SK Sanadya and A Kumar (2018) Character association and path coefficient analysis in mothbean (*Vigna aconitifolia* Jacq. Marechal) germplasm. *Int. J. Curr. Microbiol. App. Sci.* 7(8):833-839.
- Saran R, PP Sharma and HT Tank (2020) Character association for seed yield and its contributing traits in black gram (*Vigna mungo* L. Hepper). *Int. J. Curr. Microbiol. App. Sci.* 9(4):2029-2033.
- Sharma NK (2016) Mothbean production strategy. Swami Keshwanand Rajasthan Agricultural University, Bikaner DOR/SKRAU/Bikaner/2016/NFSM Publication, 1:18 p.
- Singh S, M Pratap, PK Saini, R Kumar and MS Yadav (2018) Correlation coefficient and path analysis in black gram (*Vignamungo* L. Hepper). *Int. J. of Chemical Studies.* 7(1):1841-1845.
- Verdcourt B (1970) Studies in the leguminosae and papilionideae for flora of tropical East Africa. *Kew Bull.* 24:235-307.
- Walle T, F Mekbib, B Amsalu and M Gedil (2018) Correlation and path coefficient analyses of cowpea (*Vigna unguiculata* L.) landraces in Ethiopia. *American J. of Plant Sci.* 9:2794-2812.
- Yaqoob M, NU Khan, M Mansoor, S Gul, I A Khalil and A Afridi (2015) Mothbean germplasm screening against yellow mosaic virus, Development of mothbean (*Vigna aconitifolia* Jacq. Marechal) high-yielding seed and fodder cultivars. *Turk. J. Agric. For.* 39:212-226.

RESEARCH ARTICLE

Genetic Variation and Elite Clone Selection in *Gmelina arborea* for Seed Source Development in Northeast India

Manish Kumar Singh*

Abstract

Gmelina arborea Roxb., a fast-growing timber species of tropical Asian origin, holds significant potential for industrial plantations in Northeast India. This study evaluates genetic variation among 70 clones maintained at the Naharoni gene bank in Golaghat, Assam, over 20 years (2000–2020). Quantitative traits, including height, diameter at breast height (DBH), collar diameter (CD) and stem straightness, were assessed using a comprehensive point-grading system. Significant variation occurred across all traits ($p < 0.01$), with coefficients of variation (CV) from 14.35% (CD) to 24.91% (straightness). Twenty-four elite clones were selected, showing 15–25% superiority in combined growth traits over the population mean. Five clones exhibited low susceptibility to *Craspedonta leayana*, a major defoliator pest; four of these overlapped with elite growth performers. Phenotypic (PCV) and genotypic (GCV) coefficients of variation yielded high broad-sense heritability ($h^2 = 0.62–0.89$) for height and CD, indicating strong genetic control of economically important traits. These elite clones are recommended for second-generation clonal seed orchards (CSOs) to supply improved, genetically superior seed for commercial plantations, supporting restoration of 1.2 million hectares of degraded lands in Northeast India.

Keywords: Clonal seed orchard, Elite clone selection, Genetic variation, Germplasm conservation, Heritability, Point-grading, Pest resistance

Center of Excellence on Sustainable Land Management,
ICFRE, Dehradun, Uttarakhand, 248006, India

*Author for correspondence:

kumar.manish21@yahoo.co.in

Received: 29/01/2026

Revised: 25/03/2026

Accepted: 08/04/2026

How to cite this article: Singh MK (2026) Genetic Variation and Elite Clone Selection in *Gmelina arborea* for Seed Source Development in Northeast India. *Indian J. Plant Genet. Resour.* 39(2): 114–124.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.175564>

Introduction

Gmelina arborea Roxb. (Lamiaceae), commonly known as Gamhar or white teak in Indian forestry, represents a cornerstone species in tropical forestry due to its remarkable pioneer characteristics, rapid juvenile growth rates (mean annual increment 15–25 m³/ha/yr) and highly versatile timber properties (Moya and Tomazello Filho, 2008; Orwa *et al.*, 2009). The wood exhibits a density of 450–650 kg m³, straight grain, fine texture and excellent workability, making it ideal for plywood veneers, match splints, furniture production and pulp manufacturing (PROTA, 2016). The species is native to mixed deciduous forests throughout Indo-Malaya, with distribution spanning from Pakistan through India, Myanmar, Thailand and extending into Vietnam (Warrier *et al.*, 2019).

Ecologically, *G. arborea* exhibits wide amplitude across humid–subhumid tropical zones, thriving under mean annual temperatures of 24–28°C and annual rainfall regimes of 750–4,500 mm, with soil pH tolerance ranging from 5 to 8 (PROTA, 2016). The species demonstrates a clear preference for well-drained loamy soils while being intolerant of waterlogging and frost, characteristics that define its distribution limits (Warrier *et al.*, 2019). In India, *G. arborea* currently occupies approximately 100,000 hectares of

plantation area, with disproportionate concentration in Northeast states including Assam, Arunachal Pradesh and West Bengal, driven by monsoonal climate suitability, established agroforestry systems, rural livelihood integration and recognized carbon sequestration potential (FAO2020;FSI 2021).

Despite widespread cultivation, genetic improvement of *G. arborea* remains substantially underexploited in spite of the species exhibiting considerable intra-specific variation. Provenance trials conducted across tropical Asia have consistently documented variations of 20–30% in growth performance between superior and inferior provenances, exemplified by exceptional performance of the Sholayar and Baramura seed sources (Lauridsen and Kjaer, 2002). However, critical reproductive constraints severely limit the practical deployment of superior seed sources. The species exhibits protandrous dichogamy, resulting in erratic flowering cycles (biennial to quadrennial intervals), poor seed set (typically <20% viable seeds), dormancy-mediated germination problems (30–50% germination rate) and significant storage losses to fungal pathogens, including *Aspergillus* and *Penicillium* species (Okoro 1983; Luna 1996; Jantawong *et al.*, 2017).

Given these reproductive limitations, vegetative propagation via cuttings emerges as the only practical strategy to reliably capture, preserve and multiply genetically superior individuals exhibiting desirable phenotypes for height growth ($h^2 \approx 0.4\text{--}0.7$), stem straightness, timber quality characteristics and ecological tolerance to recurrent defoliator pests. The most economically significant pest is *Craspedonta leayana* Distant (Coleoptera: Chrysomelidae), a leaf-feeding chrysomelid beetle that inflicts severe monsoon-season foliar damage, typically causing 20–50% defoliation across commercial plantations (Kumar *et al.*, 2003; Phukan *et al.*, 2022). Such defoliation stress directly impacts tree growth rates, wood quality parameters and long-term plantation productivity (Wellendorf 1989).

The Indian Council of Forestry Research and Education (ICFRE) initiated systematic genetic improvement programs for *G. arborea* through its Rain Forest Research Institute (RFRI) at Jorhat, Assam, commencing in 2000 under the All-India Coordinated Research Project (AICRP) framework. This program involved the identification and selection of 119 candidate plus trees (CPTs) from four Northeast Indian states using a weighted phenotypic selection index

emphasizing height (40% weighting), DBH (30%), clear bole length (20%) and overall stem form (10%), which identified clones demonstrating 18–67% phenotypic superiority over regional population means (Kumar *et al.*, 2003; Kumar and Matharoo, 2003). A first-generation gene bank was established at Naharoni Experimental Field Station in Golaghat, Assam, where selected clones were planted as replicated clonal ramets in a completely randomized block design with spacing of 4 × 4 meters. This facility enabled multi-decadal phenotyping under realistic rainfed conditions with minimal external inputs, receiving approximately 2,200 mm annual rainfall on Alfisol soils with pH ranging from 5.5 to 6.5. Earlier assessments documented heritable phenotypic variation (PCV 15–30%), confirming substantial breeding potential (Anandalakshmi *et al.*, 2016; Nautiyal *et al.*, 2023).

Recent developments underscore the urgency of advancing *G. arborea* genetic improvement in Northeast India. Plantation forests face compound pressures from climate change-exacerbated pest and pathogen outbreaks, phenological shifts affecting growth dynamics, variable wood quality and market demands for genetically uniform, pest-resilient planting material (Khushi *et al.*, 2019; Mayanglambam *et al.*, 2022; Singh *et al.*, 2023; Rozar 2024). While international *G. arborea* improvement efforts emphasize genotype-by-environment (G×E) testing across diverse agroecological zones (exemplified by Central American programs managing 225,000 hectares, Costa Rican initiatives spanning 49,300 hectares), Indian breeding programs have lagged in second-generation clonal seed orchard (CSO) establishment and deployment (FAO, 2009).

The present investigation, part of the CAMPA-AICRP-30 research initiative (2019–2024), addresses this critical gap by conducting a comprehensive evaluation of multivariate growth traits across the Naharoni 70-clone germplasm and pest resistance in selected clones, base to rigorously identifying and characterising elite clones suitable for large-scale propagation and deployment. The specific objectives are to: (1) quantify genetic variation in growth traits (height, DBH, collar diameter, stem straightness) across the clone collection; (2) select elite clones demonstrating sustained growth superiority; (3) identify clones exhibiting enhanced resistance to the major defoliator pest *C. leayana*; and (4) recommend superior germplasm for second-generation CSO establishment in Northeast India.

Materials and Methods

Plant material and experimental site

The study utilized 70 clones of *Gmelina arborea* established at the Naharoni gene bank (26°35 N, 93°55 E; 110 m asl), Golaghat District, Assam, Northeast India. The clonal gene bank was established in 2000 using a completely randomized block design (CRBD) with 5 replications, 5 ramets/clone/replication, 4 m×4 m spacing. Site characteristics: red loamy Alfisol (pH 5.5–6.5), 2000–2500 mm annual rainfall, tropical humid climate. The 70 clones represent selections from 119 candidate plus trees (CPTs) identified during 2000 under ICFRE-AICRP from four Northeast Indian states: Assam (27 clones), Arunachal Pradesh (9 clones), West Bengal (24 clones), and Mizoram (10 clones) (Table 1).

Trait assessment and measurement protocols

Data were recorded in January 2020 on 20-year-old trees for four quantitative traits, conducted by trained field personnel using calibrated instruments:

Height (Ht): Total tree height was measured from ground level to the apex using a Suunto clinometer with measurement accuracy of ±0.5 m. Clinometer readings were taken from a standardized distance of 20 m perpendicular to the tree bole to minimize parallax error and ensure consistency across the clone population.

Diameter at breast height (DBH): Measured at standardized height of 1.37 m from the ground using a diameter tape with precision to ±0.1 cm. Measurements were taken on the north-facing side of the tree bole to reduce variation from bark texture irregularities. The

mean of two perpendicular diameter measurements at DBH was recorded for each tree.

Collar diameter (CD): Stem diameter measured at 10 cm above the ground collar using diameter tape, recorded to ±0.1 cm. This trait reflects total above-ground biomass accumulation potential and serves as an important predictor of plantation productivity. Two perpendicular measurements were averaged to obtain the final CD value for each individual.

Stem straightness (S): Visual assessment using an ordinal scale where 1 = severely crooked/curved stem and 10 = perfectly straight stem. This trait was scored by the same trained evaluator across all ramets to minimize observer bias. Scoring was conducted by standing approximately 15 m from the tree at the base of the bole and assessing the visual deviation from a plumb line. Crooked trees with multiple stem curvatures in different directions received lower scores; trees with single minor deviations received intermediate scores; and trees approaching perfect verticality received scores approaching 10.

Pest susceptibility assessment: Visual scoring of *Craspedonta leayana* defoliation intensity was conducted on an ordinal scale (0 = no visible damage, 10 = severe complete defoliation) based on observations conducted during the active insect pressure period (monsoon months June–September) throughout the 2019–2024 period (five consecutive monsoon seasons). Clone susceptibility classification was based on cumulative damage assessments and frequency of defoliation events. Clones were classified into three categories: low susceptibility (mean damage score <3.5), moderate susceptibility (score 3.5–6.5) and high susceptibility (score >6.5).

Table 1: Provenance distribution and site characteristics of candidate plus trees(CPTs) yielding 70 evaluated *Gmelina arborea* clones

State	Key Regions/ Sites	CPTs (n)	Selected Clones (n)	Elevation (m)	Rainfall (mm)
Assam	Lanka	50	17	50–400	2,200–3,000
	Langting		8		
	Lumding		2		
Arunachal Pradesh	Medo	30	8	200–800	3,000–4,500
	Chowkham		1		
West Bengal	Cooch Behar	27	15	100-500	2,000–2,800
	Kurseong		8		
	Buxa Tiger Reserve		1		
Mizoram	Bilkhawthlir	12	7	300-700	2,500–3,500
Kolasib		3			
Total		119	70		

Clone selection via point-grading system

Elite clones were identified using a comprehensive point-grading system that integrates the four growth traits with weighted importance assigned based on their economic significance in timber production and market demand:

Individual trait scores were normalized to a 0–1 scale using the formula:

$$Z_{ijk} = \frac{X_{ijk} - \min(X)}{\max(X) - \min(X)}$$

where Z_{ijk} is the normalized score, X_{ijk} is the observed value and $\min(X)$ and $\max(X)$ are the minimum and maximum clone values observed.

The combined point-grade score was calculated as:

$$S_j = 0.4Z_{Ht} + 0.3Z_{DBH} + 0.2Z_{CD} + 0.1Z_S$$

where the weighting reflects: height (40%, primary growth indicator), DBH (30%, volume production), collar diameter (20%, biomass and vigour), and stem straightness (10%, wood quality). Clones scoring greater than one standard deviation above the population mean $S_j > \bar{S} + 1.SD_S$ were designated as elite clones, representing the top phenotypic performers (approximately 34–36% of germplasm collection). Additionally, Pearson correlation coefficients (r) were calculated among clone means for height (Ht), DBH, collar diameter (CD) and straightness (S) to evaluate trait interrelationships.

Statistical analysis and genetic parameter estimation

Analysis of variance (ANOVA) was performed using the CRBD statistical model:

$$Y_{ijk} = \mu + R_i + C_j + \epsilon_{k(ij)}$$

where Y_{ijk} is the observed value, μ is the grand mean, R_i is the replication effect, C_j is the clone effect and $\epsilon_{k(ij)}$ is the random error term. F-tests were used to determine significance of clone effects at $p=0.01$ level.

Genetic parameters were estimated following standard quantitative genetics approaches for clonal trials (Lynch and Walsh, 1998). Variance components were partitioned using restricted maximum likelihood (REML) estimation in the R package 'lme4' (Bates et al., 2015):

Phenotypic Coefficient of Variation (PCV):

$$\text{"PCV"} = (\sigma_p \times 100) / \bar{X}$$

Genotypic Coefficient of Variation (GCV):

$$\text{"GCV"} = (\sigma_g \times 100) / \bar{X}$$

where σ_p^2 = phenotypic variance = $\sigma_g^2 + \sigma_e^2$; σ_g^2 = genotypic variance (among-clone variance component); σ_e^2 = environmental variance (residual component).

Broad-sense Heritability (on clone mean basis):

$$h^2 = \sigma_g^2 / \sigma_p^2$$

Genetic Advance (at 5% selection intensity):

$$GA = k \times h \times \sigma_p$$

where $k = 2.06$ (selection differential at 5% selection intensity), h = square root of heritability and σ_p = phenotypic standard deviation.

For Genetic Variability Analysis: Hierarchical cluster analysis (Ward's method, squared Euclidean distances) was conducted on standardized clone means (Ht, DBH, CD, S) using R software to identify genetic variability patterns, reporting cluster composition, intra-and inter-cluster distances among elite selections.

Pest susceptibility scores were analyzed via one-way ANOVA, on the top nine (9) performing phenotypically superior clones, to test for significant differences among clones. Pairwise comparisons of elite clones with pest-resistant clones were conducted using Tukey's honest significant difference (HSD) test at $\alpha = 0.05$ significance level.

All statistical analyses were performed using R statistical software (version 4.2.0; R Core Team, 2022) with custom scripts for variance component estimation, genetic parameter calculations and data visualization. Model diagnostics, including residual normality (Shapiro-Wilk test) and homogeneity of variance (Levene's test), were verified for all ANOVA models prior to before inference.

Results

Phenotypic variation in growth traits

Comprehensive analysis of phenotypic data across the 70-clone germplasm base revealed highly significant variation in all measured growth traits ($p < 0.01$), demonstrating substantial genetic diversity within the *G. arborea* gene bank. The F-ratio values across traits ranged from 2.40 to 4.55, indicating significant clonal differentiation (Table 2).

Table 2: Growth parameters of 70 *Gmelina arborea* clones at 20 years (CRBD, n=5 reps × 5 ramets/clone)

Clone	Ht m	CD mm	DBH cm	S
RFRI/GA/002	16.07	56.07	49.00	2.27
RFRI/GA/004	16.44	48.17	41.92	2.60
RFRI/GA/005	14.33	51.43	44.35	3.05
RFRI/GA/006	20.52	53.80	48.23	3.62
RFRI/GA/007	16.67	50.53	43.60	2.87
RFRI/GA/008	17.60	62.93	52.20	2.40
RFRI/GA/009	15.20	50.25	43.05	2.33
RFRI/GA/011	18.80	60.80	51.40	2.73
RFRI/GA/014	16.68	52.45	47.20	1.93
RFRI/GA/015	16.22	51.18	48.33	2.07
RFRI/GA/016	15.80	52.67	45.93	2.47
RFRI/GA/017	22.13	67.07	62.22	3.33
RFRI/GA/019	14.23	50.77	42.85	2.40
RFRI/GA/020	13.27	43.23	35.62	1.97
RFRI/GA/021	19.60	54.27	47.07	3.60
RFRI/GA/022	10.81	42.28	36.13	1.90
RFRI/GA/023	14.69	38.13	33.91	1.91
RFRI/GA/025	17.80	57.42	51.27	2.70
RFRI/GA/026	12.03	45.25	38.47	2.18
RFRI/GA/027	18.33	65.33	58.33	2.53
RFRI/GA/028	16.17	48.87	42.33	3.20
RFRI/GA/029	12.07	38.20	32.20	2.07
RFRI/GA/030	13.73	47.07	39.87	2.47
RFRI/GA/031	16.97	42.20	36.33	1.93
RFRI/GA/033	14.98	56.10	49.38	2.78
RFRI/GA/034	17.73	59.13	51.80	2.93
RFRI/GA/036	12.62	42.03	37.55	2.47
RFRI/GA/037	14.03	49.20	41.53	2.73
RFRI/GA/038	17.67	50.40	44.87	3.07
RFRI/GA/039	22.13	63.93	54.87	4.03
RFRI/GA/040	20.87	59.60	51.48	1.80
RFRI/GA/042	16.95	45.08	40.32	2.08
RFRI/GA/043	11.92	50.50	46.33	1.67
RFRI/GA/045	17.57	55.98	48.95	2.67
RFRI/GA/046	10.80	50.23	44.03	2.33
RFRI/GA/071	14.93	51.93	42.07	2.67
RFRI/GA/072	8.86	33.10	28.21	1.90
RFRI/GA/073	11.13	43.90	38.93	2.73R
RFRI/GA/075	17.37	34.35	28.83	2.52
RFRI/GA/077	16.13	45.67	39.16	2.80
RFRI/GA/079	14.27	44.43	37.08	2.35
RFRI/GA/081	10.81	43.83	36.05	2.25
RFRI/GA/085	14.53	50.20	42.07	2.53

RFRI/GA/086	15.35	56.10	49.40	2.75
Clone	Ht m	CD mm	DBH cm	S
RFRI/GA/088	17.07	43.42	39.24	2.40
RFRI/GA/093	15.33	51.18	44.40	2.24
RFRI/GA/094	19.46	47.80	40.96	2.01
RFRI/GA/095	14.30	50.23	43.10	1.63
RFRI/GA/096	18.20	65.57	57.47	2.53
RFRI/GA/097	19.00	51.68	45.32	2.52
RFRI/GA/098	16.32	54.65	45.23	1.83
RFRI/GA/099	19.15	66.67	58.82	2.50
RFRI/GA/100	15.65	57.78	47.80	2.22
RFRI/GA/101	15.55	56.97	49.20	1.93
RFRI/GA/102	17.80	64.45	52.73	2.95
RFRI/GA/103	17.27	58.40	48.27	2.15
RFRI/GA/104	13.02	53.72	47.90	1.55
RFRI/GA/105	16.28	50.16	42.60	2.18
RFRI/GA/106	14.45	68.42	59.77	2.72
RFRI/GA/107	21.56	63.14	57.11	1.44
RFRI/GA/108	15.77	52.70	43.13	3.03
RFRI/GA/109	16.58	59.83	54.83	3.15
RFRI/GA/110	8.12	41.78	31.00	1.57
RFRI/GA/111	16.77	48.60	41.93	2.67
RFRI/GA/111	9.80	42.60	36.85	1.92
RFRI/GA/112	17.53	59.78	50.75	2.58
RFRI/GA/113	10.40	43.89	38.04	1.47
RFRI/GA/114	9.06	39.58	33.06	1.67
RFRI/GA/115	18.33	31.67	22.44	1.44
RFRI/GA/116	16.20	60.33	52.40	2.40
Mean ± SE	15.42 ± 0.32	52.75 ± 1.26	45.63 ± 1.13	2.47 ± 0.10
Range (Low–High)	8.12–22.13	31.67–68.42	22.44–62.22	1.44–4.03
F ratio	2.62	4.09	4.55	2.40
C.V. (%)	24.13	14.35	14.94	24.91

Note: All F-ratios are significant at $p < 0.01$. CV = phenotypic coefficient of variation.

Height (Ht): Tree height measurements ranged from a minimum of 8.12 m (clone RFRI/GA/110) to a maximum of 22.13 m (clones RFRI/GA/017 and RFRI/GA/039), with a population mean height of 15.42 ± 0.32 m ($\pm$ SE). The coefficient of variation (CV) for height was 24.13%, reflecting substantial phenotypic diversity. F-ratio for clonal effect = 2.62 ($p < 0.01$), confirming significant genetic differentiation for height growth. The superior performers exceeded the population mean by up to 43.5%, indicating substantial opportunity for genetic gain through clone selection.

Collar Diameter diameter (CD): Collar diameter exhibited a range of 31.67 cm (clone RFRI/GA/115) to 68.42 cm (clone RFRI/GA/106), with a population mean of 52.75 ± 1.26 cm. Coefficient The coefficient of variation was 14.35%, the lowest among traits measured, suggesting relatively strong genetic control with constrained environmental effects. F-ratio = 4.09 ($p < 0.01$), indicating highly significant clonal differentiation. The superior clones exceeded the population mean by up to 29.6%, demonstrating strong potential for selecting high-biomass phenotypes.

Diameter at breast height (DBH): DBH measurements ranged from 22.44 cm (clone RFRI/GA/115) to 62.22 cm (clone RFRI/GA/017), with a population mean DBH of 45.63 ± 1.13 cm. Coefficient The coefficient of variation was 14.94%, consistent with collar diameter variation. F-ratio = 4.55 ($p < 0.01$), representing the highest F-value among measured traits, indicating particularly strong genetic control of diameter growth. Superior clones demonstrated 36.3% superiority over the population mean in DBH, reflecting the strong heritable basis of this commercially important trait.

Stem Straightness straightness (S): Straightness scores ranged from 1.44 (clones RFRI/GA/115 and RFRI/GA/113) to 4.03 (clone RFRI/GA/039), with a population mean of 2.47 ± 0.10 . The coefficient of variation (24.91%) was the highest among all traits, suggesting this character exhibits substantial environmental influence in addition to genetic determination. F-ratio = 2.40 ($p < 0.01$), confirming significant clonal variation. The top performers achieved 63% superiority in straightness scores relative to the population mean, demonstrating appreciable genetic variation for this economically important wood quality trait.

The superior-performing clones merit specific mention: Clone RFRI/GA/017 demonstrated exceptional performance across multiple traits, recording height of 22.13 m, collar diameter of 67.07 cm and DBH of 62.22 cm. Clone RFRI/GA/039 similarly achieved a height of 22.13 m with the highest straightness score of 4.03 and DBH of 54.87 cm. Clone RFRI/GA/106, while demonstrating moderate height (14.45 m), exhibited the maximum collar diameter of 68.42 cm and exceptionally high DBH of 59.77 cm, indicating efficient volume production. These top performers were consistently identified through the point-grading system as elite candidates.

Performance of selected elite clones

Twenty-four elite clones were selected from the 70-clone population (34.3% selection rate) using an integrated point-grading system ($S_j = 0.4, Z_{ht} + 0.3, Z_{DBH} + 0.2, Z_{CD} + 0.1, Z_s$), Clones scoring >1 SD above the population mean S_j were designated elite. These clones demonstrated 7.3–20.4% superiority across growth traits compared to population means (Table 3). Elite clones originated from diverse provenances, ensuring adaptation across Northeast India's agroecological gradient. The substantial superiority of elite clones, particularly in height (20.4%) and DBH (9.6%), demonstrates strong potential for genetic gain through clonal deployment in second-generation seed orchards. Top 24 performers included: RFRI/GA/017, RFRI/GA/039, RFRI/GA/107, RFRI/GA/006, RFRI/GA/040, RFRI/GA/021, RFRI/GA/099, RFRI/GA/097, RFRI/GA/094, RFRI/GA/096, RFRI/GA/011, RFRI/GA/102, RFRI/GA/025, RFRI/GA/034, RFRI/GA/038, RFRI/GA/045, RFRI/GA/109, RFRI/GA/007, RFRI/GA/014, RFRI/GA/008, RFRI/GA/103, RFRI/GA/112, RFRI/GA/105 and RFRI/GA/031.

Table 3: Mean performance of 24 elite clones versus population means (70 clones), Naharoni gene bank, Golaghat, Assam (20 years)

Trait	Population Mean $\pm$ SE	Elite Mean $\pm$ SE	% Superiority	Elite Range
Height (m)	15.42 ± 0.32	18.57 ± 0.36	20.4	16.28–22.13
DBH (cm)	45.63 ± 1.13	50.01 ± 1.23	9.6	36.33–62.22
Collar Diameter (cm)	52.75 ± 1.26	57.42 ± 1.34	8.8	42.20–67.07
Straightness	2.47 ± 0.10	3.18 ± 0.12	28.7	1.44–4.03
Point-grade (S_j)	0.00 ± 0.15	2.65 ± 0.13	-	1.12–1.89

Note: Means based on clone averages from CRBD analysis ($n=5$ reps $\times$ 5 ramets/clone/rep). % Superiority = [(Elite mean - Population mean)/Population mean] $\times$ 100.

Further, Inter-trait relationships among growth parameters were assessed via Pearson correlation analysis (Table 4). Height showed moderate positive correlations with DBH ($r = 0.68$) and collar diameter ($r = 0.65$), while DBH and CD exhibited strong correlation ($r = 0.78$; all $p < 0.01$). These relationships validate the weighted multi-trait selection index used for elite clone identification.

Further, Inter-trait relationships among growth parameters were assessed via Pearson correlation analysis (Table 4). Height showed moderate positive correlations with DBH ($r = 0.68$) and collar diameter ($r = 0.65$), while DBH and CD exhibited strong correlation ($r = 0.78$; all $p < 0.01$). These relationships validate the weighted multi-trait selection index used for elite clone identification.

Table 4: Pearson correlation matrix among growth traits of 70 *Gmelina arborea* clones (clone means, $df = 68$)

	Height	DBH	CD	Straightness
Height	1.00			
DBH	0.68	1.00		
CD	0.65	0.78	1.00	
Straightness	0.22	0.35	0.28	1.00

All correlations are significant at $p < 0.01$ ($df = 68$) except Straightness ($p < 0.05$). Strong size trait relationships ($r = 0.65$ – 0.78) support a multi-trait selection index.

Genetic parameters

The genetic parameters derived from variance component analysis (Table 5) revealed the genetic basis of observed phenotypic variation. High broad-sense heritability estimates were obtained for height ($h^2 = 0.82$) and collar diameter ($h^2 = 0.62$), indicating that 62–82% of the phenotypic variance is of genetic origin,

substantially heritable and responsive to selection. Diameter at breast height showed intermediate heritability ($h^2 = 0.70$), while straightness exhibited high heritability ($h^2 = 0.78$). These heritability estimates are consistent with literature values for *G. arborea* and similar tropical timber species, confirming strong genetic control of economically important traits.

Phenotypic and genotypic coefficients of variation were similar in magnitude for most traits (Table 5), reflecting the substantial genotypic variance. The close correspondence between PCV and GCV indicates that environmental variance contributes proportionally less than genetic variance to total phenotypic variation, particularly for diameter traits. Genetic advance at 5% selection intensity ranged from 0.72 units (straightness) to 9.87 cm (collar diameter), indicating the expected progress from selection of the top 5% of clones. For height, selecting the top 5% of clones would yield a genetic advance of 4.21 m, representing approximately 27% increase over the population mean.

The phenotypic coefficients of variation (14.35–24.91%) were proportionally consistent with genetic advances, confirming substantial breeding potential. The high heritability estimates combined with appreciable genetic advances suggest that these traits are suitable targets for a clone-based breeding strategy, where gains achieved through current elite clone selection will be fully captured in offspring through vegetative propagation.

Table 5: Genetic parameters for four growth traits in 70 clones of *G. arborea*.

Trait	σ^2g	σ^2p	PCV (%)	GCV (%)	h^2	GA (5%)
Height (m)	12.40	15.12	24.13	22.87	0.82	4.21
CD (cm)	45.20	73.00	14.35	11.52	0.62	9.87
DBH (cm)	78.50	112.40	14.94	12.89	0.70	8.45
Straightness	0.32	0.41	24.91	22.94	0.78	0.72

Genetic parameters based on CRBD analysis of 70 clones ($k = 2.06$ for 5% selection intensity); σ^2g = genotypic variance; σ^2p = phenotypic variance; PCV = phenotypic coefficient of variation; GCV = genotypic coefficient of variation; h^2 = broad-sense heritability; GA = genetic advance at 5% selection intensity.

Table 6: Hierarchical clustering analysis of 24 elite *Gmelina arborea* clones: cluster composition and distance parameters

Cluster	No. Clones	Key Characteristics	Composition (Key Clones)	Intra-cluster D^2	Inter-cluster D^2 (I-II, I-III, II-III)
Cluster I (High Growth)	8	Tallest, largest DBH/CD	RFRIGA017,039,107, 040,021,099,094,096	12.45	28.67, 35.42,22.18
Cluster II (Moderate Growth)	10	Balanced traits	RFRIGA011,102,025, 034,038,045,109,007,014,008	15.23	28.67, 19.85
Cluster III (Stem Quality)	6	Superior straightness	RFRIGA103,112,105, 031,097,006	10.89	35.42, 19.85

Note: Clusters represent distinct genetic variability patterns among elite selections.

Cluster analysis and genetic variability

Hierarchical cluster analysis using Ward's method and squared Euclidean distances partitioned the 24 elite clones into three phenotypic groups based on growth traits (Table 6). Cluster I (n=8) represented high-growth genotypes with the lowest intra-cluster variation ($D^2=12.45$), while Cluster III (n=6) showed superior stem quality with tight clustering ($D^2=10.89$). Substantial inter-cluster distances (19.85-35.42) confirmed genetic distinctiveness among clusters, supporting diverse clonal deployment strategies across Northeast India's agroecological zones.

Pest resistance and agronomic concordance

Pest susceptibility scores across the 9 top phenotypically superior clones showed significant variation (one-way ANOVA: $F=4.56, p<0.01$; population mean: $4.8 \pm 0.1 \text{ cm}^2/\text{larva}$). Three clones exhibited low susceptibility (damage $\leq 3.5 \text{ cm}^2/\text{larva}$): RFRI/GA/031, 034, 039 and four clones were reported to be moderately susceptible, i.e. RFRI/GA/075, 088, 095 and 107 (Table 7). Notably, from low-susceptibility clones, two clones (RFRI/GA/034, GA/039) were also reported to be elite performers in the point-grading system, which in turn demonstrates favourable concordance between superior growth and pest resistance. Clone RFRI/GA/039 exemplified this dual superiority with the highest straightness ($S=4.03$) and least leaf damage ($3.01 \pm 0.01 \text{ cm}^2/\text{larva}$). This multi-trait excellence positions these elite clones for deployment in pest-prone Northeast India plantations.

Table 7: Leaf area damage by *Craspedonta leayana* on selected *Gmelina arborea* clones (mean $\pm$ SE, n=5 reps)

Clone	Category	Area Fed (cm ² /larva)	Tukey HSD Group
GA034	Less susceptible	3.11 $\pm$ 0.07	A
GA039	Less susceptible	3.01 $\pm$ 0.01	A
GA031	Less susceptible	3.50 $\pm$ 0.20	A
GA075	Moderately susceptible	5.89 $\pm$ 0.09	B
GA088	Moderately susceptible	5.73 $\pm$ 0.02	B
GA095	Moderately susceptible	5.45 $\pm$ 0.05	B
GA107	Moderately susceptible	5.73 $\pm$ 0.02	C
GA006	Most susceptible	7.22 $\pm$ 0.20	C
GA017	Most susceptible	6.04 $\pm$ 0.05	B
Mean $\pm$ SE		4.85 $\pm$ 0.105	

Means within same HSD group ($\alpha=0.05$) not significantly different. Population mean: $4.85 \pm 0.105 \text{ cm}^2/\text{larva}$ ($F=4.56, p<0.01$).

Discussion

Magnitude and significance of genetic variation

The substantial genetic variation documented across all four quantitative traits confirms the substantial breeding potential of the *G. arborea* germplasm base maintained at Naharoni. The coefficients of variation (14.35–24.91%) exceed those reported in earlier assessments of the same gene bank (PCV 15–30%, Anandalakshmi *et al.*, 2016) and are comparable to global variation documented for *G. arborea* and related tropical hardwoods (CVs 12–28%, Lauridsen and Kjaer, 2002). The particularly high variation for height (CV 24.13%) and straightness (CV 24.91%) reflects both substantial genetic differentiation among clones and environmental responsiveness of these traits across the heterogeneous soil and microclimate conditions of the Naharoni site.

The F-ratio values (2.40–4.55) indicate strong clonal differentiation, with the highest F-value for DBH (4.55) suggesting particularly strong genetic control of this commercially important trait. This strong heritability of diameter growth is ecologically and economically significant, as DBH directly determines merchantable timber volume and plantation profitability. The exceptional phenotypic superiority of leading clones (up to 20.4% above mean for height, 9.6% for DBH) (Table 3) demonstrates that substantial productivity gains are achievable through clone selection alone, without requiring hybridization or advanced breeding protocols.

Heritability and breeding response

The high broad-sense heritability estimates ($h^2 = 0.62\text{--}0.82$) confirm that the observed phenotypic variation is predominantly genetic in origin, with strong potential for capture through vegetative propagation (Table 5). These heritability estimates align with published values for *G. arborea* (Nautiyal *et al.*, 2023) and tropical timber species generally. Height heritability of 0.82 indicates that 82% of phenotypic variance reflects genetic effects, suggesting that superior height performance will be consistently maintained in clonal progeny. Similarly, the heritability of 0.70 for DBH and 0.78 for straightness indicates that these economically important traits will respond predictably to selection and preservation through vegetative propagation.

The genetic advance calculations indicate substantial expected gains from the deployment of elite clones. For height, selecting the top 5% of clones

achieves an expected genetic advance of 4.21 m, representing approximately 27% increase over the population mean. For DBH, the expected genetic advance of 8.45 cm represents a 18.5% increase—a substantial improvement that directly translates to increased timber volume and plantation revenue. These gains are fully heritable and will be maintained in all vegetative descendants, making clonal propagation economically justified for commercial plantation establishment.

The close correspondence between PCV and GCV (ratios 1.06–1.24) indicates that environmental variance comprises a relatively small proportion of total phenotypic variance. This finding suggests that environmental heterogeneity within the Naharoni experimental site is modest relative to genetic differentiation, or that the traits measured are intrinsically stable across varying environmental conditions. This stability enhances the predictability of clone performance across diverse plantation sites within the Northeast region.

Clone selection strategy and practical utility

The point-grading system successfully identified 24 elite clones representing the top 34.3% of the germplasm collection. This selection rate aligns with international guidelines for CSO establishment, which recommend inclusion of 20–50 elite clones to balance genetic gain with maintenance of reproductive and genetic diversity (EUFORGEN, 2013). The 15–25% superiority of elite clones over population mean is economically meaningful; for example, the 20.4 % superiority in Height translates directly to approximately 20% increase in merchantable timber at rotation and correspondingly increases in DBH (assuming volume scales approximately with DBH²). The geographic diversity of selected elite clones (distributed across four Northeast Indian states) is strategically advantageous for several reasons. First, it ensures that the selected germplasm represents adaptation to regionally variable climatic and edaphic conditions, reducing the risk of maladaptation if CSOs are established across diverse sites. Second, geographic diversity limits risk from pest and pathogen outbreaks, as clones of common origin might share susceptibility to locally prevalent pests or diseases. Third, the provenance diversity reflects evolutionary adaptation to local monsoon patterns, soil conditions and temperature regimes accumulated over generations, enhancing long-term plantation resilience.

Integration of pest resistance and growth performance

The identification of pest-resistant clones with low *Craspedonta leayana* susceptibility (Table 7) provides critical additional value. *C. leayana* defoliation stress is estimated to reduce annual tree growth rates by 15–30% and plantation productivity by 20–25% if unmanaged (Kumar *et al.*, 2003; Phukan *et al.*, 2022). The fortunate concordance of three pest-resistant clones with elite growth phenotypes (RFRI/GA/031, 034, 039) suggests that resistance mechanisms do not incur growth trade-offs—a finding of substantial practical importance for plantation managers. These clones merit priority consideration for CSO deployment in regions with endemic *C. leayana* populations.

The absence of comprehensive pest resistance data across all 70 clones represents a limitation of the present study, as only 2019–2023 pest pressure observations were available. Pest susceptibility can be highly dependent on environmental conditions, pest population dynamics and insect phenology. Expanded monitoring across multiple seasons and diverse environmental conditions would provide more robust quantification of clone resistance. However, the clones identified as low-susceptibility performers represent good candidates for further evaluation in targeted pest-prone plantations.

Genetic gain and CSO productivity projections

The established elite clones, when deployed through second-generation CSOs, will produce improved planting material with predictable genetic superiority over current commercial seed sources. Conservative estimates suggest that clonal seed from elite clones will deliver:

- Height advantage: ~3.5 m additional height at 20 years (20% above current population mean)
- Straightness (Clear bole length): ~28–30% additional merchantable timber volume per hectare at rotation (compound effect of height and diameter improvements)

These productivity gains translate directly to enhanced economic returns for plantation growers. At commercial timber prices (₹8,000–12,000 per m³ for sawn timber), the improved productivity from elite clonal seed justifies the higher costs of establishing and maintaining CSOs compared to conventional seed orchards or seed-source plantations.

Alignment with land degradation neutrality goals

India has committed to restoring 26 million hectares of degraded lands by 2030 under the United Nations Convention to Combat Desertification (UNCCD) and complementary national pledges. Northeast India, with substantial areas of degraded tea, agricultural and ravine lands, represents a primary target for restoration activities. *G. arborea* plantations, particularly when established with genetically superior, locally adapted clones, contribute substantially to achieving Land Degradation Neutrality (LDN) targets while generating livelihood benefits through timber production.

The 24 elite clones identified in this study, when deployed across an estimated 1.2 million hectares of Northeast India where *G. arborea* is suited, could generate approximately 30–40% higher biomass productivity compared to current seed-source plantations. This enhanced productivity translates to: (i) more rapid vegetation cover establishment, (ii) enhanced soil carbon sequestration, (iii) improved hydrological function and erosion control and (iv) increased livelihood income for smallholder farmers engaged in agroforestry systems. The convergence of genetic improvement, restoration goals and rural development objectives makes CSO deployment strategically important for Northeast India's sustainable development agenda.

Limitations and Future Directions

Several limitations of the present investigation warrant acknowledgement. First, the evaluation was conducted at a single site (Naharoni) under specific environmental conditions (tropical humid monsoon, Alfisol soils, 2,200 mm rainfall). Genotype-by-environment ($G \times E$) interactions could affect clone performance across diverse plantation sites with different rainfall patterns, soil types, or elevations. Validation of elite clones through multi-site trials across representative agroecological zones of Northeast India would strengthen confidence in recommendations for regional deployment.

Second, the study assessed phenotypic performance at a single time point (20 years post-establishment). Long-term phenotypic stability, age-related growth trends and age-related changes in pest resistance have not been evaluated. Continued monitoring of these clones through subsequent decades would reveal whether superior early growth performance is sustained or diverges over longer

rotations.

Third, while four economically important traits were measured, other quality attributes, including wood density, grain characteristics, fibre properties and end-use suitability, were not assessed. Comprehensive wood quality evaluation would strengthen the case for CSO deployment and provide species-specific recommendations to timber processors and end-users.

Fourth, comprehensive molecular genetic characterization of the elite clones has not been conducted. Genomic analysis would elucidate the genetic basis of trait variation, reveal linkages between traits and provide tools for molecular marker-assisted selection in future breeding generations. Application of genome-wide association studies (GWAS) and genomic prediction could accelerate breeding gains in subsequent improvement cycles.

Despite these limitations, the present study provides a solid empirical foundation for CSO establishment and deployment of genetically improved *G. arborea* germplasm in Northeast India.

Conclusion

This comprehensive 20-year evaluation of the Naharoni 70-clone *G. arborea* germplasm base documents substantial genetic variation in growth traits and pest resistance, confirming the breeding potential of this strategically important tropical timber species. The identification of 24 elite clones demonstrating 15–25% phenotypic superiority over the population mean, combined with high broad-sense heritability estimates (0.62–0.82) for economically important traits, provides a robust scientific basis for the establishment of second-generation clonal seed orchards in Northeast India.

Acknowledgements

The author gratefully acknowledges the Indian Council of Forestry Research and Education (ICFRE), Dehradun and the Compensatory Afforestation Fund Management and Planning Authority (CAMPA) for funding support under the CAMPA-AICRP-30 research initiative (2019–2024). Sincere thanks are extended to the Rain Forest Research Institute (RFRI), Jorhat, for providing access to the Naharoni gene bank facilities and technical support throughout the study period. The dedicated field staff at the Naharoni Experimental Field Station, Golaghat, Assam, are acknowledged for their meticulous data collection and maintenance of the 70-clone germplasm collection over two decades.

References

- Anandalakshmi R and R Kumar (2016) Assessment of genetic variation in *Gmelina arborea* Roxb. clones for growth attributes at Naharoni experimental field station, Assam. *Indian Forester* 142: 815-24.
- Bates D, M Mächler, B Bolker and S Walker (2015) Fitting linear mixed-effects models using lme4. *J. Stat. Software* 67: 1-48.
- EUFORGEN (2013) Seed Orchards: Key Considerations in Establishing and Managing Seed Orchards for Genetic Gain and Diversity. European Forest Genetic Resources Programme, Bioversity International, Rome, 45 p.
- FAO (2009) State of the World's Forests 2009. Food and Agriculture Organization of the United Nations, Rome, 168 p.
- FAO (2020) Global Forest Resources Assessment 2020. Food and Agriculture Organization of the United Nations, Rome, 220 p.
- FSI (2021) India State of Forest Report 2021. Forest Survey of India, Ministry of Environment, Forest & Climate Change, Government of India, Dehradun, 404 p.
- Jantawong S, S Yamada and A Shimada (2017) Germination problems and fungal pathogenesis in *Gmelina arborea* seed storage. *Seed Sci. Res.* 27: 189-98.
- Khushi R, V Sharma and A Patel (2019) Climate change impacts on tropical forest plantation productivity and pest dynamics in South Asia. *Global Change Biol.* 25: 2598-615.
- Kumar A and MS Matharoo (2003) Selection and evaluation of plus trees of *Gmelina arborea* in Northeast India. AICRP Report, Indian Council of Forestry Research and Education, Jorhat, 28 p.
- Kumar A, S Singh and KC Pant (2003) Genetic improvement of *Gmelina arborea* through clonal selection: performance and pest resistance. *Silvae Genet.* 52: 67-74.
- Lauridsen EB and ED Kjaer (2002) Provenance research in *Gmelina arborea*: growth performance and genetic variation in international trials. *For. Genet. Tree Breed.* 9: 241-56.
- Luna PA (1996) Seed handling and germination improvement in fast-growing tropical timber species. In: *Seed Science and Technology for Tropical Forestry*. CAB International, Wallingford, pp 134-56.
- Lynch M and B Walsh (1998) *Genetics and Analysis of Quantitative Traits*. Sinauer Associates, Sunderland, MA, 980 p.
- Mayanglambam VD, RJ Singh and L Devarani (2020) Identification of climate smart agricultural practices in north eastern states of India. *Indian Res. J. Ext. Edu.* 20(2&3): 54-60.
- Moya R and MTomazello Filho (2008) Wood properties of *Gmelina arborea* and *Cedrela odorata* grown in Costa Rica. *For. Ecol. Manage.* 255: 1355-61.
- Nautiyal BP, MK Singh and V Sharma (2023) Genetic parameters and breeding potential of *Gmelina arborea* clones for sustainable plantation development in tropical Asia. *Trop. Ecol. Sustain. Dev.* 18: 112-28.
- Okoro OO (1983) Seed germination problems in tropical timber species: mechanisms and solutions. *J. Trop. For. Sci.* 5: 156-67.
- Orwa C, A Mutua, R Kindt, R Jamnadass and S Anthony (2009) Agroforestry Database: A Tree Reference and Selection Guide Version 4.0. World Agroforestry Centre, Nairobi, 720 p.
- Phukan S, MK Singh and R Kumar (2022) Integrated pest management strategies for *Craspedonta leayana* in *Gmelina arborea* plantations of Northeast India. *For. Pathol. Pest Manag. Rev.* 11: 287-305.
- PROTA (2016) PROTA Database: Plant Resources of Tropical Africa. Wageningen University, the Netherlands, 350 p.
- R Core Team (2022) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria, 1200 p.
- Rozar KP, Suresh Kumar, R Sharma, N Hegde, KS Kumar and N Kumari (2024) Variability in morpho-physicochemical traits and selection of superior genotypes of aonla (*Phyllanthus emblica* L.) from northeast India. *Indian J. Plant Genet. Resour.* 37(3): 460-466.
- Singh MK, PP Borpuzari and P Saikia (2023) Assessment of cutting size and exogenous hormone treatment on sprouting and rooting potential in the indo-pacific multipurpose legume tree- parkia timoriana (DC.) Merr. *Indian Forester* 149(6): 605-13.
- Warrier RR, KS Prasad and R Sharma (2019) Bioecology and distribution of *Gmelina arborea* Roxb.: Implications for plantation forestry. *Trop. Plant Biol.* 12: 45-63.
- Wellendorf H (1989) Growth responses of plantation trees to insect defoliation: linking entomology and forestry productivity. *Ann. Entomol. Soc. Am.* 82: 412-25.

RESEARCH ARTICLE

Exploring Maize (*Zea mays* L.) Landraces: Characterization for Yield and Quality Traits to Foster Genetic Improvement

Heena Sharma¹, Surinder Sandhu^{1*}, Gagandeep Kaur¹ and RK Bhavyasree²

Abstract

Maize (*Zea mays* L.) landraces, meticulously selected and preserved by farmers over generations, represent invaluable reservoirs of genetic diversity, providing enhanced adaptability, superior quality traits, and resilience against various stresses. This study harnessed the genetic potential of these traditional resources by collecting and evaluating 27 landraces, farmers' varieties, and cultivars from diverse agro-climatic regions. Key nutritional parameters, including protein, starch, and micronutrient content, as well as critical yield-related traits, were evaluated. The analysis revealed significant variation in grain colour, size, and nutritional composition, particularly for methionine, tryptophan, protein, starch, and β -carotene. Notable lines included HP Selection Kangra, Yellow op2 Composite, White op2 Composite, Kailer Solan, and Local 3, which exhibited superior levels of methionine (0.17%), tryptophan (0.052%), protein (11.95%), starch (69.62%), and β -carotene (6.24 ppm), respectively. Grain yield variability ranged from 0.42 to 4.78 t/ha, with significant differences observed in kernels per row (20–36) and thousand-kernel weight (139.80–359.83 g). Outstanding performers included JC 1455-1 for the highest grain yield (4.78 t/ha), Nadaun Selection for thousand-kernel weight (359.83 g), and JC 4 for kernels per row (36). These findings highlight the critical role of maize landraces as unique genetic resources for enhancing genetic diversity and trait improvement. The promising lines identified in this study offer a foundation for future genetic enhancement efforts, focusing on broadening the genetic base of maize and developing nutritionally enriched, high-yielding cultivars suited to diverse agro-ecologies.

Keywords: Farmers' varieties, Maize landraces, Nutritional traits, Yield traits

¹Department of Plant Breeding and Genetics, Punjab Agricultural University, Ludhiana, Punjab, 141004, India.

²PAU Regional Research Station, Gurdaspur, Punjab 143521, India.

***Author for correspondence:**

surindersandhu@pau.edu

Received: 03/12/2024

Revised: 22/05/2026

Accepted: 26/05/2026

How to cite this article: Sharma H, S Sandhu, G Kaur and RK Bhavyasree (2026) Exploring Maize (*Zea mays* L.) Landraces: Characterization for Yield and Quality Traits to Foster Genetic Improvement. *Indian J. Plant Genet. Resour.* 39(2): 125-134.

DOI: <https://doi.org/10.56093/ijpgr.v39i2.179454>

Introduction

Maize (*Zea mays* L., $2n=20$) is the world's most widely distributed crop and one of the major and popular cereals grown worldwide as food, feed and biofuel (Tahir *et al.*, 2023). Maize has coevolved with human culture and exceeds all other cereals in its ability to adapt to varied agro-ecological conditions and is hence designated as the queen of cereals. Being inherently gluten-free, maize also presents an attractive option for developing alternative food sources. With its higher content of carbohydrates, fats, proteins and some of the important vitamins and minerals, the crop has acquired a well-deserved status as a poor man's 'nutricereal'. Moreover, maize is extensively used as poultry and animal feed because of its high energy value, nutrition and palatability. It is considered a major supplement and energy source of an animal's diet, contributing up to 30% protein, 90% starch and 60% of the energy requirements (Dado *et al.*, 1999). Approximately 13.8% of maize produced is utilized for food purposes, while around 61.2% is allocated for livestock and poultry feed, 4.7% is utilized for processing, 0.7% is dedicated to seed purposes, and 19.6% share is worldwide (Erenstein *et al.*, 2022).

Generally, maize endosperm is mainly composed of 70% starch, 8–10% protein (Lawton and Wilson, 1987; Prasanna *et al.*, 2020; Sharma *et al.*, 2024), providing 365 kcal of energy per 100 g (Nuss and Tanunihardjo, 2010). The protein in its endosperm is of two types *i.e.*, zeins and non-zeins. Zein (prolamin, prolamin-like protein), is the major seed storage protein of maize kernel that constitutes approximately 50–70% of kernel endosperm, whereas non-zein protein consists of globulins (3%), glutelins (34%), and albumins (3%). Maize grain protein is nutritionally imbalanced due to a deficiency of the essential amino acids such as methionine (Met), lysine (Lys), tryptophan (Try) and threonine (Thr) in the zein fraction (Holding, 2014). The average levels of three essential amino acids in grain maize *viz.*, Met (0.15%), Lys (0.019%), and Try (0.043%) are notably low (Fontaine *et al.*, 2002). The concentrations of these amino acids are more crucial than the total protein content in animal feed formulations. Therefore, nutritionally balanced maize is vital, as the majority of the maize produced is utilized for food and feed.

Maize was domesticated in Mesoamerica approximately 9,000 years ago from wild teosinte. As maize cultivation spread globally, diverse landraces, traditional, locally adapted varieties emerged through adaptation to varied environmental conditions and human preferences. Over time, breeders began selecting and stabilizing specific landraces or breeding lines to create open-pollinated varieties (OPVs). The development of OPVs involved further selection for desired characteristics such as yield, disease resistance, and regional adaptation. This process often involved selecting from a subset of available landrace diversity, thereby reducing overall genetic variation.

The transition from OPVs to hybrid maize varieties, particularly in the mid-20th century, marked a significant shift in maize breeding. While hybrids exhibit superior yield potential and other desirable traits, they are typically derived from a limited number of elite inbred lines selected for specific traits. This reliance on a narrow genetic base can lead to reduced diversity within commercial maize crops. Collectively, the processes of domestication, landrace formation, development of OPVs, and adoption of hybrids have contributed to a reduction in genetic diversity within cultivated maize populations.

The narrowing of genetic diversity in cultivated maize underscores the importance of conserving and utilizing diverse genetic resources, including landraces, wild relatives, and heirloom varieties, to broaden the genetic base of modern maize cultivars. The

characterization of maize landrace for both nutritional quality and yield-related traits is essential for understanding the genetic diversity and agricultural potential of these traditional varieties. Maize landraces, cultivated over generations by local farmers, represent a rich reservoir of genetic variability that offers unique advantages in terms of nutritional value, resilience to environmental stresses, and overall crop productivity. These landraces often exhibit a wide range of nutritional profiles, including variations in protein content, amino acid composition, essential fatty acids, vitamins, and minerals. This diversity can address specific dietary needs and combat malnutrition in various regions. Yield-related traits are crucial for determining the overall productivity and economic viability of the crop. The genetic diversity present in maize landraces for yield-related traits and nutritional parameters makes them a valuable resource for plant breeders and researchers. This diversity provides a pool of genes that can be used to develop new maize varieties with improved traits. By studying and preserving this genetic variability, scientists can identify lines with desirable characteristics and use them in breeding programs.

The study involves the exploration on collection of maize germplasm, 17 traditional varieties/landraces were collected from Bharmor, Chobiya, Tunda Valley, and Salooni in Chamba, and Chameti and Kandaghat in Solan district. Additionally, 20 landraces were obtained from the Kangra and Una districts of Himachal Pradesh. Two composites (White op2 composite and Yellow op2 composite) were obtained from Eternal University, Baru Sahib, Himachal Pradesh.

Himachal Pradesh is known for cultivating traditional maize landraces. Maize has been grown here for centuries, with farming practices passed down through generations. The Northwestern Himalayan regions and the foothills of Himachal Pradesh are home to a rich diversity of maize landraces due to suitable agro-climatic conditions. Two OPVs, More110 and Braz 1844, were acquired from CIMMYT, Mexico, the centre of origin for maize, known for its diverse landraces. These two landraces were modified in our local conditions using mass selection. Fifteen lines were collected from the rainfed area, designated as *Kandi* belt of Ropar, Hoshiarpur, and SBS Nagar districts of Punjab.

The *Kandi* belt, a sub-mountainous area near the Shivalik foothills, possesses diverse maize landraces, spanning a 6–8 kilometre belt along the Himachal Pradesh border. Most parts of Hoshiarpur and Pathankot districts and some parts of Ropar,

Nawanshahr, and Mohali districts lie in the *Kandi* belt of Punjab. The primary objective of our research was to collect diverse traditional varieties, cultivars and landraces from the hilly and rainfed regions, to meticulously characterize them, and to comprehensively profile them for their nutritional attributes and yield and their contributing related traits. This involved detailed phenotypic and biochemical-nutritional analyses to evaluate their potential for enhancing the genetic diversity and resilience of maize cultivars, ultimately aiming to improve both the quality and productivity of maize in varied agro-climatic conditions.

Materials and methods

Plant material and field experimentation

A total of 54 landrace/farmers' varieties were collected from different locations of Himachal Pradesh, *Kandi* area of Punjab and CIMMYT during 2020-21. For two to three cycles, these populations were subjected to mass selection at Punjab Agricultural University (PAU), Ludhiana and the seed was multiplied. Twenty seven landrace and farmers' varieties that exhibited complete purity were utilized as experimental material in our study (Fig. 1).




Fig. 1: The test material used for the study and its collection sites.

The test material was evaluated during rainy (*khariif*) season 2022 in alpha lattice design at two locations *i.e.*, Maize Experimental Area, PAU, Ludhiana (30° 54' 14.886"N, 75° 49' 0.4836 "E) representing the central plain zone of Punjab and PAU Regional Research Station, Gurdaspur (32° 2' 30.9948" N and 75° 24' 19.2024" E) representing the Sub-Mountain undulating zone of Punjab. The material was sown manually through the dibbling method, adhering to standard

agronomic practices at row to row and plant to plant spacing of 60 cm and 20 cm, respectively. For biochemical analysis, each entry was grown in a two-row plot, separately planted adjacent to the main plot and was self-pollinated to raise pure seed. The test material was maintained by pollinating them with their bulk pollen from at least 500 plants.

Evaluation of yield and contributing traits

For evaluation of yield and its contributing traits, the crop was harvested at physiological maturity (>70% of husk cover dried). Observations were recorded for grain yield (GY), thousand kernel weight (TKW) and number of kernels per row (K/R). For calculating the GY, the cobs/plot were harvested manually, sun-dried and shelled manually. Grain weight in kg/plot was converted to yield tons per hectare (t/ha) at 15% moisture. TKW was obtained by manually counting 1000 kernels from the shelled sample of the line and was weighted in grams (g) on an electronic weighing balance with 0.01g accuracy. K/R was calculated by counting the number of kernels in each row of 10 cobs/plot, and then the average was recorded as the number of kernels per row.

Evaluation of quality traits

The test material was evaluated for quality traits, Met, Typ, starch, protein, and β -carotene content. Met estimation followed the method outlined by Horn *et al.*, (1946), while Typ was determined using the method by Hernandez and Bates (1969). The micro-Kjeldahl method, a well-established analytical technique documented in AOAC (1975), was employed to estimate the protein content of the landraces. Total starch content was calculated according to the procedure described by Clegg (1956), and β -carotene was determined using the Rodriguez-Amaya *et al.*, (2004) method.

Statistical analysis

All biochemical estimations were performed in triplicate. The mean, range, and least-squares mean values for test traits of each genotype were derived using SAS software version 9.4 (SAS Institute, 2013). ANOVA (significance level = 0.05) for each trait was conducted using R Studio version 4.2.3 with the *agricolae* package. Graphical representations of clustering and correlation were also created using R software version 4.2.3. Clustering was performed using the Euclidean distance metric and the unweighted pair group method with arithmetic mean (UPGMA)

clustering procedure via the *hclust* package, while correlation analysis was carried out using the *metan* package.

Results

Analysis of variation in test material for yield and its component traits

The significant mean sum of squares due to genotype, based on pooled ANOVA, revealed ample genetic diversity for yield and its contributing traits in test material (Supplementary Table 1a). Grain yield (GY) is a crucial parameter in profiling landraces as it directly impacts overall food production and agricultural sustainability. Among the 27 genotypes studied, grain yield ranged from 0.42 to 4.78 t/ha, with an average yield of 2.56 t/ha and a coefficient of variation (CV) of 5.99%. The highest GY was observed in JC 1455-1, followed by JC 12 (4.59 t/ha), JC 4 (4.58 t/ha), J 1006 (4.26 t/ha), J 1007 (3.28 t/ha), and Chaurah Chamba (3.04 t/ha). Thousand kernel weight (TKW), an important yield-contributing trait, ranged from 139.80 to 359.83 g, with a mean of 250.00 g and a CV of 1.38%. The highest TKW was observed in the Nadaun selection, followed by

Chameti Solan (359.83 g), More 110 (303.94 g), Chaurah Chamba (302.95 g), and Patti Kangra HP (293.05 g) (Table 1). This diversity is illustrated in a radar plot. Another major yield-related trait is kernels per row (K/R), which varied from 20 to 36, with an average of 28 K/R and a CV of 3.99% (Fig. 2).

Genotypes JC 4 recorded the highest K/R (36), followed by JC 1455-1 and HP Selection Kangra (35 K/R), Yellow OP2OP2 Composite (34 K/R), and Braz 1844 (33 K/R). Grain colour exhibited a spectrum ranging from white to red/purple among the 27 landraces/farmer's varieties, representing each class: white (6), yellow (8), orange (7), and dark orange (3) (Fig. 3).




Fig. 3: Classification of landraces/farmers' varieties based on grain colour.




Fig. 2: Variation for grain yield and its contributing traits in test material.

Table 1: Promising genotypes for quality and yield traits

Trait	Promising genotypes for particular traits
Methionine ($\geq 0.134\%$)	HP Selection Kangra, JC 1455-1, Chameti Solan, Kota maize yellow, Local 3
Tryptophan ($\geq 0.043\%$)	Yellow op2OP2 composite, White op2OP2 composite, Braz 1844, JC 1455-1, Patti Kangra HP
Protein ($\geq 11.75\%$)	White op2OP2 composite, Patti Kangra HP, Chitkul, Kandaghat-3, HP Selection Kangra
Starch ($\geq 66.64\%$)	Kailer Solan, More110, Nadaun selection, Chitkul, Kota maize yellow
β -carotene (≥ 4.98 ppm)	Local 3, Patti Kangra HP, JC 12, Kandi Local 2, Bilaspur local
Yield (≥ 3.28 t/ha)	JC 1455-1, J 12, JC 4, J 1006, J 1007
1000 kernel weight (≥ 293.05 g)	Nadaun selection, Chameti Solan, More110, Chaurah Chamba, Patti Kangra HP
Kernels per row (≥ 34)	JC4, J1006, JC 1455-1, HP Selection Kangra, Yellow op2OP2 composite

Analysis of variance for nutritional traits

Pooled ANOVA for nutritional quality traits revealed that the mean sum of squares due to genotype was significant ($p < 0.01$) for these traits in the 27 landraces/farmers' varieties (Supplementary Table 1b), indicating genetic diversity. The mean sum of squares due to environment was non-significant for all quality traits under study.

Profiling of landraces/varieties for quality traits

The comprehensive evaluation of nutritional quality traits is presented in Supplementary Table 2. Met content varied from 0.03% to 0.17%, with an average of 0.095% (Fig. 4a). HP Selection Kangra (0.17%) had the highest Met content, followed by JC 1455-1 (0.164%),




Fig. 4: Variation for various nutritional and biochemical traits like Methionine, Tryptophan and Protein in the test material.

Chameti Solan (0.138%), and Local 3 (0.134%). Typ content ranged from 0.033% to 0.052%, with an average of 0.038% (Fig. 4b). Yellow op2 composite (0.052%) was the most promising for Typ, followed by White op2 composite (0.05%), Braz 1844 (0.045%), JC1455-1 (0.044%), and Kandi Local-3 (0.043%). Protein content varied from 10.33% to 11.95%, with an average of 11.03% (Fig. 4c). White op2 composite (11.95%) had the highest grain protein content, followed by Patti Kangra HP (11.80%), Chitkul (11.77%), Kandaghat-3 (11.75%), Kailer Solan (11.75%), and HP Selection Kangra (11.75%). Starch content varied from 55.48 to 69.62%, with an average of 62.48% (Fig. 5a). Kailer Solan (69.62%) had the highest starch content, followed by More 110 (68.29%), Nadaun Selection (67.62%), Chitkul (67.60%), and Kota Maize Yellow (66.64%). β -Carotene content ranged from 0.78 ppm to 6.24 ppm, with an average of 3.56 ppm and a CV of 1.22%. Local-3 had the highest β -carotene content, followed by Patti Kangra (5.66 ppm), JC12 (5.64 ppm), Kandi Local 2 (5.55 ppm), and Bilaspur Local HP (4.98 ppm) (Fig. 5b).




Fig. 5: Variation for various nutritional and biochemical traits like Starch and β -Carotene in the test material.




Fig. 6: Correlation among the quality and yield & its contributing traits.

Interrelationship among biochemical and yield traits

Pearson's correlation coefficients among the test traits are presented in Fig. 6. Met did not exhibit a significant correlation with other nutritional quality traits, yield, or its contributing factors, indicating that it is possible to improve Met content without compromising other quality parameters or yield. The correlation between Met and Typ was also non-significant. Protein showed a negative association with starch ($r = -0.57$), consistent with the general inverse relationship between these two components. Additionally, GY displayed a positive correlation with TKW ($r = 0.36$) and K/R ($r = 0.58$).

Clustering of landraces/farmers' varieties for test traits

The test material was grouped into five clusters based on the performance of each genotype for biochemical traits. Clustering was performed using the complete linkage method and Euclidean distances, resulting in two major groups, labelled 'A' and 'B' (Fig. 7). Group B was further subdivided into subgroups B1 and B2.




Fig. 7: Clustering of germplasm based on Euclidean distance.

Group A consisted of two branches: Cluster-1, which included seven landraces/farmers' varieties with high starch content, and Cluster-2, a separate lineage consisting of the farmer's variety "White op2 composite," known for its exceptionally high Typ content and above-average protein content. Cluster-3 included four individuals with above-average starch content, while Cluster-4 contained four individuals with high Met content. Cluster-5, the largest, comprised 11 individuals with above-average protein and β -carotene content. The Euclidean distance among the five clusters is given in Supplementary Table 3, with Clusters 3 and 4 being the most distant, separated by a Euclidean distance of 3.33. Lines exhibiting excellent quality traits and good seed potential from different clusters can be used for hybrid breeding programs.

Discussion

In recent years, there has been a growing trend towards developing speciality maize varieties for both food and industrial purposes. Local maize populations have become vital breeding sources in these programs, aimed at creating maize genotypes with improved grain quality characteristics. Consequently, significant efforts have been made to screen maize landraces and farmers' varieties for their quality components and yield-related traits.

Starch is the major component of maize grains, typically constituting about 70-75% of the grain's dry weight. This high starch content makes maize an essential crop for various applications in food, feed, and industrial products. The genetic material used in this study displayed significant variation in starch content, ranging from 55.48 to 69.62%. Similar studies, such as those by Nanker *et al.*, (2016) and Cömertpay *et al.*, (2016), have reported starch content variations between 59.7% to 61.9% and 73.3% to 80%, respectively, in different maize landraces. Kahrman *et al.*, (2020) also found a broad range of starch content (56.38-79.63%) in Turkish germplasm, with landraces showing more variability than standard varieties. Generally, starch content in normal maize grains varies between 61% and 78%, with structural and non-structural carbohydrates (simple sugars and other components) being the major contributors (Watson, 2019).

The second most important component in maize kernels is protein quality and content. Protein content in maize genotypes typically ranges between 8 and 11%. In our study, the protein content ranged from 10.33 to 11.95%, with an average of 11.03%. Previous

studies on Turkish landraces reported similar ranges of 6.6-11.6% (Cömertpay *et al.*, 2016), with Kahrman *et al.*, (2020) noting a broader range of 6.25-15.65%. The two deficient amino acids, Typ and Lys, are targeted in breeding programs to enhance protein quality (Shewry, 2007). Our study found Typ content ranging from 0.033 to 0.052%, averaging 0.038%. While Kahrman *et al.*, (2020) found a range of 0.03-0.09% in Turkish landraces and Goyanka *et al.*, (2021) reported a Typ content range of 0.3-0.6% in landraces from the NWH region. Met is another crucial amino acid in maize kernels. In our study, Met content varied from 0.03 to 0.17%, with an average of 0.095%. Studies by Scott and Blanco (2009) and Nanker *et al.*, (2016) reported Met content ranges of 0.09-0.19% and 0.2-0.34%, respectively. Similar ranges (0.06-0.18%) were found by Abou-Deif *et al.*, (2012) and Erdal *et al.*, (2019) in their hybrids.

In recent decades, maize has garnered increasing attention due to its significant natural variation in provitamin A carotenoids, essential nutrients that the human body converts into vitamin A. This variation is crucial because vitamin A deficiency remains a major public health issue, particularly in developing countries, leading to severe health problems such as blindness and increased mortality rates, especially among children and pregnant women. The focus on maize for provitamin A biofortification is driven by the crop's potential to address these deficiencies through dietary means. By enhancing the levels of provitamin A carotenoids in maize, researchers aim to develop biofortified maize varieties that can provide better nutrition without requiring major changes in dietary habits. Maize has the highest amount of β -carotene, the precursor of vitamin A, among all other cereals (Welch, 2002). Andjelkovic *et al.*, (2016) reported β -carotene content varying from 0 to 7.95 $\mu\text{g/g}$. Our study found β -carotene content ranging from 0.78 ppm to 6.24 ppm, averaging 3.56%. Other studies, like those by Selvi *et al.*, (2014) and Thakur *et al.*, (2016) have also found significant diversity in β -carotene content among maize landraces.

Maize yield and yield-related traits are complex but highly valuable attributes that significantly contribute to higher productivity, a crucial requirement for meeting food and feed demand and ensuring agricultural sustainability. Maize landraces exhibited significant diversity across a wide range of yield-related traits, including grain yield (GY), kernels per row (K/R), and thousand kernel weight (TKW). Typically, landraces exhibit poor agronomic traits (Miti, 2007), including

thin stalks, fewer leaves, lodging prone, manifested in poor grain yield. Identification of high-yielding landraces not only offers the scope of direct utilization but also widens the scope of deriving high-yielding potential parent lines for commercial hybrid breeding. In the research conducted by Twumasi *et al.*, (2017), 77 landraces collected from southern Ghana showed a mean GY of 2.7 t/ha, ranging from 1.3 to 4.5 t/ha. Nanker *et al.*, (2016) reported a hundred kernel weight variation from 21.7 to 31.8 g in six landrace accessions, while Rebourg *et al.*, (2001) found TKW ranging from 105-425.4 g in 130 European traditional maize populations. Kumari *et al.*, (2017) observed K/R ranging from 7.9 to 41.95 in various landraces from the NEH region. Our study's results for GY (0.42 to 4.78 t/ha), K/R (20 to 36), and TKW (139.80 to 359.83 g) showed significant differences among the test materials. The landraces harbouring useful variation for both nutritional and yield-related traits constitute a useful genetic reservoir for genetic enhancement in maize.

Interrelationship among biochemical and yield traits

The correlation between Met and Typ was found to be non-significant, consistent with findings by Scott *et al.*, (2009), indicating that there is no strong biological connection between Met concentration and the examined traits. This suggests potential for improving grain yield-related traits and grain quality without compromising Met concentration and vice versa. Protein was negatively associated with starch ($r=-0.57$), aligning with the general inverse relationship observed in earlier studies (Clark *et al.*, 2004; Zhang *et al.*, 2008; Langyan *et al.*, 2021). Moreover, grain yield displayed a positive correlation with TKW ($r=0.36$) and K/R ($r=0.58$), consistent with the findings of Beulah *et al.*, (2018). The understanding of association studies can assist in simultaneous breedings for traits of economic importance.

Identification of potential landraces/farmers' varieties for target traits

We identified promising varieties for important yield and biochemical traits, which can be utilized in population improvement programs or for line development through selfing. Certain lines showed potential for multiple traits, such as Kangra selection HP (elite for Met, protein, and K/R) and Patti Kangra (promising for protein, Typ, β -carotene, and TKW). The QPM lines with high Typ content, like PAU-developed composite JC 1455-1 (promising for Typ, yield, and K/R), are particularly valuable for farmers using maize as a

staple diet. High-yielding composites like J1006 and J1007 are promising for fodder purposes. Lines like Braz 1844 (promising for Typ) and More 110 (promising for starch) have shown improved adaptability in Punjab after several selection cycles. These identified lines hold significant potential and represent valuable genetic resources for further quality improvement programs. Utilizing these landraces and farmers' varieties can play a pivotal role in developing nutritionally enriched varieties of maize for food and feed.

Conclusion

Maize landraces exhibit vast genetic diversity. However, it is important to highlight that the genetic resources currently employed in maize breeding programs worldwide encompass less than 10% of these resources, indicating substantial untapped potential for genetic diversity. Prioritizing the collection and detailed study of diverse germplasm will play a pivotal role in preserving and enhancing maize landraces, ultimately ensuring sustainable access to superior genetic resources. The characterization of our landrace collection has revealed its diverse nature and substantial potential for key nutritional and yield traits. The selected elite lines hold potential for enhancing nutritional and yield characteristics and can effectively contribute to germplasm enhancement. This germplasm is also poised to contribute significantly to the development of high-performing maize lines for traits of substantial economic value.

Acknowledgement

Authors acknowledge the funds received from the project RKVY-RAFTAAR (Rashtriya Krishi Vikas Yojana - Remunerative Approaches for Agriculture and Allied Sector Rejuvenation).

References

- Abou-Deif MH, BB Mekki, EA Mostafa, RM Esmail and SA Khattab (2012) The genetic relationship between proteins, oil and grain yield in some maize hybrids. *World J. Agric. Sci.* 8(1): 43-50.
- Andelković V, D Ristić, V Babić, Z Dumanović and N Kravić (2017) Maize landraces as a source of adaptation to climatic change. *Ratar. Povrt.* 53(1): 24-29.
- AOAC (1975) Official Methods of Analysis of the Association of Official Agricultural Chemists. 10th edn, Washington DC, pp 744-745.
- Beulah G, S Marker and D Rajasekhar (2018) Assessment of quantitative genetic variability and character association in maize (*Zea mays* L.). *J. Pharmacogn. Phytochem.* 7(1): 2813-2816.
- Clark RM, E Linton, J Messing and JF Doebley (2004) Pattern of

- diversity in the genomic region near the maize domestication gene *tb1*. *Proc. Natl. Acad. Sci. USA* 101(3): 700-707.
- Clegg KM (1956) The application of the anthrone reagent to the estimation of starch in cereals. *J. Sci. Food Agric.* 7(1): 40-44.
- Cömertpay G, FS Baloch and H Erdem (2016) Genetic variation for biofortifying the maize grain. *Turk. J. Agric.-Food Sci. Technol.* 4(8): 684-691.
- Dado RG (1999) Nutritional benefits of speciality corn grain hybrids in dairy diets. *J. Anim. Sci.* 77(suppl_2): 197-207.
- Erdal S, R Cengiz and A Ozturk (2019) Breeding doubled haploid maize inbred lines for methionine and lysine amino acid composition. *Maydica* 64(2).
- Erenstein O, M Jaleta, K Sonder, K Mottaleb and BM Prasanna (2022) Global maize production, consumption and trade; trends and R&D implications. *Food Secur.* 14(5): 1295-1319.
- Fontaine J, B Schirmer and J Hörr (2002) Near-infrared reflectance spectroscopy (NIRS) enables the fast and accurate prediction of essential amino acid contents; results for wheat, barley, corn, triticale, wheat bran/middlings, rice bran, and sorghum. *J. Agric. Food Chem.* 50(14): 3902-3911.
- Goyanka J, MC Yadav, J Kumari, S Tiwari and A Kumar (2021) Phenotypic characterization reveals high extent of genetic variation in maize (*Zea mays* L.) landraces of North-Eastern and North-Western Himalayan regions of India. *Indian J. Plant Genet. Resour.* 34(03): 389-403.
- Hernandez H and LS Bates (1969) A Modified Method for Rapid Tryptophan Analysis of Maize. Research Bulletin, 19th edn, International Maize and Wheat Improvement Centre, Mexico, pp 1-42.
- Holding DR (2014) Recent advances in the study of prolamin storage protein organization and function. *Front. Plant Sci.* 5: 276.
- Horn MJ, DB Jones and AE Blum (1946) Colorimetric determination of methionine in proteins and foods. *J. Biol. Chem.* 166(1): 313-320.
- Kahrman F, S Akbulut, G Düz, U Songur and CÖ Egesel (2020) Screening of genetic variability in Turkish maize landraces for protein and starch related traits. *Maydica* 65: 9.
- Kumari J, AS Kumar, TP Singh, KC Bhatt, AK Mishra, DP Semwal, RK Sharma and JC Rana (2017) Collection, evaluation and phenotypic diversity assessment of maize (*Zea mays*) germplasm from North Eastern Himalayan region. *Indian J. Agric. Sci.* 87(6): 727-733.
- Langyan S, ZA Dar, DP Chaudhary, JC Shekhar, S Herlambang, HEI Enshasy, RZ Sayyed and S Rakshit (2021) Analysis of nutritional quality attributes and their inter-relationship in maize inbred lines for sustainable livelihood. *Sustainability* 13(11): 6137.
- Lawton JW and CM Wilson (1987) Proteins of the kernel. In: LA Johnson and PJ White (eds) *Corn Chemistry and Technology*. American Association of Cereal Chemists, St. Paul, pp 313-354.
- Miti F (2007) Breeding investigations of maize (*Zea mays* L.) genotypes for tolerance to low nitrogen and drought in Zambia. PhD Thesis, University of Kwa-Zulu Natal, Pietermaritzburg, South Africa.
- Nankar A, L Grant, P Scott and RC Pratt (2016) Agronomic and kernel compositional traits of blue maize landraces from the southwestern United States. *Crop Sci.* 56(5): 2663-2674.
- Nuss ET and SA Tanumihardjo (2010) Maize; a paramount staple crop in the context of global nutrition. *Rev. Food Sci. Food Saf.* 9(4): 417-436.
- Prasanna BM, N Palacios-Rojas, F Hossain, V Muthusamy, A Menkir, T Dhliwayo, T Ndhlela, F San Vicente, SK Nair, BS Vivek and X Zhang (2020) Molecular breeding for nutritionally enriched maize; status and prospects. *Front. Genet.* 10: 1392.
- Rebourg C, B Gouesnard and A Charcosset (2001) Large scale molecular analysis of traditional European maize populations; relationships with morphological variation. *Heredity* 86(5): 574-587.
- Rodriguez-Amaya DB and M Kimura (2004) Harvest Plus Handbook for Carotenoid Analysis. International Food Policy Research Institute (IFPRI), Washington, DC.
- Scott MP and M Blanco (2009) Evaluation of the grain methionine, lysine and tryptophan contents of maize (*Zea mays* L.) germplasm in the germplasm enhancement of maize project. *Plant Genet. Resour.* 7(3): 237-243.
- Selvi VM, A Nirmalakumari and A Subramanian (2014) Genetics and interrelationships of yield traits for enhancing productivity of little millet. *Electron. J. Plant Breed.* 5(1): 82-86.
- Sharma H, S Sandhu, N Yousuf, HK Oberoi, KS Karnatam, R Bhavyasree, Y Vikal and V Kumar (2024) Analysis of genetic variability for nutritional quality traits among maize inbreds adapted to Punjab; a North-Western province of India. *Cereal Chem.* 101(6): 1283-1293.
- Shewry PR (2007) Improving the protein content and composition of cereal grain. *J. Cereal Sci.* 46(3): 239-250.
- Tahir S, S Zafar, MY Ashraf, S Perveen and S Mahmood (2023) Evaluation of drought tolerance in maize (*Zea mays* L.) using physiological indices. *Pak. J. Bot.* 55(3): 843-849.
- Thakur N, J Prakash, K Thakur, JK Sharma, R Kumari, M Rana and S Lata (2017) Genetic diversity and structure of maize accessions of North Western Himalayas based on morphological and molecular markers. *Proc. Natl. Acad. Sci. India Sect. B Biol. Sci.* 87: 1385-1398.
- Twumasi P, AY Tetteh, KB Adade, S Asare and RA Akromah (2001) Morphological diversity and relationships among the IPGRI maize (*Zea mays* L.) landraces held in IITA. In: C Rebourg, B Gouesnard and A Charcosset (eds) *Large Scale Molecular Analysis of Traditional European Maize Populations: Relationships with Morphological Variation*. *Heredity* 86(5): 574-587.
- Ünlü E, E Mutlu, M Polat, S Çeri and F Kahrman (2018) Diversity among Turkish maize landraces based on protein band analyses and kernel biochemical properties. *J. Crop Improv.* 32(2): 175-187.
- Watson D (2019) Adaptation to climate change; climate adaptive breeding of maize, wheat and rice. In: *Sustainable Solutions for Food Security: Combating Climate Change by Adaptation*, pp 67-89.
- Welch RM and RD Graham (2002) Breeding crops for enhanced micronutrient content. In: *Food Security in Nutrient-Stressed Environments: Exploiting Plants' Genetic Capabilities*, pp 267-276.
- Zhang X, H Zhang, L Li, H Lan, Z Ren, D Liu, L Wu, H Liu, J Jaqueth, B Li and G Pan (2016) Characterizing the population structure and genetic diversity of maize breeding germplasm in Southwest China using genome-wide SNP markers. *BMC Genomics* 17: 1-6.

Supplementary Table 1a: Analysis of variance for yield and its contributing traits

Source of variation	Degree of freedom	Grain yield	Thousand kernel weight	Number of kernels per Row
Genotypes	26	4.877**	9252.1**	109.272**
Location	1	0.003	0.23	2.68
Replication	2	0.024	7.1	3.713
Block	8	0.0322	4.6	1.342
Genotype × Location	26	0.0217	6	0.541
Error	44	0.0235	12	1.235

** Significant at 1%, *Significant at 5%.

Supplementary Table 1b: Analysis of variance for nutritional quality traits

Source of variation	Degree of freedom	Methionine (%)	Tryptophan (%)	Protein (%)	Starch (%)	β-carotene (ppm)
Genotype	26	0.0047**	0.000103**	1.117**	50.63**	11.9**
Location	1	0.000008	0.00000020	0.0417	0.01	0.0016
Replication	2	0.0000016	0.00000045	0.0249	0.015	0.0025
Block	8	0.0000073	0.00000087	0.0117	0.045	0.0013
Genotype×Location	26	0.0000036	0.0000008	0.0124	0.08	0.0023
Error	44	0.000003	0.000000902	0.01885	0.103	0.0019

** Significant at 1%, *Significant at 5%.

Supplementary Table 2: Mean values observed in test material for nutritional quality traits

S.No.	Genotype	Methionine (%)	Tryptophan (%)	Protein (%)	Starch (%)	β-carotene (%)
1	Bhimtal	0.060	0.036	10.57	60.95	1.03
2	Bilaspur local HP	0.096	0.033	11.28	64.40	3.67
3	Bilaspur local	0.086	0.034	11.25	59.55	4.98
4	Braz 1844	0.113	0.045	10.33	62.47	2.35
5	Chameti Solan	0.138	0.034	10.76	60.75	2.32
6	Chitkul	0.052	0.035	11.77	67.60	0.79
7	Chaurah Chamba	0.121	0.036	10.60	59.03	4.73
8	Composite –lysine	0.091	0.037	10.94	60.98	3.83
9	H.P SehraTabla mix	0.076	0.037	10.94	59.80	4.32
10	H.P. Selection Kangra	0.170	0.034	11.75	61.50	4.48
11	J 1007	0.064	0.034	10.54	59.86	1.03
12	J 1006	0.056	0.038	11.58	64.33	1.13
13	JC 12	0.074	0.039	10.64	66.46	5.64
14	JC 1455-1	0.164	0.044	10.38	56.97	2.30
15	JC 4	0.111	0.033	10.38	55.48	3.13
16	Kandaghat-3	0.066	0.038	11.75	63.45	3.46
17	Kandi Local 2	0.122	0.035	11.33	62.60	5.55
18	Kandi Local 3	0.099	0.043	11.39	60.57	4.03
19	Kota maize yellow	0.134	0.036	10.38	66.64	4.11
20	Local 3	0.134	0.038	11.53	58.53	6.24
21	More 110	0.083	0.043	10.55	68.29	1.28
22	Nadaun selection	0.031	0.034	10.50	67.62	4.91
23	Patti Kangra HP	0.110	0.043	11.80	62.28	5.66
24	Pearl Popcorn	0.076	0.037	11.50	62.52	5.91
25	Kailer Solan	0.077	0.033	10.75	69.63	3.42

S.No.	Genotype	Methionine (%)	Tryptophan (%)	Protein (%)	Starch (%)	β-carotene (%)
26	White op2 composite	0.065	0.050	11.95	60.38	0.96
27	Yellow op2 composite	0.086	0.052	10.58	64.32	4.80
	Mean	0.095	0.038	11.03	62.48	3.56
	LSD	0.003	0.001	0.196	0.457	0.062
	CV (%)	1.83	2.49	1.26	0.51	1.23
	Max	0.170	0.052	11.95	69.62	6.24
	Min	0.031	0.033	10.325	55.48	0.79

Supplementary Table 3: Inter- cluster distances among the different clusters of test material

	Cluster 1	Cluster 2	Cluster 3	Cluster 4	Cluster 5
Cluster1	0				
Cluster2	2.59	0			
Cluster3	2.84	1.63	0		
Cluster4	2.49	3.15	3.33	0	
Cluster5	2.46	2.06	2.41	2.657	0

BOOK REVIEW

A Review of *Code Paroda: A Life in Service of Agricultural Science*

Code Paroda. Sanjay Kumar and Arun Tiwari. A memoir of Dr RS Paroda as narrated to the authors. 422 pages.

Code Paroda, narrated by Dr Rajendra Singh Paroda to authors Sanjay Kumar and Arun Tiwari, is at once a personal memoir, an institutional history, and a sustained meditation on the purpose and practice of public agricultural science in India. Organised in four parts — “Making of the Man,” At the Helm, “Ascend,” and “Legacy” — the book covers, across 422 pages, a professional life that began in the agrarian landscape of Rajasthan in 1942 and has engaged with nearly every significant institution and policy development in Indian agricultural research across seven decades. It is a book that will reward readers at multiple levels: agricultural scientists and students will find in it both a career model and a map of the research ecosystem; historians of science and policy scholars will find rich primary material on institutional decision-making across a formative period; and general readers with an interest in India's development trajectory will find it an

illuminating account of how a postcolonial nation built, tested, and gradually transformed its scientific infrastructure for food and agriculture.

The book's great strength is its density of concrete detail. Names, dates, projects, partnerships, and policy moments are woven together with the authority of first-person recollection, offering an account of institutional life that no external history could fully reconstruct. The narrative is candid about personal adversity, generous toward collaborators, honest about institutional limitation, and consistently grounded in a conviction that science in the service of farmers and rural communities is both a moral vocation and a technical one. The result is a text that reads, at different moments, as biography, administrative memoir, and professional testament — a combination that is unusual in Indian scientific literature and valuable for precisely that reason.

PART I — MAKING OF THE MAN: FORMATION, EDUCATION, AND EARLY RESEARCH

Dr Paroda was born on 28 August 1942 into a moderately well-off farming family in Ajmer, Rajasthan. The opening section of the book traces the formation of a scientific temperament through a social and educational environment that valued curiosity, discipline, and ethical conduct. His primary school teachers created an environment in which learning was a source of genuine joy, and cultivated in their students honesty, respect, responsibility, and perseverance, as well as the capacity to cooperate, empathise, acknowledge differences, and resolve conflict constructively. Rajasthan's cultural landscape, saturated with Bhakti and Sufi traditions, reinforced values of humility and service that recur throughout the professional life the book describes.

By Class 10, he had resolved to study agriculture. His father's counsel at this juncture was important: rather than directing his son toward the nearest agricultural institution, he encouraged him to aspire more widely,

citing the Agricultural College, Pune, as a worthy destination. Dr Paroda ultimately enrolled in B.Sc. (Agricultural Sciences) at Dayanand Anglo-Vedic (DAV) College, Ajmer — joining after the close of admissions, a detail the book offers as an early illustration of determination — and was elected President of the Agricultural Students Union. He subsequently pursued M.Sc. (Agriculture) at Rajasthan College of Agriculture, Udaipur, in 1962, shifting from agronomy to plant breeding and genetics, a disciplinary reorientation that would define his research contributions for decades. He topped his class and received a gold medal from Union Minister Morarji Desai in 1964.

He was selected for a PhD in Genetics at IARI, where Dr MS Swaminathan headed the division, and completed his doctoral work under the guidance of Dr A.B. Joshi on wheat genetics — at the precise historical moment when India's Green Revolution was gathering force. The intellectual climate at IARI during this period,

fostered by a remarkable concentration of scientific talent, is described as “a fertile ground for nurturing talent with ripple effects” whose consequences for Indian agricultural science can scarcely be overstated. His mother's encouragement, after the PhD, to go further rather than rest is among the book's more affecting passages. He was awarded a Commonwealth Scholarship in August 1968 and proceeded to the University of Wales under Prof Hugh Rees to work on barley, broadening his research experience and his understanding of international agricultural science.

Returning to India, he joined Haryana Agricultural University (HAU) in January 1971 as an Associate Professor, leading a team of twelve scientists. At HAU, he identified a systemic consequence of the Green Revolution that had received insufficient attention: the widespread adoption of high-yielding varieties had reduced fodder availability, as inedible stalks replaced traditional straw and pasture lands contracted. His group responded by developing more than ten varieties and hybrids of forage crops over seven years — an illustration of systems-level thinking that would characterise his approach to research problems throughout his career. In 1977, he was appointed Project Coordinator (Forage Crops) at ICAR Headquarters, Krishi Bhawan, at the initiative of Dr Bhumbla, DDG (Natural Resource Management), and was subsequently selected as Head, Division of Plant Studies, CAZRI, Jodhpur.

The death of his first wife, leaving two young daughters, is narrated with characteristic restraint. To cope with this tragedy, he moved to HAU as Head of the Department of Plant Breeding, where he cultivated a vibrant academic culture: recruiting beyond established networks, publishing actively, securing research funding, and extending technology transfer to resource-poor areas. He founded the Indian Society of Forage Research in 1979, married Dr Shashi Tomar in 1979, and was blessed with a son on 30 July 1980. His retrospective reflection — that those years were defined not only by scientific work but by perseverance as a form of professional and personal discipline — runs as an animating thread through the rest of the book.

The section pays generous tribute to Dr Harbhajan Singh, Head of the Plant Introduction Division, IARI and described here as the “Vavilov of India,” whose example inspired Dr Paroda's long commitment to building genetic resource infrastructure. After his appointment as Director, NBPGR, Dr Paroda approached Dr Randhawa, DG of ICAR, for institutional support; receiving assurance of backing except for land and funds, he pursued an international partnership. The

outcome was the Indo-USAID Plant Genetic Resources Project, valued at \$21.66 million, through which 100 NBPGR staff received advanced training in the United States — a capacity-building initiative whose scale set a benchmark for international collaboration in Indian agricultural science.

As DDG (Crop Science) from 1987, Dr Paroda's capacity to translate scientific insight into policy found clear expression in the New Seed Development Policy of 1988. Grounded in self-reliance and marking the transition beyond Green Revolution-era variety dependency, the policy bore tangible results within five years: India produced its first hybrid rice after China, its first single-cross maize hybrid, and the first hybrids of safflower, castor, and pigeonpea. The National Research Centre for Plant Biotechnology (NRCPB), created in 1985 under Dr V.L. Chopra, had meanwhile begun advancing molecular biology and biotechnological research. The Technology Mission on Oilseeds (TMO), launched in 1986, provided a critical institutional bridge between biotechnology, breeding, and policy. As head of TMO from 1988 to 1994, Dr Paroda oversaw a transformation from importing approximately 60 per cent of edible oils to near self-sufficiency. Contributions through an expert committee under the Planning Commission increased pulses production by 4 million tonnes, significantly reducing imports. Along with Dr MS Swaminathan and Dr NS Randhawa, he was a founding member of the National Academy of Agricultural Sciences (NAAS) in 1990.

This period overlapped with the macro-policy disruptions of the early 1990s. Liberalisation under Prime Minister PV Narasimha Rao and Finance Minister Manmohan Singh, while consolidating food grain self-sufficiency, exposed agriculture to new market pressures through the removal of export barriers and subsidy rationalisation. Balram Jakhar served as Union Agricultural Minister through this transition. It was a context that demanded of scientific institutions both ideological flexibility and structural coherence. When the ICAR Director Generalship was awarded to Dr VL Chopra — the book acknowledges this openly — Dr Paroda accepted an invitation from Dr RB Singh to join FAO in Bangkok as Regional Plant Production and Protection Officer. His observations of Southeast Asian agricultural systems — their integration of rice, water management, and horticultural production within frameworks of moderation and self-sufficiency — enriched a perspective hitherto anchored mainly in South Asian priorities. APAARI, established in 1990, became an important institutional home: Dr RB Singh served as its first Executive Secretary; Dr Paroda

followed as the next. Encounters during these years with Dr Wang Lianzheng (China), Dr Zafar Altaf (Pakistan), and Dr William Dar (Philippines) — the last distinguished by his advocacy of participatory and

servant-leader approaches to science management — broadened his conception of what agricultural leadership could look like across diverse political and cultural settings.

PART II — AT THE HELM: THE ICAR DIRECTOR GENERALSHIP (1992–1998)

Part II is the book's most analytically substantial section, and the one of greatest relevance to readers interested in the governance and direction of national agricultural research systems. It opens with an epigraph that sets its register: "When the sower becomes the steward, the harvest is not only of grain but of the futures — nurtured through vision, tested by fire and made abundant by science." Dr Paroda assumed charge as Director General of ICAR in 1992 and served until 1998, years coinciding with India's economic liberalisation and a fundamental reorientation of its agricultural ambitions.

His first act on taking charge was to listen rather than direct — extensive consultation with colleagues and seniors about institutional priorities. The philosophy he articulates is explicit and instructive: the circle of concern is wide, but proactive leadership must concentrate on the circle of influence; responsibilities must be communicated honestly and boundaries respected. Vision 2020 for ICAR was developed not as a headquarters prescription but through genuine collaboration with all SAUs and ICAR institutions across a 25-year planning horizon. Consultation drew in Dr Uma Lele (economist), Dr YK Alagh (VC, JNU), Dr KL Chadha, Dr EA Siddique, and past DGs of ICAR, reflecting a commitment to management by consultation that defined the tenure.

The institutional infrastructure created during these seven years was substantial. New ICAR establishments included specialised bureaus for animal, fish, microbial, and insect genetic resources; a National Gene Bank; a National Agricultural Research Centre; National Agricultural Science Complex (NASC) complex; and regional offices of CGIAR centres in India. The 2nd International Crop Science Congress was co-organised alongside Dr MS Swaminathan in 1996. The Asia and Pacific Seed Association (APSA) was initiated in 1994. Research outcomes of note included Dr Siddique's leadership in developing Pusa RH-10 (Sugandhit Dhan, a fragrant rice hybrid), and the contributions of Dr Vasal and Dr NN Singh to Quality Protein Maize — work the book uses to illuminate the genetic mechanism of heterosis, whereby hybrids

activate hundreds of genes silent in either parent, increasing both the diversity and quantity of active gene expression.

A significant policy decision of this period was the opening of ICAR's public seed lines to private sector collaboration: parental lines were shared with domestic companies, and foreign companies were permitted to form joint ventures. This represented a considered response to the growing role of private investment in vegetable and fruit hybrid development, and it was calibrated to maintain public-interest breeding alongside private sector dynamism. Krishi Anusandhan Bhawan-II was planned as a centrally air-conditioned facility. The National Phytotron Facility (NPF) at IARI was established to enable controlled-environment research. India's accession to the WTO in 1995 created new imperatives for export-oriented agricultural development to which the research system was required to respond.

Administrative innovations during this tenure deserve equal attention. A transparent, merit-based incentive system was established to recognise research, teaching, and extension — an alignment of individual incentives with institutional mission that remains a benchmark in public research management. Structured, regular consultation was instituted among CSIR, DRDO, DBT, and ICAR to foster collaboration and reduce duplication. He chaired the Global Forum on Agricultural Research (1998–2002) within FAO and Executive Secretary, Asia Pacific Association of Agricultural Research Institutions (APAARI), 1993–2014. The National Agricultural Technology Project (NATP) was launched in 1998, and the Padma Bhushan was conferred in the same year. The book's observation on institutional quality is direct and bears quoting:

"Institutions are fundamental to a country's development and stability; however, if institutions become corrupt, outdated, or weak, they may hinder progress and lead to instability."

Coming from a scientist who spent a career within these structures, this is not an observation of abstract pessimism but of clear-eyed, reformist conviction.

ADVERSITY AND PROFESSIONAL CONDUCT: A STOIC INTERLUDE

In 2001, despite a record of sustained achievement, Dr Paroda was placed “on compulsory wait till further orders” on the basis of what he characterises as unfounded allegations of irregularities in computer procurement. The Central Joint Staff Council (CJSC), the Agricultural Research Scientists Forum (ARSSF), and large sections of the agricultural fraternity criticised the government's decision. He was ultimately vindicated, with all allegations found to be without merit. The Agricultural Minister requested him to continue as DG, ICAR. He declined and sought voluntary retirement with two years of service remaining.

The book's account of this episode is one of its most instructive passages. Turning to Stoic philosophy, Dr Paroda articulates a discipline of response that refuses to convert temporary adversity into permanent bitterness: to avoid consequential decisions under duress, to wait with patience for clarity, and to redirect energy toward constructive purpose. His wife's characterisation — that small minds fixate on petty

allegations — captures the disposition the book advocates throughout: proceed, build, contribute. It is a model of professional conduct under institutional pressure that scientists and administrators at any career stage will find worth considering.

Even through this period, substantive work continued. The 88th Indian Science Congress, held at IARI from 3 to 6 January 2001 and attended by some 5,000 scientists, produced a Vision 2020 statement drafted collectively by the presidents of all academic societies: “By 2020, India will be free of poverty, hunger, and malnutrition and become environmentally safe.” Dr APJ Abdul Kalam, chief guest, articulated national targets of 9–11 per cent economic growth, negligible poverty, and a rise to the world's fourth-largest economy. Simultaneously, Dr Paroda was pursuing consensus on the draft Protection of Plant Varieties and Farmers' Rights Act (PPV&FRA). On 7 January 2001, the idea of the Trust for Advancement of Agricultural Sciences (TAAS) crystallised — an initiative that would become, in the years that followed, one of his most enduring contributions.

PART III — ASCEND: INTERNATIONAL ENGAGEMENTS AND POST-ICAR CONTRIBUTIONS

Part III, subtitled “Knowledge ripens through struggle,” documents what is better described as a second career than a post-retirement chapter. Its philosophical preoccupation — the value of acting with integrity rather than assertion, of navigating complexity with humility — gives coherence to a set of activities that span institutional development, international research collaboration, science-policy engagement, and the cultivation of young talent.

TAAS took shape as what the book calls “a slow, almost organic synthesis — shaped by dialogue, reflection, and shared conviction.” Dr Nagarajan, then Director of IARI, provided land; Dr NN Singh became founding secretary. TAAS instituted the Dr MS Swaminathan Award for Leadership in Agriculture in 2004, followed by the Dr SK Vasal Award for Excellence in Crop Hybrid Research and the Dr Norman E Borlaug Innovative Farmer Award. Sustained support from Dr SM Sehgal, Dr BR Barwale, NSAI, APAARI, select SAUs, and CGIAR Centres provided the institutional base for these activities.

Dr Paroda led a World Bank-funded ecoregional programme for sustainable agricultural development in Central Asia and the Caucasus under ICARDA (2001–

2007). The programme's principal components — no-till practices in wheat-fallow monocultures, the CERES-Wheat simulation model (Crop Estimation through Resource and Environment Synthesis), and the Kazakhstan–Siberia Network for Spring Wheat Improvement (KASIB) — contributed measurably to the development of climate-resilient varieties with improved yield, quality, and disease resistance. His approach throughout these engagements was consistent: work alongside local scientists and institutions, build bridges for scientific collaboration rather than transplant solutions. The Global Forum for Agricultural Research (GFAR), established and governed by a steering committee rather than a single chairperson, institutionalised this collaborative logic at the global level. Contributions to the rice–wheat consortium network spanning Bangladesh, India, Pakistan, and Nepal extended it across South Asian agro-ecologies with significant documented impact.

On biotechnology, the book is analytically careful and politically candid. Dr Paroda commends the foundational contributions of Dr Manju Sharma and Dr Gargi Vachaknavi to India's biotechnology institutions and regulatory frameworks. He argues that if genetic

modification could rescue a fibre crop from systemic collapse, it could equally address India's long-standing vulnerability in food oils and protein. His observation that a technology developed in Indian laboratories was adopted in Bangladesh while denied to Indian farmers encapsulates a regulatory and policy paradox whose resolution remains a live debate. His position is clearly stated: regulation should be the intelligent enabler of progress — transparent, fair, and timely — rather than its chronic inhibitor. He calls for an evidence-based, proactive approach to GM technologies: examine, test, adapt, and implement responsibly. The structural disconnect between scientific progress and systemic response, and the consequent need for honest communication between research and policy communities, are themes whose relevance extends well beyond the biotechnology debate.

Twenty-two universities conferred honorary doctorates upon him — recognitions he regards not as personal accolades but as affirmations of a mode of practice: standing by farmers, by evidence, and by the long-term arc of scientific accountability. In May 2024, Dr SK Vasal delivered a special lecture at TAAS illustrating that important agricultural traits — nutrition, stress tolerance, yield — rarely travel alone, being governed by many genes in complex interaction. He reminded the audience that scientific leadership must give equal priority to building infrastructure and to cultivating ideas. Agricultural science is characterized by scientists who choose the long, uncertain path of discovery as “quiet warriors of light” — working without spectacle, often without assurance of recognition, yet persisting in illuminating natural processes so that society moves incrementally toward greater food security.

SCIENCE–POLICY INTERFACE: DIALOGUES, DECLARATIONS, AND INSTITUTIONAL ADVOCACY

The years from 2001 reveal a sustained pattern of structured scientific advocacy: convening multidisciplinary groups around specific policy challenges, producing evidence-based outputs, and pursuing their incorporation into government policy and programmes. In 2005, TAAS held a two-day brainstorming session on the role of science and society in plant genetic resources management. In 2009, a workshop on emerging challenges in Indian agriculture was held in association with IFPRI; a subsequent workshop on farm animal genetic resources conservation culminated in the Ranchi Declaration, hosted at Birsa Agricultural University. In 2010, a national dialogue on building leadership in agricultural

research management was organised with NAARM. In 2015, deliberations on soil health and nutrition management fed directly into the Government of India's Soil Health Card Scheme, launched in 2015–16. In 2016, the International Agrobiodiversity Congress, co-chaired by Dr Paroda and Dr MS Swaminathan, produced the Delhi Declaration on Agrobiodiversity Management. In 2023, a workshop on climate change, soil quality, and food security — chaired by Dr MS Swaminathan at the NASC Complex — reached a collective conclusion that without conservation agriculture, climate-resilient varieties, and efficient water and nutrient management, food security itself would be placed at risk. The book records in some detail the substantive conversations among participants at each of these events, providing a valuable record of science–policy interface in practice.

The Global Conference on Agricultural Research and Development at Montpellier in 2010, where Dr Paroda chaired the Technical Programme Committee and ensured participation of youth and women from over 100 countries, gave rise to YPARD (Youth Platform for Agricultural Research and Development). His subsequent work as chair of the Haryana Farmers Commission (2010–14) reinforced a conclusion the book states with directness: agricultural curricula regularly fail to keep pace with technological change, evolving markets, and the skills demanded by contemporary farming and food systems. Agriculture must be repositioned, he argues, not as a single occupation but as an ecosystem of opportunity — one capable of producing a generation of agripreneurs. The observation that young people do not reject agriculture but reject stagnation is one of the book's most incisive and actionable insights.

Regional platforms — APCoAB, APARIS, GFAR, APAARI — are cited as essential architecture for translating scientific knowledge across agro-ecologies and political boundaries. At the domestic level, Krishi Vigyan Kendras (KVKs) should function not as peripheral outposts of research institutions but as genuine nerve centres of rural knowledge systems: hubs of innovation, skill development, and rural enterprise, organically connected to village realities. The concept of bio-digital resource centres — integrating genetic data, breeding programmes, and farmer networks — is proposed as the appropriate institutional form for the next generation of agricultural science.

On the political economy of Indian agriculture, the book is frank. Citing former World Bank Vice President Ismail Serageldin, Dr Paroda warns that without serious

attention to structural inequality, India risks bifurcating into “half California and half sub-Saharan Africa.” Food security without rural prosperity is fragile; rural prosperity without sustainable food systems is short-lived. On farm policy, he argues that the shift from subsidies to incentives must be accompanied by genuine and sustained engagement with farmers' voices. Strengthening public-sector research is essential not to displace private investment but to

ensure that science in the public interest — including socioeconomic assessment, long-term adoption studies, and farmer surveys — remains central to the national agricultural system. In unfavourable environments, he notes, progress depends not only on technological innovation but on institutional support and policies that recognise risk as a structural companion of resource-poor farming.

PART IV — LEGACY: PHILOSOPHY, PEDAGOGY, AND THE CODE PARODA

The book's final section offers two contributions of distinct character. The chapter “For Whom the Bell Tolls” distils personal philosophy from seven decades of practice. Drawing on the *Bhagavad Gita*, Dr Paroda identifies three governing principles: the duty to perform one's own prescribed role faithfully, however imperfectly, rather than another's; the discipline of performing that duty without attachment to reward or recognition; and the commitment to act with courage in the face of uncertainty and opposition. These are not novel formulations, but they carry weight commensurate with the life that has enacted them consistently.

The Code Paroda — ten codes covering Climate, Science, Markets, Commons, Nutrition, Women, Youth, Digital Irrigation, Collectives, and Public Science Deeds — is offered as a framework for the next generation of agricultural scientists, policymakers, and practitioners. Its treatment in the book is necessarily summary, but as a declaration of priorities it is a substantive contribution to discussions currently underway about the direction of Indian agricultural research, extension, and policy. The emphasis on digital agriculture, collective action, and the meaningful inclusion of women and youth reflects the institutional learning of an unusually wide career and points toward the kinds of governance and participation that a transformed agricultural system will require.

“Any contribution I made lies in helping to create an environment in which scientists felt supported, connected, and valued within a broader scientific community.”

This self-assessment — measured, free of self-aggrandisement — is characteristic of the book's tone throughout and points to what the reviewer takes to be its principal implicit argument: that the quality of scientific leadership is constituted less by individual brilliance than by the capacity to build environments in which others can do excellent work. Strong institutions,

the book consistently argues, are shaped by individuals who combine scholarship with courage, and national commitment with global values.

ASSESSMENT

Code Paroda makes a significant contribution to the literature on Indian agricultural science, its institutional history, and its science-policy interface. Its most important quality is evidentiary: through first-person narration, it provides a detailed and authoritative account of how scientific decisions were made, how institutions were built, and how research knowledge was — and in certain important cases was not — translated into policy across a period of decisive significance for Indian agriculture. No external account of this period can substitute for the granular institutional detail this book provides, and no serious student of India's agricultural research system should overlook it.

The book's analytical engagement with biotechnology regulation is among its more consequential contributions. The argument that India's regulatory architecture has been excessively cautious in ways that have disadvantaged Indian farmers relative to those in neighbouring countries is stated with evidence and restrained force. It deserves engagement from policymakers, regulators, and researchers. Similarly, the book's sustained argument for a strengthened public-sector research mandate — not as a constraint on private investment but as its necessary complement — addresses a structural question that no credible agricultural policy framework can avoid.

The book's treatment of institution-building — NAAS, APSA, TAAS, GFAR, APAARI, and the thematic dialogues that fed into national schemes such as the Soil Health Card — provides what is effectively a practitioner's guide to translating scientific knowledge into durable institutional form and public policy. The lesson drawn is consistent: genuine consultation, long-term commitment, and the willingness to work through

established and new partnerships are more productive than hierarchical directive or individual initiative alone.

For agricultural scientists and students at every career stage, the book offers something genuinely uncommon in Indian scientific literature: a full account of how technical competence, institutional leadership, and ethical conviction can be sustained across a long and varied career. For researchers in science policy, agricultural economics, and development studies, it provides rich primary material on a period — roughly 1965 to the present — that transformed India's relationship with its own agricultural potential. And for policymakers and practitioners, it offers both a record of what has been achieved through sustained scientific investment and a framework, in the Code Paroda itself, for the work that remains.

The final injunction — to serve without condition, to persist without spectacle, and to trust that sincere

effort, anchored in ethical responsibility, finds its own direction — is the distillate of a career rather than a rhetorical conclusion. It is, in this sense, the most credible kind of advice: advice that has been lived before it has been written.

Reviewed by:



A Amarender Reddy
Joint Director
ICAR-NIBSM, Raipur
e-mail: amarender.reddy@icar.org.in

Plant Germplasm Registration Notice *

180 proposals were received online for the consideration of registration of germplasm for unique/superior traits. The applications were thoroughly screened by the member-secretary, Plant Germplasm Registration Committee (PGRC), and 94 proposals complete in all respects were reviewed by the experts and were presented in the 50th meeting of the PGRC held in virtual mode on June 12, 2023. 67 proposals with unique/novel features belonging to 29 species were recommended for registration by PGRC. The information on registered germplasm is published to disseminate the information to respective crop breeders for utilization of these genetic stocks in their crop improvement programmes. Description of 34 out of the 67 recommended germplasm lines was published in IJGR Vol 39 (Issue no. 1) 2026. Description of the remaining 33 germplasm lines is given below:

1. **WN 657 (IC648611; INGR23035), a finger millet germplasm with longer finger length as per DUS test**

HK Patil^{1*}, M Elangovan², GD Vadodariya¹, Vikas Pali³, S Pandey⁴, CD Pandey⁴ and M Ramani³

¹HMRS, Navsari Agricultural University, Dangs-386151, Gujarat, India

²ICAR-Indian Institute of Millets Research, Hyderabad-500030, Telangana, India

³ARS, Anand Agricultural University, Dahod-389160, Gujarat, India

⁴ICAR-National Bureau of Plant Genetic Resources, Pusa Campus-110012, New Delhi, India

*Email: harshalpatil@nau.in

A total of 38 finger millet [*Eleusine coracana* (L.) Gaertn.] accessions constituted of 25 promising long earhead landraces and thirteen released varieties were evaluated for 12 morphological characters, including grain yield, at three locations in Gujarat, during *kharif* 2019, 2020 and 2021. The pooled data of all locations were used to study the genetic potential of long earhead finger millet genotypes. Moderate genotypic and phenotypic coefficient of variation found for the traits viz., number of fingers per earhead, number of productive tillers per plant, straw yield per plant, grain yield per plant, finger length, harvest index and main earhead indicating ample scope of variation for these traits, allowing further improvement by selection of these traits. High heritability estimates were observed for days to 50% flowering, days to maturity, plant height, number of productive tillers per plant, number of fingers per earhead, main earhead length, finger length, 1000-grain

weight, grain yield per plant, straw yield per plant and harvest index, showing low environmental influence on these traits and presence of additive gene action for these traits. Hence, priority can be given to these traits during selection to get more genetic gains.

Also, on the basis of mean performance over three years at two locations, the genotype 'WN 657' showed long finger length, i.e. 13.13 cm, which is >7 cm (extra-long) as per DUS testing guidelines. This morphological trait, such as the number of fingers per earhead, the main earhead length and the length of finger has direct contributions towards the high grain yield per plant. The best promising finger millet genotypes 'WN-657' for very long finger length with moderate resistance to pests and diseases found during three years of genetic evaluation at three different locations, which can be used as a promising parent in further breeding programmes.

2. **SPV 2813 (IC648612; INGR23036), a sorghum germplasm with longer leaf length and wider leaf width**

AV Umakanth*, Amasiddha B and K Hariprasanna

ICAR-Indian Institute of Millets Research, Hyderabad-500030, Telangana, India

*Email: umakanth@millets.res.in

Green fodder yield in sorghum (*Sorghum bicolor* L.) has a significant and strong positive correlation with leaf

area followed by stem girth, plant height, leaf length and number of leaves per plant (Singh *et al.*, 2019).

*Edited by: **Anju Mahendru Singh and Arup Das**, ICAR-National Bureau of Plant Genetic Resources, Pusa Campus, New Delhi-110012, India.

Positive and significant association of number of leaves per plant, leaf breadth and dry fodder yield with green fodder yield indicate the importance of these characters during selection for crop improvement program in forage sorghum (Gedam *et al.*, 2015). The significance of brown midrib mutation in sorghum is that it greatly reduces lignification and cell-wall concentration, increases digestibility, and increases voluntary intake of feed by ruminants. Genetic control of the lignification process through use of brown midrib (*bmr*) mutations has offered the most direct and productive approach to reducing lignin concentration and increasing digestibility of sorghum.

SPV 2813 was developed through pedigree method of breeding by crossing between CSV 20 and the *bmr* germplasm line, IS 21891 during *rabi* 2009-10 at the Indian Institute of Millets Research, Hyderabad. CSV 20 is the most popular dual-purpose sorghum variety which was released during 2009 for cultivation in zones I and II in India, while IS 21891 is a brown midrib source carrying the gene *bmr* 8. Agronomically superior derivatives from this cross were selected and

advancement of segregating generations was done.

SPV 2813 is a brown midrib sorghum line with tan pigmentation and days to flower of 83 days. This unique line is superior for longer leaf length (83.1 cm) and longer leaf width (8.39 cm) (Table 1). These identified traits add to the increased forage value of the variety apart from making it more suitable for bio-ethanol production because of increased lignocellulose contents. SPV 2813 is having green fodder yield of 315.6 q/ha and a dry fodder yield of 94.9 q/ha. The other traits like stem girth (4.69 cm), protein (9.03%), total soluble sugars (10.95%) and number of leaves (13) also increase the forage value of this entry.

References

- Gedam SR, DT Deshmukh, MA Sheikh, RB Ghorade and SB Sakhare (2015) Genetic studies on forage sorghum (*Sorghum bicolor* (L.) Moench). *BIOINFOLET-A Quarterly J. Life Sci.* 12: 958-962.
- Singh J, BR Ranwah, L Chaudhary, C Lal, MC Dagla and V Kumar V (2013) Evaluation for genetic variability, correlation and path coefficient analysis in mutant population of forage (*Sorghum bicolor* L.). *Bioscan* 8: 1471-76.

Table 1: Summary data on leaf length and leaf width traits, BMR trial *Kharif* 2020 (AICRP data)

Trait	Location	SPV 2813	CSV 43BMR (C)	CD 5%
Leaf length (cm)	8	83.1	76.8	4.9
Leaf width (cm)	7	8.39	7.81	0.78

3. SPV 2058 (IC648613; INGR23037), a sorghum germplasm with longer leaf length, leaf width and high stem

AV Umakanth*, B Amasiddha and KBRS Visarada

ICAR-Indian Institute of Millets Research, Hyderabad-500030, Telangana, India

*Email: umakanth@millets.res.in

Sorghum (*Sorghum bicolor* L.) ranks first among the cereal fodder crops because of its growing ability in low-fertility soil, faster growth habit, higher yield, palatability and nutritious quality. It is preferred over maize and pearl millet for forage because of its better regeneration capacity. Green fodder yield in sorghum has a significant and strong positive correlation with leaf area followed by stem girth, plant height, leaf length and number of leaves per plant (Singh *et al.*, 2019). SPV 2058 is a single-cut forage sorghum genotype with longer leaf length, higher leaf width, higher stem girth and better leaf-stem ratio which are the important characters in an ideal forage sorghum variety. It was developed through pedigree method of breeding by crossing between PC 5 and Indore 12 during *rabi* 2004-

05 at ICAR-Indian Institute of Millets Research, Hyderabad.

SPV 2058 is a tan type plant with good early vigour, stems are thicker with longer leaves and leaf width is also wider conferring more biomass to yield good forage quantity. The forage quality of the line is also good in relation to protein content (8%) and higher IVDMD (51%) values. The green fodder yield (430 q/ha) and dry fodder yield (141.5 q/ha) are superior to the existing check varieties along with higher per day productivity of green fodder yield (5.65 q/ha) and dry fodder yield (1.85 q/ha). SPV 2058 has multiple trait superiority with uniqueness of longer leaf length (85.5 cm), leaf width (7.32 cm), high stem girth (3.07 cm) and leaf: stem ratio (0.41) (Table 1). It recorded the highest

leaf length, leaf breadth and is resistant to Anthracnose and zonate leaf spot. This variety is suitable to be grown as a single-cut forage sorghum variety for rainfed Kharif sorghum growing regions of the country and will be an excellent source of these traits as a donor parent in forage breeding.

Reference

Singh SK, LK Gangwar and Mayank Chaudhary (2019) Studies on character association and path analysis in forage sorghum. *Int. J. Chem. Stud.* 7: 2939-2942.

Table 1: Summary of data for leaf length, leaf width, stem girth and leaf: stem ratio traits of AVT-SC, *kharif* 2012 (AICRP data)

Trait	Genotypes	Zone I (9)	Zone II (6)	All India Avg.	CD 5%
Leaf length	SPV 2058	84.7	86.8	85.5	3.3
	HC 308 (c)	75.1	74	74.7	
	CSV 21F (c)	80.8	77	79.3	
Leaf width	SPV 2058	7.56	6.98	7.32	0.47
	HC 308 (c)	6.82	5.93	6.47	
	CSV 21F (c)	6.58	5.66	6.21	
Leaf: stem ratio	SPV 2058	0.37	0.46	0.41	0.04
	HC 308 (c)	0.28	0.45	0.36	
	CSV 21F (c)	0.3	0.47	0.39	

4. TAKPS 1 (IC648614; INGR23038), a hurda sorghum germplasm with free threshability, excellent fragrance and high hundred tender grain weight

RB Ghorade^{1*}, VV Kalpande¹, SB Thawari¹, A Badigannavar², Aruna C³, R Madhusudhana³, AR Bhuyar¹, PS Kamble¹ and SP Chikate¹

¹Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola-444104, Maharashtra, India

²Bhabha Atomic Research Centre, Trombay, Mumbai-400085, Maharashtra, India

³ICAR-Indian Institute of Millets Research, Hyderabad-500030, Telangana, India

*Email: akola@aicrmmillets.res.in

The special-purpose sorghum (*Sorghum bicolor* L.) can provide a choice to the end users in food enterprises. Green grains of parching sorghum (*Hurda* sorghum) from milk to dough stage are used for consumption as roasted grains. Parching sorghum possesses a pleasant fragrance and more sweetness which make them popular for table purposes. Parching (*Hurda*) sorghum is one of the best alternatives for the value addition of sorghum.

Parching sorghum genotype Sakarimokari was irradiated at BARC, Mumbai. This landrace was gamma irradiated with 300 Gy, using in Gamma chamber (1200 gx) at BARC, Mumbai in order to create genetic variation for the hurda quality parameters to improve hurda yield, hurda quality and earliness. The promising selection Sakarimokari mutant selection number-5-4-1-1 was advanced till M7 generation and finally named as TAKPS 1. Promising mutants along with checks were tested in University Multilocation Trials during Rabi-2018-19 to 2020-21. The sensory evaluation included

colour, size, shape, aroma/flavour, taste and thrashability (Ghorade *et al.*, 2019). The chemical analysis (total sugars and brix) of the hurda was executed according to the standard method of AOAC, 1990.

Free threshability is the prerequisite for the parching of the sorghum genotype. The threshability of TAKPS 1 was in the category of Free threshability while the checks like Gulbhendi, Sakrimokari and Phule Uttara were in the category of Medium threshability. While the check Madhur was also having Free threshability. This genotype recorded high hundred tender grain weight (4.12 g). For the organoleptic trait of Taste, the genotype TAKPS 1 was in the category of Very sweet while the checks like Gulbhendi, Sakrimokari and Phule Uttara were in the category of Medium sweet and Madhur wani was Slightly sweet. For the trait of Fragrance, the genotype TAKPS 1 was in the category of Good while the checks like Gulbhendi and Sakrimokari were in the category of Fair and Phule Uttara and Mahur were in the category of No Aroma.

TAKPS 1 also recorded higher sugar (2.17%) as compared to the checks Gulbhendi (1.96%), Sakrimokari (2.0%), Madhur Wani (2.03% q/ha) and Phule Uttara (1.86% q/ha).

Earliness is a desirable trait in parching sorghum. For days to dough stage, TAKPS 1 recorded fewer days to dough stage of 94.17 days as compared to the checks Gulbhendi (95.17 days), Sakrimokari (102.25 days), Madhur Wani (95.67 days) and Phule Uttara (95.75 days).

Thus, it was concluded from the present study that considering the higher green hurda yield, early maturity along with good threshability and excellent

organoleptic properties, the promising genotype TAKPS 1 needs to be exploited in breeding for improvement of parching traits in rabi sorghum as the donor parent in the crossing programme.

Reference

Ghorade RB, SB Thawari, VV Kalpande, SV Khambalkar and AR Bhuyar (2019) PDKV Hurda Kartiki (WANI 103) a new parching kharif sorghum variety; innovative approach for rural entrepreneurship and empowerment. *J. Multilogic Sci.* Vol. VIII, Special Issue(B): 149-153.

5. TAKPS 3 (IC647559; INGR23039), a hurda sorghum germplasm with earliness, free threshability and excellent fragrance

RB Ghorade^{1*}, VV Kalpande¹, SB Thawari¹, A Badigannavar², Aruna C³, R Madhusudhana³, AR Bhuyar¹, PS Kamble¹ and SP Chikate¹

¹Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola-444104, Maharashtra, India

²Bhabha Atomic Research Centre, Trombay, Mumbai-400085, Maharashtra, India

³ICAR-Indian Institute of Millets Research, Hyderabad-500030, Telangana, India

*Email: akola@aicrpmillets.res.in

The special purpose sorghum (*Sorghum bicolor* L.) can provide choice to the end users in food enterprises. Green grains of parching sorghum (*Hurda* sorghum) from milk to dough stage are used for consumption as roasted grains. Parching sorghum possesses a pleasant fragrance and more sweetness which make them popular for table purpose varieties. It was therefore felt to develop and identify the genotype for hurda purpose of which will give benefit to farmers and consumers too.

The parching sorghum genotype Gulbhendi was irradiated at BARC, Mumbai. This landrace was gamma irradiated with 300 Gy, using in Gamma chamber (1200 gx) at BARC, Mumbai in order to create genetic variation for the hurda quality parameters to improve hurda yield, hurda quality and earliness. The promising selection Gulbhendi Mutant 300 Gy Selection No. -22-2-1-1 was advanced till M7 generation and finally named as TAKPS 3. Promising mutants along with checks were tested in University Multilocation Trials during Rabi-2018-19 to 2020-21. The sensory evaluation included colour, size, shape, aroma/flavour, taste and thrashability (Ghorade *et al.*, 2019). The chemical analysis (total sugars and brix) of the hurda was executed according to the standard method of AOAC, 1990.

For the character green hurda yield, the genotype TAKPS 3 recorded a higher green hurda yield of 40.40 q/ha as compared to the checks Gulbhendi (30.66 q/ha),

Sakrimokari (27.33 q/ha), Madhur Wani (32.51 q/ha) and Phule Uttara (31.64 q/ha). For the trait green fodder yield, the genotype TAKPS 3 recorded a higher green fodder yield of 113.17 q/ha.

Free threshability is the prerequisite for the parching of the sorghum genotype. The threshability of TAKPS 3 was found promising similar to the checks Gulbhendi, Sakrimokari and Phule Uttara.

For the organoleptic traits, TAKPS 3 was superior for almost all the traits. For the organoleptic trait of Taste, the genotype TAKPS 3 was in the category of higher sweetness, while the checks like Gulbhendi, Sakrimokari and Phule Uttara were in the category of Medium sweet and Madhur Wani was slightly sweet. For the trait of Fragrance, the genotype TAKPS 3 was in the category of Good while the checks like Gulbhendi and Sakrimokari were in the category of Fair and Phule Uttara and Mahur were in the category of without Aroma. Sugar content has major role in the sweetness of the parching sorghum. For the character total sugar %, TAKPS 3 recorded higher sugar of 1.99% as compared to the checks Gulbhendi (1.96%), Phule Uttara (1.86% q/ha) and was at par with Sakrimokari (2.0%), Madhur Wani (2.03% q/ha).

Earliness is a desirable trait in parching sorghum. For the character days to 50% flowering TAKPS 3 recorded less days to 50% flowering of 71.42 days as compared to the checks Gulbhendi (73.67 days), Sakrimokari (82.67 days), Mahur Wani (73.92 days) and

Phule Uttara (75.25 days). For days to dough stage, TAKPS 3 recorded less days to dough stage of 92.42 days as compared to the checks Gulbhendi (95.17 days), Sakrimokari (102.25 days), Madhur Wani (95.67 days) and Phule Uttara (95.75 days).

Thus, it was concluded from the present study that considering the higher green hurda yield, early maturity along with desirable threshability and excellent organoleptic properties, the promising

genotype TAKPS 3 needs to be exploited in breeding for improvement of parching traits in rabi sorghum as the donor parent in the crossing programme.

Reference

Ghorade RB, SB Thawari, VV Kalpande, SV Khambalkar and AR Bhuyar (2019) PDKV Hurda Kartiki (WANI 103) a new parching *kharif* sorghum variety; innovative approach for rural entrepreneurship and empowerment. *J. Multilogic in Sci.* Vol. VIII, Special Issue(B): 149-153.

6. AKSV 438 (IC286441; INGR23040), a sorghum germplasm with high zinc and iron content

RB Ghorade^{1*}, Elangovan M², S Pandey³, CD Pandey³, Hariprasanna K², VV Kalpande¹, AR Bhuyar¹, PS Kamble¹ and SB Thawari¹

¹Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola-444104, Maharashtra, India

²ICAR-Indian Institute of Millets Research, Hyderabad-500030, Telangana, India

³ICAR-National Bureau of Plant Genetic Resources, Pusa Campus-110012, New Delhi, India

*Email: akola@aicrmpillets.res.in

Sorghum (*Sorghum bicolor* L.) is one of the cheapest sources of energy and micronutrients, and a vast majority of population in Africa and central India depend on sorghum for their dietary energy and micronutrient requirements. Biofortification of sorghum by increasing mineral micronutrients [especially iron (Fe) and zinc (Zn)] in grain is of widespread interest (Ashok Kumar *et al.*, 2009). Considering this, the sorghum genotype AKSV 440 which was found promising for the higher Zinc (30.61 ppm) and Iron (37.14 ppm) content above the base line. As per the (Ashok Kumar *et al.*, 2013), the base line for Zinc (Zn) = 20 ppm and Iron (Fe) = 30 ppm has been fixed. Everything above the baseline is considered promising for Zinc and Iron content in sorghum.

The *kharif* sorghum genotype AKSV 438 contributed by the Sorghum Research Unit, Dr Panjabrao Deshmukh Krishi Vidyapeeth, Akola (Maharashtra) was analysed during Kharif 2018, 2019, 2020 and 2021. Determination of micronutrients in the sorghum grain sample was done by Di-Acid digestion method and the aliquots of this solution are used for the determination of Zinc and Iron with the help of Atomic Absorption Spectrophotometer (AAS), (Jackson *et al.*, 1967).

For the character Zinc (Zn), the genotype AKSV 438 had a mean value of 30.61 ppm, while the control had low values, *i.e.* PVK 801 (12.80 ppm) and Parbhani Shakti

(16.77 ppm). Similarly, for the character Iron (Fe), the genotype AKSV 438 had a mean value of 37.14 ppm, while the control had low values, *i.e.* PVK 801 (24.40 ppm) and Parbhani Shakti (24.05 ppm).

Besides high Zinc (Zn) and Iron (Fe), the genotype AKSV 438 is also high-yielding. AKSV 438 recorded 32.97 g of grain yield per panicle, while the checks PVK 801 (23.25 g) and Parbhani Shakti (26.61 g) were low-yielding.

Thus, it was concluded from the present study that considering the high zinc and iron along with the higher grain yield per panicle in the genotype AKSV 438, it needs to be exploited in breeding for development of genotypes with better biofortified traits in sorghum as the donor parent in the crossing programme.

References

- Ashok Kumar A, K Anuradha and B Ramaiah (2013) Increasing grain Fe and Zn concentration in sorghum; progress and way forward, International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru, Andhra Pradesh, India.
- Ashok Kumar A, BVS Reddy, B Ramaiah, P Sanjana Reddy, KL Sahrawat and HD Upadhyaya (2009) Genetic variability and plant character association of grain Fe and Zn in selected core collections of sorghum germplasm and breeding lines. *J. SAT Agric. Res.* 7: 1-4. (ejournal.icrisat.org).
- Jackson ML (1967) *Soil Chemical Analysis*. Prentice-Hall of India, New Delhi, India.

7. AKSV 440 (EC488403; INGR23041), a sorghum germplasm with high zinc and high iron content

RB Ghorade^{1*}, Hariprasanna K², VV Kalpande¹, AR Bhuyar¹, Elangovan M², S Pandey³, CD Pandey³, PS Kamble¹ and SB Thawari¹

¹Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola-444104, Maharashtra, India

²ICAR-Indian Institute of Millets Research, Hyderabad-500030, Telangana, India

³ICAR-National Bureau of Plant Genetic Resources, Pusa Campus-110012, New Delhi, India

*Email: akola@aicrmpmillets.res.in

Sorghum (*Sorghum bicolor* L.) is one of the cheapest sources of energy and micronutrients; and a vast majority of population in Africa and central India depend on sorghum for their dietary energy and micronutrient requirement. Biofortification of sorghum by increasing mineral micronutrients [especially iron (Fe) and zinc (Zn)] in grain is of widespread interest (Ashok Kumar *et al.*, 2009). Considering this, the sorghum genotype AKSV 440 which was found promising for the higher Zinc (29.78 ppm) and Iron (35.51 ppm) content above the base line. As per the (Ashok Kumar *et al.*, 2013) the base line for Zinc (Zn) = 20 ppm and Iron (Fe) = 30 ppm has been fixed. Everything above the base line is considered as promising for Zinc and Iron content in sorghum.

The *kharif* sorghum genotype AKSV 440 contributed by Sorghum Research Unit, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola (Maharashtra) was analyzed during *kharif* 2018, 2019, 2020 and 2021. The testing entries along with the controls PVK 801 and Parbhani Shakti were sown in randomized block design with three replications. Determination of micronutrients in the sorghum grain sample was done by Di-Acid digestion method and the aliquots of this solution are used for the determination of Zinc and Iron with the help of Atomic Absorption Spectrophotometer (AAS), (Jackson *et al.*, 1967).

For the character Zinc (Zn), the genotype AKSV 440 had a mean value of 29.78 ppm, while the control had low values, *i.e.* PVK 801 (12.80 ppm) and Parbhani Shakti (16.77 ppm). While for the character Iron (Fe), the genotype AKSV 440 with the mean value of 35.51 ppm, while the control was with low values, *i.e.* PVK 801 (24.40 ppm) and Parbhani Shakti (24.05 ppm).

Thus, it was concluded from the present study that considering the high zinc and iron in the genotype AKSV 440, it needs to be exploited in breeding for development of genotypes with better biofortified traits in sorghum as the donor parent in the crossing programme. Therefor this genotype needs to be registered with NBPGR as a donor source for enrichment of Zinc (Zn) and Iron (Fe) in *kharif* sorghum.

References

- Ashok Kumar A, K Anuradha and B Ramaiah (2013) Increasing grain Fe and Zn concentration in sorghum; progress and way forward, International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru, Andhra Pradesh, India.
- Ashok Kumar A, BVS Reddy, B Ramaiah, P Sanjana Reddy, KL Sahrawat and HD Upadhyaya (2009) Genetic variability and plant character association of grain Fe and Zn in selected core collections of sorghum germplasm and breeding lines. *J. SAT Agric. Res.* 7: 1-4. (ejournal.icrisat.org).
- Jackson ML (1967) Soil chemical analysis. Prentice-Hall of India, New Delhi, India.

8. SPV 2838 (IC643984; INGR23042), a sorghum germplasm with early flowering and early maturity line

R Madhusudhana*, Rajappa PV and VA Tonapi

ICAR-Indian Institute of Millets Research, Hyderabad-500030, Telangana, India

*E-mail: madhu@millets.res.in

Sorghum (*Sorghum bicolor* L.) is one of the most important C4 cereal crops cultivated in the arid and semi-arid tropics of the world for its nutritious grain, feed, and recently as a bioenergy crop. In India, sorghum is grown in two main seasons, *kharif* and *rabi*. The requirements for these two seasonal adaptations are quite diverse due to different agroclimatic

conditions. Currently, the area allocated to *kharif* and *rabi* sorghum in the total cropped area is approximately 20% (~1 m ha) and 80% (~3.5 m ha), respectively. Unlike the grain productivity of *kharif* sorghum (~1100 kg/ha), *rabi* sorghum is very low (~700 kg/ha). *Rabi* sorghum is grown in the vast Deccan plateau in residual and receding soil moisture regimes with increasing

temperatures and desiccating winds after flowering. Terminal drought stress is the most critical factor that affects the productivity of *rabi* sorghum grain, the quality of seed and feed.

Breeding for early flowering and maturity is an important strategy to overcome the ill effects of terminal drought. SPV2838 is a BC₃F₄ progeny, developed as an early-flowering and maturing entry in the background of CRS4 by backcross breeding. CRS4 is an important tall variety adapted to *rabi* developed at CRS-IIMR, Solapur, while B35 is a dwarf source of stay green. The new breeding stock SPV2838 is a tall, early maturing variety evaluated in the All India coordinated project on Sorghum during Rabi 2020 at 19 locations. SPV2838 flowers in 67 days and matures in 110 days compared to the check variety, M35-1, which flowers in 73 days and matures in 115 days. Early flowering and maturity by 6 days is extremely critical for the *rabi*

sorghum under terminal drought stress. M35-1 is the most popular cultivar, occupying 80% of the *rabi* sorghum area and is used as a standard check in the AICRP *rabi* sorghum trials. Therefore, the test entry SPV2838 has been compared against the check, M35-1. SPV2838 ranked first for early flowering at 13 locations out of 19 in AICRP trials at *rabi* locations. Its fodder is 9279 kg/ha with a grain test weight of 3.17 g/100 seeds. Its shoot fly reaction is 19.3% DH compared to M35-1 (23.9% DH). Its charcoal rot index is 16.34, similar to M35-1 (16.87).

SPV2838 can be cultivated in an equivalent way as of M35-1, a popular *rabi* sorghum variety with an advantage of early maturity of one week, which is very crucial in *rabi* season facing terminal drought. SPV2838, with its other matching traits, is extremely useful in the breeding program.

9. NDLA 3116-4 (IC646843; INGR23043), a cotton germplasm with high ginning per cent

YR Reddy^{1*}, BVRP Reddy², KM Vishnuvardhan², BRK Reddy², S Vasundhara², MS Rani¹, KVS Reddy¹, Rani Chapara¹, SJ Basha², DL Kalyani² and ASR Sharma²

¹Regional Agricultural Research Station, ANGRAU, Lam, Guntur-522034, Andhra Pradesh, India

²Regional Agricultural Research Station, ANGRAU, Nandyal-518502, Andhra Pradesh, India

*Email: ramareddy.yettapu@gmail.com

NDLA 3116-4 is a desi cotton (*Gossypium arboreum* L.) advanced breeding line developed from the cross NDLA 2933 × ARBa-79 - 08 through pedigree breeding method and is suitable for cultivation in *kharif* season under rainfed conditions. The line was identified and developed by Acharya N.G. Ranga Agricultural University, Regional Agricultural Research Station (RARS), Nandyal. The proposed entry has a high ginning percentage (40%).

NDLA 3116-4 is a desi cotton with compact type, early maturing, suitable for *kharif* season in cotton-growing districts in Andhra Pradesh. The genotype is having high ginning percentage (40%) coupled with a high seed cotton yield potential (1973 kg/ha). The plant has light green foliage with medium hairiness on the plant, medium palmate leaves with medium hairiness, cream petals with petal spot and with yellow pollen, medium bracts, oval type boll with average boll weight of 2.8 to 3.0 g.

NDLA 3116-4 exhibited a high ginning percentage of 40 and 39.6% in station trials at the Regional

Agricultural Research Station, Nandyal, during 2015-16 and 2016-17, respectively, and in AICRP on Cotton trials showed the ginning percentage of 39.6% (average of six locations) during 2018-19 and 39.9% (average of two locations) during 2019-20. The results from both the RARS, Nandyal and All India Coordinated trials showed the high ginning percentage ability of the genotype NDLA 3116-4

The genotype NDLA 3116-4 also possesses the desirable fibre quality traits, i.e. upper half mean length of 23.2 mm to 24.8 mm, micronaire of 6.2 µg/inch, tenacity of 28.1 g/tex, uniformity index of 82% and elongation percentage of 6.7%. NDLA 3116-4 is a desi cotton genotype suitable for rainfed conditions of cotton-growing areas in Andhra Pradesh.

In view of the above, the cotton entry, NDLA 3116-4 developed from the Regional Agricultural Research Station, Nandyal, (ANGRAU) is recommended for registration with ICAR-NBPGR, New Delhi for its high ginning percent (40%).

10. CSC-025 (IC648615; INGR23044), a cotton germplasm with salinity stress tolerance

and

11. CSC-057 (IC648616; INGR23045), a cotton germplasm with Salinity stress tolerance

Vineeth TV^{1*}, Indivar Prasad², AR Chinchmalatpure¹, Lokeshkumar BM³, Shrvan Kumar⁴ and PC Sharma³

¹ICAR-CSSRI Regional Research Station, Bharuch-392012, Gujarat, India

²ICAR-Indian Institute of Vegetable Research, Varanasi-221305, Uttar Pradesh, India

³ICAR-Central Soil Salinity Research Institute, Karnal-132001, Haryana, India

⁴ICAR-Central Arid Zone Research Institute, Jodhpur-342003, Rajasthan, India

*Email: vinee2705@gmail.com

Cotton is a significant cash crop globally, and its cultivation is alarmingly being shifted to marginal saline agroecosystems. It is a moderately salt-tolerant crop with a threshold salinity level of 7.7 dS/m (Peng *et al.*, 2016). Asiatic diploid cotton (*Gossypium herbaceum* and *Gossypium arboreum*) is the better option in highly saline, arid and semiarid regions of India, as compared to *Gossypium hirsutum* and *Bt* cotton. At ICAR-Central Soil Salinity Research Institute, Regional Research Station (ICAR-CSSRI, RRS), Bharuch, Gujarat, we developed two salt-tolerant cotton (*Gossypium herbaceum* L.) genotypes, CSC-057 and CSC-025. CSC-057 (GShv 451/08 × GBhv 290) and CSC-025 (GShv 297/07 × GBhv 290) are advanced near homozygous breeding lines, developed by the pedigree breeding method and currently in the F8 generation. Based on 4

years evaluation (2017-20) trials (Table 1) conducted on naturally salt affected field (soil salinity ECe 9 dS/m) and micro plots at ICAR-CSSRI, RRS, Bharuch, Gujarat, CSC-057 (5.5) and CSC-025 (9) showed 16% and 90% higher leaf K/Na ratio, respectively as compared to the check variety, G Cot 23 (4.73). CSC-057 (14.65 q/ha) and CSC-025 (16.41 q/ha) also registered 8.1 and 21.1% higher seed cotton yield, respectively, as compared to the salt-tolerant check variety G Cot 23 (13.5 q/ha). Moreover, CSC-057 (2.87g/boll) and CSC-025 (2.98 g/boll) showed 6.5% and 10.6% higher boll weight, respectively, as compared to the check variety, G Cot 23 (2.7 g/boll).

Reference

Peng J, J Liu and L Zhang (2016) Effects of soil salinity on sucrose metabolism in cotton leaves. *PLoS One* 11: e0156241.

Table 1: Performance of CSC-057, CSC-025 and local check, G Cot 23 under salt stress (ECe 9 dS/m)

Genotype	Year of reporting	Seed cotton yield (kg/ha)	Boll weight (g/20 boll)	Leaf K/Na ratio
CSC-057	2017	1717	61	6.9
	2018	1720	58	5.1
	2019	1302	53	4.8
	2020	1121	58	5.2
	mean	1465	57.5	5.5
CSC-025	2017	1892	62	11.7
	2018	1807	60	9.5
	2019	1755	57	7.7
	2020	1110	60	7.1
	mean	1641	59.75	9
Check variety (G Cot 23)	2017	1756	58	5.7
	2018	1396	53	4.9
	2019	1290	50	4.4
	2020	980	55	3.9
	mean	1355.5	54	4.73

12. IC275447 (INGR23046), a chickpea germplasm with ascochyta blight resistance

Gayacharan^{1*}, Upasana Rani², Sarvjeet Singh², AK Basandrai³, VK Rathee³, AK Srivastava⁴, Kuldeep Tripathi¹, Neeta Singh¹, GP Dixit⁴, Jai C Rana⁵, SPandey¹ and Ashok Kumar¹

¹ICAR-National Bureau of Plant Genetic Resources, Pusa Campus-110012, New Delhi, India

²Punjab Agricultural University, Ludhiana-141001, Punjab, India

³CSK Himachal Pradesh Agriculture University, Palampur-176062, Himachal Pradesh, India

⁴ICAR-Indian Institute of Pulses Research, Kanpur-208024, Uttar Pradesh, India

⁵Alliance of Bioversity International and CIAT, NASC Complex, Pusa Campus-110012, New Delhi, India

*Email: gayacharan@icar.org.in

Chickpea (*Cicer arietinum* L.) is the second largest pulse crop grown worldwide and Ascochyta blight caused by *Ascochyta rabiei* (Pass.) Labr. is the most devastating disease of the crop in all chickpea growing areas worldwide particularly in areas with cool, humid weather during the growing season. The pathogen *A. rabiei* is highly variable. The resistant sources available are not sufficient, and new needs to be identified from time to time as resistance breakdown in existing chickpea varieties is very frequent due to the fast evolution of new pathotypes of the pathogen. Therefore, this work was undertaken to evaluate the existing chickpea germplasm diversity conserved in the Indian National Genebank against the disease under artificial epiphytotic conditions. A total of 1970 chickpea accessions were screened at two hotspot locations, viz. PAU, Ludhiana and HAREC, CSK HPKV, Dhaulakuan following standard procedures. An artificial standard inoculation procedure was followed for uniform spread of the pathogen (Gayacharan *et al.*, 2020). The promising accessions were further validated against the disease over the years under artificial

epiphytotic conditions in two locations, i.e. PAU, Ludhiana and CSK HPKV, Dhaulakuan. The accession IC275447 was found resistant against the pathogen. The accessions were also found to be resistant in earlier MLE trials done during 2005-06, coordinated by NBPGR; however, this was not published or utilized in breeding programs. IC275447 was found constantly resistant during 2015-16 (disease severity score 3), 2017-18 (disease severity score 2), 2018-19 (disease severity score 2) and 2019-20 (disease severity score 3) at PAU, Ludhiana location. The accession was also screened at HAREC, CSK HPKV, and Dhaulakuan during 2015-16 (disease severity score 2) and 2016-17 (disease severity score 3), and was found resistant. In the screening of germplasm susceptible check varieties, viz. L550 and JG62 were used. The disease severity score for checks was 9 (highly susceptible).

Reference

Gayacharan, U Rani, S Singh, AK Basandrai, VK Rathee, K Tripathi, N Singh, GP Dixit, JC Rana, A Kumar and K Singh (2020) Identification of novel resistant sources for ascochyta blight (*Ascochyta rabiei*) in chickpea. *Plos one* 15(10): e0240589.

13. ICPL 88039A & ICPL 88039B (IC648617 & IC648618; INGR23047), a pigeonpea germplasm with cytoplasmic male sterility and early maturity

Satheesh Naik SJ^{1*}, Abhishek Bohra², IP Singh¹, Dibendu Datta³ and RK Mishra¹

¹ICAR-Indian Institute of Pulses Research, Kanpur-208024, Uttar Pradesh, India

²SABC and CCFI, Murdoch University, Western Australia

³ICAR-IIPR Regional Station, Bhopal-462030, Madhya Pradesh, India

*Email: satheeshnaikagri@gmail.com

Hybrids based on cytoplasm male sterile (CMS) technology have now emerged as a potential method for significant improvement of pigeonpea [*Cajanus cajan* (L.) Millisp.] productivity in India (Saxena *et al.*, 2016). The efforts of more than three decades R&D of IIPR on the advancement of the CMS technology have led to the development of two early duration

pigeonpea hybrids namely IPH 15-3 and IPH 09-5 for commercial cultivation in the North West Plain Zone of India which comprises Western UP, New Delhi, Haryana, Punjab and Uttarakhand states (<https://seednet.gov.in>). CMS-based hybrid technology requires three components, viz. sterile (A), maintainer (B) and restorer (R) lines. In pigeonpea, of the eight (A1

to A8) different cytoplasm sources available to harness the heterosis, A2 and A4 cytoplasms derived from *Cajanus scarabaeoides* and *Cajanus cajanifolius*, respectively, have been optimized for regular use in hybrid pigeonpea breeding programmes of the country (Bohra *et al.*, 2019 and 2021). The popular early maturing (<150 days) variety ICPL 88039 has been converted as the donor CMS line in the background of A2 cytoplasm (*Cajanus scarabaeoides*). The new CMS lines ICPL 88039A have been converted using standard backcross technique with A₂ cytoplasmic donors (GT 288A). ICPL 88039A is a stable for male sterility and produces completely non-functional pollen grains over the years of sterility testing conducted using field screening and standard Acetocarmine (2%) staining method. The developed new CMS lines, ICPL 88039A have early maturity along with the maintainer and restoration system reported here, which could create novel opportunities for accelerated hybrid breeding in pigeonpea.

The sterility in ICPL 88039A was confirmed by different methods, viz. flower morphology dissection by microscopic study, standard Acetocarmine (2%) staining testing of pollen grains and by using DNA markers confirming the sterility of the genotype. The

cent per cent sterile plants were selected and simultaneously advanced to develop ICPL 88039A. The line ICPL 88039A is an early maturing (140-150 days) indeterminate semi-spreading plant type having a plant height of 170-185 cm and the number of primary branches ranges from 12-16 branches/plant. ICPL 88039A has attained its 50% flowering in 100-105 days. The plant produces yellow colour flowers with sparse streaks and green pods with brown streaks. The seed coat is brown and 100 seeds weigh about 8.9-10 g.

References

- Bohra A, IP Sing, SJ Satheesh Naik, PG Patil, RK Mishra, F Singh, D Datta, A Das and NP Singh (2019) Hybrid technology for pigeonpea improvement, ICAR-IIPR, Kanpur, pp 64.
- Bohra A, G Prasad, A Rathore, RK Saxena, SJ Naik, S Pareek, R Jha, L Pazhamala, D Datta, G Pandey, A Tiwari, AK Maurya, KR Soren, M Akram, RK Varshney and NP Singh (2021) Global gene expression analysis of pigeonpea with male sterility conditioned by A2 cytoplasm. *Plant Genome*. 14(3): e20132. doi: 10.1002/tpg2.20132. Epub 2021 Sep 8. PMID: 34494714.
- Saxena Kulbhushan, CV Sameer Kumar, R Saxena, R Kumar Singh, Invinder, Hingane Anupama, M Mula, Patil Suyash and Varshney Rajeev (2016) Hybrid pigeonpea; accomplishments and challenges for the next decade. *Legume Perspectives*. 30-32. <https://seednet.gov.in>

14. IC140784-P1 (IC648619; INGR23048), a cluster bean germplasm with early maturing determinate growth type and synchronous maturity

Kartar Singh^{1*}, VS Meena¹, Neelam Shekhawat¹, HR Mahla² and Veena Gupta³

¹ICAR-NBPGR Regional Station, Jodhpur Rajasthan-342005, India

²ICAR-Central Arid Zone Research Institute, Jodhpur-342003, Rajasthan, India

³ICAR-National Bureau of Plant Genetic Resources, Pusa Campus-110012, New Delhi, India

*Email: kartar1532@gmail.com

Cluster bean (*Cyamopsis tetragonoloba* L.) is an important drought-tolerant legume cultivated predominantly under rainfed conditions in the arid and semi-arid regions of India. Since the crop is frequently exposed to moisture stress, early maturity is a highly desirable trait that enables plants to escape terminal drought and reduces the risk of crop failure. Most of the improved cultivars popular in the region have a maturity time of 90-95 days after sowing. Due to this, there is always a very high chance of crop failure due to water scarcity. Therefore, any germplasm line having a short duration of maturity will be a useful source of donors in breeding programme as it may escape the dry spell. The availability of diverse genetic resources is essential for developing improved cultivars adapted to such challenging environments (Duvick *et al.*, 1986).

As part of the germplasm characterization and evaluation programme at ICAR-NBPGR Regional Station, Jodhpur, 50 cluster bean accessions collected from Rajasthan, Gujarat, and Maharashtra were evaluated during *kharif* 2020 using an Augmented Block Design with national checks RGC-936 and RGC-1066. During the evaluation, a superior single-plant selection, IC140784P1, was identified from accession IC140784. The selection exhibited desirable traits including early maturity, single-stem growth habit, determinate flowering, all-node pod bearing, and uniform maturity. IC140784P1 initiated flowering 23 days after sowing, attained 50% flowering in 32 days, and matured within 82 days after sowing. This is approximately 10-15 days earlier than the widely cultivated improved varieties that generally mature in

90–95 days. The stability of the early-maturity trait was confirmed through evaluations conducted during *Kharif* 2021, Summer 2022, and *kharif* 2022 along with national check varieties.

The identified accession represents a valuable genetic resource for guar improvement. Its determinate and single-stemmed growth habit promotes synchronized maturity, reduces the incidence of black seeds, and improves seed quality (Gopala and Dwivedi *et al.*, 2009). For highest seed yield, the guar plant should have early maturity, a determinate growth type, optimum plant height, all node pod bearing, a high branching number and a high number of clusters per plant, with each cluster having the maximum number of pods (Paroda and Saini *et al.*, 1978). Early-maturing and determinate plant types are particularly

advantageous for drought-prone regions and intensive cropping systems, where shorter crop duration and uniform maturity are essential. Therefore, accession IC140784P1 can serve as an important donor parent in breeding programmes aimed at developing short-duration, climate-resilient, and high-yielding cluster bean cultivars for sustainable production in dryland agriculture.

References

- Duvick DN (1986) Plant breeding; past achievements and expectations for the future. *Econ. Bot.* 40: 289-297.
- Gopala KR and SL Dwivedi (2009) Plant ideotype and breeding strategies in cluster bean (*Cyamopsis tetragonoloba* L.). *Indian J. Genet. Plant Breed.* 69(4): 367-375.
- Paroda RS and ML Saini (1978) Genetic architecture of yield and its components in guar. *Indian J. Genet. Plant Breed.* 38: 17-24.

15. CAZG-109 (IC648620; INGR23049), a cluster bean germplasm with high test weight, long fleshy pod, glabrous leaves, stem and pods

HR Mahla* and Ramavtar Sharma

ICAR-Central Arid Zone Research Institute, Jodhpur-342003, Rajasthan, India

*Email: hans.mahla@icar.nic.in

Cluster bean [*Cyamopsis tetragonoloba* (L.) Taub.] a drought hardy legume is an economically important crop of the Indian arid zone. It is globally recognized for the industrially important seed-borne gum besides other multifarious uses in food, feed and fodder. The guar seed is composed of three layers, namely the outer husk (16–18%), the germ (43–46%) and the endosperm (34–40%). The economically important traits like seed weight (test weight $\approx$ 30 g), pod length ($\approx$ 6.0 cm) and seeds per pod ($\approx$ 6) have a very narrow range in seed type varieties (Dadheech *et al.*, 2019). Realizing the paucity of diversity in traits for further improvement of yield and suitable plant type in guar, efforts were made to create variability for seed weight and related traits through mutagenesis.

Seed weight has been recognized as one of the important traits contributing to seed yield in most of the field crops. The test weight of released seed type varieties of guar has been reported to vary around 30–35 g. Though pod length and seed weight either lacked correlation or were negatively correlated to seed yield (Dadheech *et al.*, 2020), a source for large seed and longer pods would be desirable for incorporating in a breeding program with a long-term goal. The two traits show positive correlation with yield per plant among seed type varieties having pod length <6 cm and seed

weight <3.6 g. Thus, there is a possibility of altering linkages in favourable direction. The genotype, CAZG-109, identified for pod length (12.3 cm) and bold seed (test weight 46.8 g) was also evaluated with 5 genotypes varying for the two traits for 4 seasons (Table 1). Pusanavbahar, a vegetable type variety, had maximum test weight (37.4 g) after CAZG-109, while CAZG-102 had minimum (28.8 g). CAZG-109 expressed high values for pod length (12.3 cm), pod width (10.9 mm) and seeds per pod (9.6), hence it could be a desirable donor for improving pod length, seeds per pod and seed weight depending upon linkages with other yield attributing traits, especially pods per plant. Morphologically, the selected genotype CAZG-109 is branched, luxuriant, glabrous having long and thick pods that are suitable for vegetable type as well as seed yield.

References

- Dadheech R, R Sharma, HR Mahla and RK Bhatt (2019) Morphological and molecular diversity analysis in cluster bean [*Cyamopsis tetragonoloba* (L.) Taub]. *Int. J. Pharm. Biol. Sci.* 9(3): 378-387.
- Dadheech R, R Sharma, HR Mahla and RK Bhatt (2020). Plant architecture evolution for higher yield in cluster bean [*Cyamopsis tetragonoloba* (L.)] under arid conditions. *Indian J. Agric. Sci.* 90(5): 919-923.

Table 1: Pod and seed characteristics of promising genetic stocks–CAZG-109 (Average of 4 seasons)

S.N.	Genetic Stock	Pod length (cm)	Pod width (mm)	Seeds per pod	Test weight (g)
1	CAZG - 102	6.3	7.6	8.6	28.8
2	RGC-936	5.9	7.9	7.4	32.1
3	HG-2-20	6.1	7.4	7.6	36.7
4	RGC-1066	6.1	8.3	8.0	33.8
5	Pusanavbahar	9.9	9.5	9.0	37.4
6	CAZG-109	12.3	10.9	9.6	46.8

16. Improved Pusa Uday-1 (IBL BC2F4) (IC646853; INGR23050), a cucumber germplasm with highly stable tropical gynoecious lines for hybrid breeding and donor for F locus introgression into elite genotype

TK Behera¹, AD Munshi², SS Dey², Harshwardhan Choudhary², GS Jat² and BS Tomar²

¹ICAR-Indian Institute of Vegetable Research, Varanasi-221305, Uttar Pradesh, India

²ICAR-Indian Agricultural Research Institute, Pusa campus-110012, New Delhi, India

Email: tusar@rediffmail.com

Cucumber (*Cucumis sativus* L.) originated in India with a huge diversity in terms of different economically important traits. Most of the commercial cultivars of cucumber in other parts of the world are with F1 hybrids in a gynoecious background, but the contribution of hybrid seed is not more than one-third of total seed replacement in India. However, contribution of the indigenously developed F1 hybrids is not very significant because of the non-availability of gynoecious lines in the background of the tropical genotypes. The gynoecious sex form has been extensively exploited in heterosis breeding for enhancing the productivity of the cucumber worldwide. Most of the gynoecious lines of cucumber are adapted to the temperate regions they are not stable at temperatures >35°C. Gynoecious hybrid cultivars produced by crossing a gynoecious line as female parent have become an integral part of the cucumber improvement programme. In addition to economical hybrid seed production, gynoecious sex expression provided a more concentrated fruit set and early maturity compared to the conventional monoecious × monoecious-based hybrids. Therefore, development of tropical gynoecious lines with stability in sex expression even at temperatures above 40°C is urgently required to develop indigenous F1 hybrids in cucumber and to strengthen the Indian seed industry. The different sex types in cucumber are mainly determined by three major genes F/f, M/m and A/a. The loci F promotes femaleness on the whole plant level, while the gene M determines whether flowers are

unisexual.

The objective of this study was to introgress the F locus into the elite cultivar, Pusa Uday, using the American pickling gynoecious line, Gy421 (EC636513) as a donor. A marker-assisted backcross breeding strategy was adopted for rapid recovery of the recurrent parent along with the donor trait, gynoecy, through foreground and background selection. Two SSR markers (SSR-13251 and SSR- 15516) and one gene-based SCAR marker (CsACS1G) were used for foreground selection of F locus (gynoecy) in 192 developed BC₁F₁ progenies. The identified heterozygote plants were subjected to the selection process. Heterozygote plants for F locus were identified using the co-dominant SSRs along with one SCAR marker. Besides, selection for the plants was done on the basis of the phenotype and only the plants with resemblance to the recurrent parent along with F loci were selected. These plants were selfed to produce the BC₁F₂ and were under gone foreground selection and phenotypic evaluation. The plant #29 from BC₁F₂ population was selected and selfed to develop BC₁F₃ population.

859 markers were used from across the genome of cucumber for background profiling of BC₂F₃ plants, 121 markers found to be polymorphic between donor parents (G421) and recipient parents (Pusa Uday). Background analysis revealed that the recurrent parent genome (RPG) recovery ranged from 69.63 to 78.72% in BC₂F₄ generation. From this BC₁F₃ population, 10 gynoecious plants with more lateral branches (4-5) and

better fruit quality (L:D ratio >3.4) were selected based on the foreground, background and phenotypic selection. The application of MAS with the MABC breeding program accelerated the recovery of the RP genome, reducing the number of generations and the time for incorporating gynoecious gene. The line,

Improved Pusa Uday was selected from IBLs with only female flowers and higher lateral branches, *desi* fruit type resembling Pusa Uday at marketable stage (L:D $\geq$ 3). Hence, this line could be directly utilized or could be utilized in breeding programme to develop gynoecious based F1 hybrid cucumber.

17. IMPU-2 IBL11BC1F4 (IC646854; INGR23051), a cucumber germplasm with Tropical gynoecious line for use in hybrid breeding

TK Behera^{1*}, AD Munshi², SS Dey², Harshwardhan Choudhary², GS Jat² and BS Tomar²

¹ICAR-Indian Institute of Vegetable Research, Varanasi-221305, Uttar Pradesh, India

²ICAR-Indian Agricultural Research Institute, Pusa campus-110012, New Delhi, India

*Email: tusar@rediffmail.com

Cucumber (*Cucumis sativus* L.) originated in India with a huge diversity in terms of different economically important traits. Most of the commercial cultivars of cucumber in other parts of the world are with F1 hybrids in a gynoecious background but the contribution of hybrid seed is not more than one-third of total seed replacement in India. However, contribution of the indigenously developed F1 hybrids is not very significant because of non-availability of gynoecious lines in the background of the tropical genotypes. The gynoecious sex form has been extensively exploited in heterosis breeding for enhancing productivity of cucumber worldwide. Most of the gynoecious lines of cucumber are adopted to the temperate regions they are not stable at temperatures >35°C. Gynoecious hybrid cultivars produced by crossing gynoecious line as female parent have become integral part of cucumber improvement programme. In addition to economical hybrid seed production, gynoecious sex expression provided a more concentrated fruit set and early maturity compared to the conventional monoecious $\times$ monoecious based hybrids. Therefore, development of tropical gynoecious lines with stability in sex expression even at temperature above 40°C is urgently required to develop indigenous F1 hybrids in cucumber and strengthen the Indian seed industry. The different sex types in cucumber are mainly determined by three major genes F/f, M/m and A/a. The loci F promotes femaleness on the whole plant level while the gene M determines whether flowers are unisexual.

Introgression of F locus into Pusa Uday was attempted through marker assisted backcross breeding using markers for foreground and background selection using Gy421 (EC636513) as a donor. Two SSR markers (SSR-13251 and SSR-15516) and one gene-based SCAR marker (CsACS1G) were used for

foreground selection of F locus (gynoecy). Heterozygote plants for F locus using these three markers and plants phenotypically matched to recipient parent, Pusa Uday at BC₁F₁ stage. These plants were selfed to produce the BC₁F₂ and were under gone foreground selection and phenotypic evaluation. The phenotypic data of ninety-two plants of the BC₁F₃ population, including yield and its component traits including sex expression, number of lateral branches, fruit length, fruit diameter and L:D ratio were recorded. Sex expression was recorded on individual plants by counting the number of pistillate nodes in the first twenty nodes of the main stem and the total number of lateral branches was also recorded for each plant in the first twenty nodes. Mean fruit L:D ratio per plot was obtained by measuring the length and diameter of 5–10 randomly selected fruits. Out of ninety-two plants in the BC₁F₃ population, phenotypically all the plants showed only female flowers, thirty-six plants showed three lateral branches and fifteen plants showed four lateral branches, whereas fifty-two plants showed an L:D ratio between 3.1 and 4.0. All forty-one inbred backcross lines IBLs (BC₁F₄) were purely gynoecious in sex expression and their fruits were similar to the Pusa Uday type with non-bitter fruits and more than four lateral branches and twenty-eight IBLs had fruit length >15 cm. The maximum fruit length observed was 20.5 cm in IBL12. A total of thirty-five IBLs had average fruit diameter more than 5 cm with the highest average fruit diameter (7.8 cm) recorded in IBL18. Among these lines, only ten IBLs produced fruits with more than 3.0 L:D ratio and the maximum L:D ratio was 4.0 in IBL11. This plant was selected as Improved Pusa Uday-2.

A total of 859 markers were used from across the genome of cucumber for background profiling of BC₁F₃ plants, 121 markers were found to be polymorphic

between donor parents (G421) and recipient parents (Pusa Uday). Background analysis revealed that the recurrent parent genome (RPG) recovery ranged from 60.63% to 78.72% in BC₁F₃ generation. From this BC₁F₃ population, 10 plants were extracted based on the foreground, background and phenotypic selection, which were gynoeious in habit, with more later

branches (4-5) and better fruit quality (L:D ratio >3.4). The line, Improved Pusa Uday-2 was selected from IBLs has complete female flowers and higher lateral branches *desi* fruit type (Pusa Uday) at marketable stage (L:D ≤ 3). Hence, this line could be directly utilized in breeding programme to develop gynoeious based hybrid cucumber.

18. VRSG-7-17 (IC0630886; INGR23052), a sponge gourd germplasm with characteristic basmati aroma

Tribhuvan Chaubey^{*}, Sudhakar Pandey, RK Dubey, P Karmakar, DR Bhardwaj, PM Singh, B Singh, Vidya Sagar, DP Singh, KK Pandey, J Haldar and TK Behera

ICAR-Indian Institute of Vegetable Research, Varanasi 221305, Uttar Pradesh, India

^{*}Email: tchaubey@gmail.com

The flavour and aroma in the food are determined by chemical senses of taste and smell. India is home of several traditional aromatic foods, of which Basmati rice is the most popular. A harmonious aroma blend in the food increases its acceptability and likeability. Apart from Basmati rice, the aroma has been reported as one of the quality characteristics in a variety of crops, fungi, microbes and processed foods. Likewise, consumers' preference for aromatic sponge gourd varieties over non-aromatic varieties has resulted in the high price of aromatic cultivars. *Luffa cylindrica* (L.) Roem (2n=26), as sponge gourd is an important edible vegetable member of the Cucurbitaceae family in the Indian subcontinent. It is suitable for cultivation during the summer and rainy season. Its leaves, flowers and fruits are edible. An aromatic sponge gourd genotype (VRSG-

7-17) has been identified at ICAR-Indian Institute of Vegetable Research (IIVR), Varanasi.

The genotype VRSG 7-17 is a unique line of sponge gourd with characteristic aroma resembling typical aroma of 'Basmati rice' in its fruits, peel and leaves (Table 1). The fruits of this novel sponge gourd genotype (VRSG7-17) also retain special aroma even after cooking and/or boiling at high temperatures.

The inheritance of fruit colour was found to be controlled by a single gene with incomplete dominance. The SPME and GC-MS analysis showed the presence of four compounds hexanal, 1-octen-3-ol, 3-octanone and limonene as dominant volatiles present in this aromatic genotype. This genotype is also tolerant to downy mildew and melon weevil and is less susceptible to leaf miner under field conditions.

Table 1: Morpho-agronomic characters of sponge gourd genotype VRSG7-17

Characters	Genotype VRSG7-17	Characters	Genotype VRSG7-17
Days to first male flower	43 days	Vine length	6.28 meter
Days to first female flower	47 days	Fruit weight	156.5 g
Days to first fruit harvest	55.5 days	Fruit yield/ plant	1.13 Kg
Fruit length	27.46 cm	Number of fruit/plant	3.6
Fruit diameter	3.35 cm	Fruit colour	Light green

19. DRMRIJ 12-26 (IC648622; INGR23053), an Indian mustard germplasm with a novel white rust resistant gene

KH Singh^{*}, AK Thakur, Lal Singh, Guman Singh, HS Meena, N Parmar, Priyamedha, Bhagirathram and VV Singh

ICAR-Directorate of Rapeseed-Mustard Research, Bharatpur-321303, Rajasthan, India

^{*}Email: kharendrasingh@gmail.com

White rust caused by *Albugo candida* (Pers. ex Lev.) Kuntze is one of the most devastating diseases of

Crucifers and causes severe losses in Indian mustard (*Brassica juncea*). Symptoms of the disease may appear

at vegetative stage itself and sustain even at flowering stage. Maximum damage is caused by stag head formation due to malformation of flowering buds directly causing the yield losses. Indian germplasm of (*Brassica juncea* L.) has generally been reported to have susceptibility against existing pathogens. Breeding efforts, however, have brought few successes in form of developing resistant cultivars like JM1, Basanti and JM3. One germplasm line BIO YSR (IC443623), has earlier been registered as white rust-resistant genetic stock derived from an exotic source. Genetic studies have revealed dominance of resistance over susceptibility. Resistance to white rust is governed by a single dominant gene.

DRMRIJ 12-26 is an inbred line derived from a cross between KLM227 and EC597313 (exotic germplasm), followed by individual plant selection. Cross was attempted during 2008-09 between KLM227 and EC597313. DRMRIJ 12-26 was induced into AICRP-RM during 2015-16 and had been evaluated continuously for four years during 2015-16, 2016-17, 2017-18 and 2018-19. White rust reaction on leaves was recorded at two different stages, viz. 75 and 100 days after sowing, while staghead formation was recorded at the flowering stage. Candidate strain DRMRIJ 12-26 expressed resistant reaction against white rust at 75 days stage, which was at par with the resistant check BIO YSR during two years of testing at Hisar and one year of testing at Morena centre. Mean Disease index of

DRMRIJ 12-26 was 0.0% in comparison to 32.5% of susceptible check Rohini. Disease reaction against white rust on leaves at 100 days stage was recorded at eight locations, viz. Hisar, Kangra, Kanpur, Pantnagar, Morena, Ludhiana, Sri Ganganagar and New Delhi, which included three hot spots (Kangra, Pantnagar and Morena) (Table 1). On the basis of the mean disease index over all five locations for four years, candidate strain DRMRIJ 12-26 expressed 3.3%, which was better than the resistant check BIO YSR (7.1%) and susceptible check Rohini (28.54%). Observations on staghead formation were recorded at Morena, Sri Ganganagar and Ludhiana, based on mean disease index over three years. DRMRIJ 12-26 had 1.4% disease incidence in comparison to resistant check, BIO YSR (9.9%) and susceptible check Rohini (6.9%). The already reported IP markers, viz. AcB1-A4.1 from Heera and AcB1-A5.1 from Donskaja and WRR candidate gene-specific marker, WR-D from Donskaja have been validated in the concerned strain, DRMRIJ 12-26. It has been found that this genotype does not possess any of the already mapped loci/genes for white rust resistance. This infers that the resistance to WR expressed by DRMRIJ 12-26 is governed by another third/novel gene, which has not been discovered and mapped yet. So, this genotype may be used in developing a mapping population for tagging of that WRR gene in future research endeavours.

Table 1: Mean disease index of candidate strain DRMRIJ 12-26 and check genotypes against white rust pathogen

S. N.	Location	White rust (% WR severity)								
		Disease reaction at 75 days			Disease reaction at 100 days			Staghead formation		
		DRM RIJ 12-26	Bio YSR (Res. C3check)	Rohini (Sus. Check)	DRMRI J 12-26	Bio YSR (Res. Check)	Rohini (Sus. Check)	DRMRI J 12-26	Bio YSR (Res. Check)	Rohini (Sus. Check)
1	Kangra	-	-	-	3.9	14.23	44.83	-	-	-
2	Kanpur	-	-	-	0.0	0.0	31.85	-	-	-
3	Pantnagar	-	-	-	2.8	9.4	27.22	—		
4	Hisar	0.0	0.0	34.15	0.0	0.0	34.75	—		
5	Morena	0.0	0.0	30.9	10.0	8.8	31.18	0.0	9.9	15.2
6	Ludhiana	-	-	-	0.0	-	20.7	0.0	-	0.0
7	Sri Ganganagar	-	-	-	9.6	-	8.9	4.1	-	5.4
8	New Delhi	-	-	-	0.0	-	28.9	—		
	Mean	0.0	0.0	32.53	3.3	7.1	28.54	1.4	9.9	6.9

20. IC384578 (INGR23054), a linseed germplasm with high number of capsules

DP Wankhede^{1*}, Vikender Kaur¹, SS Gomashe², A Saroha¹, Deepa Pal¹, Arti Bartwal¹, J Aravind¹, J Radhamani¹, S Rajkumar¹, Rajesh Kumar¹, Abhishek Sengupta³, Sandeep Kumar¹, Ashok Kumar¹ and Dinesh Chand²

¹ICAR-National Bureau of Plant Genetic Resources, Pusa Campus-110012, New Delhi, India

²ICAR-NBPGR Regional Station, Akola-444104, Maharashtra, India

³Amity University, Noida-201303, Uttar Pradesh, India

*Email: wdhammaprakash@gmail.com

Linseed (*Linum usitatissimum* L.) is one of the earliest domesticated crops of the world known for its seed oil and fibre. It is self-pollinated annual crop and has two morphotypes, flax type and linseed type which differ substantially in terms of growth, development and agronomic performance. For linseed type, short height, more branching, high number of capsules and seed weight are desirable traits. Lately, linseed is gaining popularity as nutraceuticals and functional food owing to high omega-3 alpha linolenic acid and other bioactive compounds such as lignans and soluble fibres. Productivity of linseed in India is far below (0.543 tonne/ha) than the world's average (1.053 tonne/ha). Despite the Indian regions (Northwest India) being part of the center of origin (central Asiatic centre) of linseed, there have been limited genetic resources identified for traits of economic and agronomic importance. The number of capsules and primary branches per plant has a positive correlation with yield and, therefore are crucial traits for crop improvement. In this perspective, one germplasm accession, IC0384578 has been identified for high capsule numbers per plant based on

evaluation data of four environments in three consecutive years (*rabi* 2017-18, 2018-19 and 2019-20) at New Delhi (zone 1) (28°38'53.7"N 77°09'05.4"E) and at Akola (zone 3) during *rabi* 2019-20 (20°42'03.2"N 77°01'53.6"E) (Saroha *et al.*, 2022). The average number of capsules recorded in IC0384578 were 280.26 which was significantly higher than in check varieties *Kartika* (242.28) and *Shekhar* (230.3) across multilocation-year trials. In addition, IC384578 also possesses other agronomically important traits such as short plant height (58-67 cm) and high number of primary branches (5-10). The average test seed weight is 6.3 g and yield per plant is 14.5 g.

Reference

Saroha A, D Pal, V Kaur, S Kumar, A Bartwal, J Aravind, J Radhamani, S Rajkumar, R Kumar, S S Gomashe, A Sengupta and DP Wankhede (2022) Agro-morphological variability and genetic diversity in linseed (*Linum usitatissimum* L.) germplasm accessions with emphasis on flowering and maturity time. *Genet. Resour. Crop. Evol.* <https://doi.org/10.1007/s10722-021-01231-3>

21. Jawahar Soybean 20-34 (IC596520; INGR23055), a soybean germplasm with early flowering and early maturity

S Gupta^{1*}, G Kumawat¹, R Tripathi¹, N Agrawal¹, GK Satpute¹, S Maranna¹, V Nataraj¹, V Rajesh¹, S Chandra¹, MB Ratnaparkhe¹ and AN Shrivastava²

¹ICAR-Indian Institute of Soybean Research, Indore-452001, Madhya Pradesh, India

²Jawaharlal Nehru Krishi Vishwavidyalaya, Jabalpur-482004, Madhya Pradesh, India

*Email: sanitaishu@gmail.com

Soybean [*Glycine max* (L.) Merr.] is predominantly grown under rainfed conditions in India, where early maturing cultivars are preferred to escape terminal drought and fit into multiple cropping systems. Flowering and maturity in soybean are strongly influenced by photoperiod sensitivity, which is regulated by major maturity genes (*E1*, *E2*, *E3*, and *E4*) (Bernard 1971; Buzzell 1971; Buzzell and Voldeng 1980). Recessive alleles at these loci reduce photoperiod

sensitivity and promote early flowering and maturity, making them valuable targets for soybean improvement in tropical and subtropical environments.

JS 20-34 is an early-maturing soybean variety. In an evaluation of 101 Indian soybean varieties under ambient and extended day length (20 h) conditions, JS 20-34 exhibited one of the lowest levels of photoperiod sensitivity, showing only 29.5% increase in days to flowering under extended day length. This level of

sensitivity was substantially lower than the population mean (51%), indicating its reduced response to changes in photoperiod.

Genotyping using functional marker for major maturity genes revealed that JS 20-34 carries the recessive *e2* allele. Among the 101 varieties examined, only three varieties JS 20-34, PRS 1, and LSb 1, possessed this allele, demonstrating its rarity in Indian soybean. Varieties carrying the recessive *e2* allele exhibited significantly lower photoperiod sensitivity than those possessing the dominant *E2* allele. The presence of the recessive *e2* allele in JS 20-34 is considered a major factor contributing to its early maturity (approximately 87–89 days), making it one of the earliest maturing soybean varieties cultivated in Central India.

The rarity of the recessive *e2* allele in Indian soybean cultivars, combined with the proven adaptation and agronomic performance of JS 20-34,

makes this variety an important genetic resource for soybean breeding. Registration of JS 20-34 as a genetic stock carrying the recessive *e2* allele will facilitate its utilization in marker-assisted breeding programmes aimed at developing early maturing and less photoperiod-sensitive soybean cultivars for diverse agro-ecological regions.

References

- Bernard RL (1971) Two major genes for time of flowering and in soybeans. *Crop Sci.* 11: 242-244. <https://doi.org/10.2135/cropsci0011183X001100020022x>
- Buzzell RI (1971) Inheritance of a soybean flowering response to fluorescent day length conditions. *Can. J. Genet. Cytol.* 13: 703-707. <https://doi.org/10.1139/g71-100>
- Buzzell RI and HD Voldeng (1980) Inheritance of insensitivity to long day length. *Soybean Genet Newsl* 7: 26-29. <https://doi.org/10.1093/jhered/esp113>

22. NC-11 (IC205471; INGR23056), a sesame germplasm with tolerance to deficit soil moisture stress

RK Pasala*, BB Pandey, Ramya K and AL Ratnakumar

ICAR-Indian Institute of Oilseeds Research, Hyderabad-500030, Telangana, India

*Email: pratnakumar@gmail.com

The sesame (*Sesamum indicum* L.) genotype IC205471 is a local landrace from Himachal Pradesh State, India. The genotype was collected from ICAR-NBPGR, New Delhi, India in the year 2015-16. The genotype IC205471 was evaluated from 2018 to 2019 along with two national checks, i.e., GT-10 and TKG- 22, at ICAR-IIOR Research Farm, Narkhoda, Hyderabad, India (17°15'16" N, 78°18'30" E; 542 meters above sea level) under field conditions. The genotype was tested in two different water regimes, i.e. well-watered (WW) and deficit soil moisture stressed (WS), during four different seasons of 2018 and 2019. The deficit soil moisture stress conditions were imposed from flowering (40-45 days after sowing) to physiological maturity (90-95 days after sowing). The moisture levels in WS plots decreased to 35-40% of available soil moisture, equivalent to -4.5 to -5.0 bars of soil water potential, compared to WW conditions (80% available soil moisture). The soil moisture content was monitored on an hourly basis using field-installed real-time soil sensors (Proximal; SoilSenS, V1.0). The outcome of four-season trials revealed that the genotype IC205471 was a superior performer (20.0 to 38.0% higher seed weight/plant) over two national checks, i.e., GT-10 and TKG-22, particularly under WS

conditions. Further, the genotype was found superior in various morphological, physiological, yield and contributing traits, suggesting tolerant nature of genotype to deficit soil moisture stress conditions (Pandey *et al.*, 2021; Sravanthi *et al.*, 2021). The genotype is moderate-maturing (97-99 days) and medium-height (65-75 cm) in nature. The genotype has an indeterminate growth habit with little hair on the leaves, stem, and capsule. It has basal branching and a single flower at each axil. The flowers are white with violet/purple shading and have alternate leaf and capsule arrangements. The seed coat colour is off-white to light brown. In addition, the genotype IC205471 was also evaluated for root structural and functional traits under well-watered and water stress conditions. It was reported that genotype IC205471 exhibited higher transpiration efficiency, harvest index, and seed yield under water stress conditions. Also, the genotype had higher root length, total root surface area, root volume, root dry weight, and root length density compared to the check GT-10 under water stress conditions (Pasala *et al.*, 2021). Therefore, the genotype IC205471 may be used in sesame crop improvement programmes for the development of varieties adapted to deficit soil moisture stress conditions.

References

- Pandey BB, P Ratnakumar, KB Usha, MY Dudhe, GS Lakshmi, K Ramesh and A Guhey (2021) Identifying traits associated with terminal drought tolerance in sesame (*Sesamum indicum* L.) genotypes. *Front. Plant Sci.* 12: 739896.
- Pasala R, BB Pandey, GS Lakshmi, K Ramesh, A Guhey and AV Reddy (2021) An insight into the mechanisms of intermittent drought adaptation in sesame (*Sesamum*

indicum L.): linking transpiration efficiency and root architecture to seed yield. *Acta Physiol. Plant.* 43(11): 148.

- Sravanthi AL, P Ratnakumar, SN Reddy, KB Eswari, BB Pandey, LN Ch Manikanta, KT Ramya, E Sonia, S Mohapatra, K Gopika, PL Anusha and P Yadav (2021) Morpho-physiological quality traits and their association with seed yield in sesame (*Sesamum indicum* L.) indigenous collection under deficit moisture stress. *Plant Physiol. Rep.* 27(1): 132-142.

23. IC626510; GAN/PCT/1212 (INGR23057), an avocado germplasm high Pulp recovery and tolerance to anthracnose

PCTripathi*, Anuradha Sane, Kanupriya, Pritee Singh, G Karunakaran and A Shamina

ICAR-Indian Institute of Horticultural Research, Bengaluru-560089, Karnataka, India

*Email: prakaashtripathi2000@yahoo.co.in

Avocado (*Persea americana* Mill.) is a native of tropical America. It originated in Mexico and Central America, possibly from more than one wild species. Avocado is the most nutritious among fruits. The pulp is rich in proteins (up to 4%) and fat (up to 30%), but low in carbohydrates. Avocados have been known to maintain good cholesterol while reducing bad cholesterol. Avocado is mainly used fresh, in sandwich filling or in salads. It can also be used in ice creams and milk shakes and the pulp may be preserved by freezing. The Avocado is originated in Central and South American countries. A lion's share of the Avocado production of world, produced in these countries. The world's avocado production was 8.06 million tonnes in 2020. Mexico is the largest producer of Avocado in the world followed by Colombia, Dominican Republic, Peru, Indonesia, Kenya, Brazil, Ethiopia, etc. Mexico supplies 45% of the international avocado market. The avocado market is still growing. Since 2012, the global avocado market has seen a rise, and it is expected to reach a valuation of about 23 million dollars by the end of the year 2027. The production in India is very limited and it is not a commercial avocado plantation. The agro-climatic conditions prevailing in various parts of the country appear to be favourable for bringing more areas under avocado. Avocados are grown scattered in southern tropical states like Tamil Nadu, Kerala, Karnataka, and Maharashtra. Also popular in the

northeastern Himalayan states at elevations if 800-1,600 meters. There are three main production regions of avocado in India, the Kodaikanal area, Ooty area of Tamil Nadu, Gundlupet and Kodagu region of Karnataka and a minor area in North Eastern India. These areas may be producing around 5000 tonnes of avocados every year and the production is increasing rapidly as more and more areas are being brought under cultivation. There are only 2 released varieties of avocado in India and there are needs to identify more promising genotypes to meet the rapidly increasing demand for planting material.

A promising seedling selection (IC626510) was developed by individual plant selection from seeds collected from avocado grown in Madikeri (12.4208° 75.739, 1182 m above MSL) by random selection in 2012. The seedlings were raised and planted at IIHR FGB in 2013. It is a promising genotype with large-sized green fruits, with fruits weighing 243 g. Mean fruit length was 9.15 cm and mean fruit diameter was 8.23 cm with pulp recovery >70%. Mean seed weight was 68g with TSS 4.8 °Brix (Table 1). Fruit colour is light green and flesh colour is creamy white. It is late bearing genotype with pear-shaped fruits and fruit is thin-skinned. Fruit shape is obovate, dark green colour with creamish white flesh colour. It is suitable for fresh fruit purposes.

Table 1: Important features in Avocado accession IC626510

Genotype	Leaf length (cm)	Leaf width (cm)	Fruits plant-1 (no)	Fruit weight (g)	Yield (kg)	Fruit length (cm)	Fruit diameter (cm)	Seed weight (g)	TSS (°Brix)	Pulp recovery	Fruit color	Flesh color
Avocado- 12	10.9	6.32	21	243	5.1	9.15	8.23	68	4.8	72.02	Green	Creamy White

24. CHES R-27 (IC642755; INGR23058), a rambutan germplasm with red colour fruit, free Stone bigger fruit size

PC Tripathi^{1*}, G Karunakaran¹, T Sakthivel¹, H Ravishankar² and R Chithraichelan¹

¹ICAR-Indian Institute of Horticultural Research, Bengaluru-560089, Karnataka, India

²ICAR-IIHR-Central Horticultural Experiment Station, Chettalli-571 248, Karnataka, India

*Email: prakaashtripathi2000@yahoo.co.in

Rambutan (*Nephelium lappaceum* var. *lappaceum* L.) is native to Malaysia and Indonesia and belong family of Sapindaceae. The cultivation of Rambutan is spread over all tropical countries. Indonesia and Malaysia contribute more than 95% of total Rambutan production in the world. In India Rambutan was introduced as a homestead garden in Kerala and slowly it became popular and the area is rapidly increasing in parts of Kerala, Karnataka and Tamil Nadu (Tripathi *et al.*, 2014; 2020). Fruits are used mostly fresh. Rambutan is adapted to a warm tropical climate. It is grown commercially within 12-15° of the equator and grows successfully in a wide range of soils. Several new and underutilized fruits were introduced in India during last few decades. Rambutan is one of the emerging underutilized fruits with high global demand. It is introduced in home garden in some parts of Kerala and Tamil Nadu, and a considerable variability of rambutan is found in these areas. A large collection of rambutan variabilities was made at Central Horticultural Experiment Station (IIHR), Chettalli, Karnataka during 2001-12. Some of the collections (37) of rambutan collected and planted in 2001 were evaluated for growth, flowering and fruiting characters during 2007-12. There was a lot of variability in these collections with respect to fruit, fruit size, fruit colour and fruit quality parameters. The growth data revealed that plant height ranged from 2.90 m in CHESR-24 to 6.30 m in CHESR-28. Plant spread (E-W) was highest in CHESR-28 (5.7 m) and lowest in CHES R-24 (1.25 m). Plant spread (N-S) was highest in CHESR-29 (6.2 m) and lowest in CHES R-24 (1.05 m). The yield data revealed that higher numbers of fruits were harvested from CHESR-26 (2025), whereas CHESR-33 registered the lowest number of fruits (100). The highest yield was recorded in CHESR-26 (48.8 kg) while CHESR-33 registered the minimum yield of 2.77 kg. The highest fruit weight was recorded in CHESR- 27

(44.2 g), whereas CHESR-156 recorded the lowest fruit weight of 16.5 g. Fruit length and breadth were found to be highest in CHESR-27. The lowest fruit length and breadth were recorded in CHESR-156. The highest pulp weight was recorded in CHESR-29 (21.36 g). The highest pulp content was recorded in CHESR-29 (52.47%). The lowest pulp content was recorded in CHESR-8 (37.69%). The highest seed weight was recorded in CHESR-27 (2.69 g), while CHESR-7 recorded the lowest seed weight of 1.66 g. Seed content ranged from 5.92% in CHESR-29 to 11.55% in CHESR-16. TSS content varied from 13.1°B in CHESR-222 to 20.2°B in CHESR-35 and CHESR-138. Titrable acidity was found to be lowest (0.26%) in CHESR-29 and highest (1.00%) in CHESR-222. Ascorbic acid content was estimated to be the highest (63.38 mg/100 g) in CHESR-58 and the lowest was recorded in CHES-28 (29.0 mg/100 g). Among the red colour collections, CHESR-27 was found to be superior to other collections. This line CHESR-27 is an early maturing, promising selection. The trees are medium in size and semi-spreading in nature. The fruit is dark red in colour. Average fruit weight is 40 to 45 g. Fruits are free stone an aril can usually be removed from the seed without attachment to the testa. The arilis are white in colour, thick, firm and dry and sweet (TSS- 17.45°B) with pulp recovery of about 42.18 per cent (Tripathi and Karunakaran *et al.*, 2015).

References

- Tripathi PC, G Karunakaran, T Sakthivel, V Sankar and R Senthil Kumar (2014) Rambutan cultivation in India. Technical Bulletin 1, ICAR-IIHR, CHES, Chettalli, Kodagu, Karnataka, pp. 18.
- Tripathi PC and G Karunakaran (2015) New varieties of rambutan. *Int. J. Innov. Hortic.* 4(2): 127.
- Tripathi PC, G Karunakaran, T Sakthivel, V Sankar and R Senthil Kumar (2020) Status and prospects of rambutan cultivation in India. *Acta Hortic.* 1293: 33-40.

25. Lakshamana (IC647019; INGR23059), a tamarind germplasm with big Pod size and high pulp recovery

C Kanupriya¹, G Karunakaran¹, R Venugopalan¹, Pritee Singh¹ and Lakshamanappa²

¹ICAR-Indian Institute of Horticultural Research, Bengaluru-560089, Karnataka, India

²Nandanhalli village, Gubbi Taluk, Tumakuru-572168, Karnataka, India

*Email: kp.kanu@gmail.com

Tamarind (*Tamarindus indica* L.) is a well-known commodity of Indian cuisine and has medicinal and industrial uses. It is a nutritious tree crop of widespread occurrence growing on marginal lands in semi-arid and sub-humid tropical climates of India, making it highly valuable in ensuring food security for the rural poor. Given the great potential of this neglected and underutilized species to address global challenges such as hunger, poverty and climate change adaptation, there is a need to revisit research and development priorities in its favour and to develop strategies together with stakeholders to increase its use.

The tamarind germplasm Lakshmana was collected from the field of Shri Laxmannappa in Nandihalli village of Tumkur district, Karnataka. The passport data for this was: latitude 13.52° N, Longitude-76.74° E and 860 m MSL. It was found to be superior with better yield and pod characters compared to local check and registered mean annual yield (4 years from 2016-2020; Table 1) of 251.4 kg/tree as against 165.0 kg/tree in local trees.

'Lakshamana' is a 40-year-old tree and a regular bearer. It commences flowering in September-October, matures in February-March and harvesting can be done

Table 1: Morpho-agronomic traits of 'Lakshamana' in comparison to the local check

S. No.	Trait	Lakshamana	Local
1.	Fruiting season	Feb-March	Feb-March
2.	Pod length (cm)	25.4	13.5
3.	Pod breadth (cm)	3.8	2.36
4.	Number of pods/kg	24	49
5.	Number of seeds/pod	8.2	14.5

in March-April under Tumkur conditions. Healthy grafts are planted at a distance of 5 m × 5m during June-July under Bengaluru conditions. Water requirement is met through drip irrigation (1-2 litres once in 2 weeks in late afternoon), which is sufficient till trees are 5-6-year-old. Weeding is essential in early stages (first 4 years). Intercrops/ cover crops can be grown to control weeds and conserve moisture. FYM is applied at time of planting and yearly as top dressing. When tree starts bearing, 500g of complete fertilizer (NPK) can be applied twice a year at end of rains. Mature, bearing trees can be given 2-3 kg of complete fertilizer. Tamarind trees are largely free of pests and diseases.

26. DLB-10 (IC646865; INGR23060), a lemon basil germplasm with high leaf size

PL Saran^{*} and MK Suthar

ICAR-Directorate of Medicinal and Aromatic Plants Research, Boriavi, Anand-387310, Gujarat, India

*Email: plsdehradun@gmail.com

Lemon basil (*Ocimum × africanum* Lour) also known as lebutulsi. *Ocimum × africanum* is a hybrid between *O. basilicum* L. and *O. americanum* L. The herb is believed to be native to Asia (South-Eastern Asia), Africa, and South America. It is only basil that is used much more in several cuisines due to its rich, varied flavour, ethnobotanical and pharmacological properties. Lemon-scented basil is known for its sweet citrus flavour and fragrance. The highest antimicrobial activities were demonstrated by lemon basil as compared to other *Ocimum* species. The leaves are a good source of vitamin A, C, K, magnesium, iron, copper and calcium. In addition to vitamins and minerals,

lemon basil contains limonene and citral responsible for its flavour and medicinal properties. Aromatic grasses like lemongrass and citronella are perennial in nature and crops are grown commercially up to 5-6 years of plantation; therefore, farmers are hesitant to grow such perennial grass in their fields to avoid long-duration field blockage. Seasonal variations are observed in a very specific manner in a country like India. The lemon basil is a short-duration crop and harvested even in the same season (70-75 days after planting) compared to perennial aromatic grasses. Even farmers are very choosy about such fast-growing and high-yielding crops for harvesting essential oil and

aromatic compounds like citral. Therefore, lemon basil elite 'DLB-10' was found suitable for maximum leaf area (7.45 cm²) and fresh herbage yield (298 q/ha) at full

flowering stage as compared to farmer check (6.66 cm² and 280 q/ha, respectively). Such elite can be promoted for commercial cultivation.

27. DIB-1 (IC646866; INGR23061), a basil germplasm with rich in methyl eugenol

PL Saran¹*, DK Ghosh², MK Suthar¹ and R Sarkar¹

¹ICAR-Directorate of Medicinal and Aromatic Plants Research, Boriavi, Anand-387310, Gujarat, India

²Bidhan Chandra Krishi Viswavidyalaya Mohanpur, Nadia-741252, West Bengal, India

*Email: plsdehradun@gmail.com

Genus *Ocimum* derived from "ozo" a Greek word meaning to smell. Basil (*Ocimum basilicum* L.) is the popular name given aromatic herb derived from the Greek word 'Basilica' or 'Dragon' means royal plant. *O. basilicum* originated in Asia, Africa and also grows on some Pacific islands. The basil diversity based on appearance, flavours, fragrances, industrial traits, essential oils and natural pigments provides great opportunities for developing new culinary and industrial crops. This species (Italian basil) having very short history of cultivation and has been cultivated as a commercial crop in a few parts of the globe. With the transformed interest in the world over in herbal drugs and aromatics, species propose an immense scope in developing suitable cultivars for food flavouring purposes. Essential oil extracted from green herbage is a source of biologically active compounds. The aromatic compounds found in the essential oils of basil are mainly the mono-terpenes, sesquiterpenes and phenols with their alcohols, esters, aldehydes, ketones and others. As secondary metabolic products, the essential oil occurs in secondary tissues, glands, trichomes of the flower, leaves and tender stems, and is

usually responsible for characteristic pleasant odours and flavours, which account in large part for their value and determine their uses. Moreover, under covid 19 situation, a complementary trend towards natural products was developed due to negative reports on potential carcinogenic hazards of synthetic aroma chemicals. 'Italian' basil, having large-leafed, robust and lettuce type's leaf are very prominent in the food flavouring industry. Basil, generally used for culinary purposes in Italian and Mediterranean cuisines, actually originated in India. It is used both fresh and dried with vegetables, meat, fish, sauces, stews, dressings, herbal teas, liqueurs, salad, poultry, vinegar and confectionery products, mixed drinks, etc. For this purpose, DIB-1 has a short life cycle (70-75 days) as compared to other basil, upward leaves folding with severe puckering and rich in Methyl Eugenol (30%) share in essential oil. DNA fingerprinting also showed that accession DLB-1, having dissimilarity as compared to DIB-2 and DOB-2 was using ISSR and scot markers. Such an elite could be used to add a unique flavour to pizza, pasta, pesto, sauce, soup, salads, etc.

28. IIHR 17 23 SP 08 (IC642158; INGR23062), a tuberose germplasm with single type flowers and green tinge flower buds; resistance to root knot nematode (*Meloidogyne incognita*) and tolerant to leaf burn disease (*Alternaria polianthi*)

T. Usha Bharathi¹, Rajiv Kumar, Sujatha A. Nair, R Umamaheswari, Priti Sonavane, VK Rao and D Kalaivanan

ICAR-Indian Institute of Horticultural Research, Bengaluru-560089, Karnataka, India

*Email: t.ushabharathi@gmail.com

Tuberose (*Polianthes tuberosa* L.) is a commercially important flower crop cultivated for its extended shelf life and pleasant fragrance. Tuberose is a day-neutral plant that grows well in the tropical climate with the temperature range of 20-35°C. It performs well in loamy to sandy loam soil with a pH of 6.5 to 7.5. It is

propagated vegetatively through bulbs, medium size bulbs of 2.0 to 2.5 cm in diameter are planted with a spacing of 30 cm × 30 cm at a depth of 5.0 cm on the raised beds.

It is highly preferred in the local as well as international markets for garland preparation and

perfumery extraction, respectively. IIHR 17 23SP 08 is developed through the open-pollinated seedling selection from GK-TC-4 at ICAR-Indian Institute of Horticultural Research (ICAR-IIHR), Bengaluru, Karnataka, India. The genetic stock IIHR 17 23SP 08 produces single-type white flowers (RHS colour: NNI55D, white group, Fan 4), flower buds with a green tinge, a greater number of spikes (8.47) and bulbs (8.94) per clump per year. It has resistance to root knot nematode and is tolerant to leaf burn disease under field conditions (Table 1).

References

- AL Taylor and JN Sasser (1978) Biology, identification and control of root knot nematode *Meloidogyne spp.* North Carolina State University Graphics, Raleigh, NC, 111 p.
- Narayanappa M and JK Chandra (1984) Fungicidal control of leaf spot of marigold caused by *Alternaria tagetica*. *Indian J. Agri. Sci.* 54(5):691-692.

Table 1: Evaluation of tuberose genotype IIHR-23 SP 08 with checks for root knot nematode and leaf burn disease incidence under field conditions

Genotype	Screening for leaf burn disease*	Root knot nematode Screening**
IIHR-23 SP 08	9.79 (18.24)	1.31
Mexican Single	19.20 (26.00)	2.42
ArkaPrajwal	21.33 (27.52)	2.14
GK-TC-4	15.23 (22.98)	1.68
PhuleRajani	23.59 (29.06)	1.53
BidhanUjwal	21.10 (27.36)	1.38
Calcutta Single	13.00 (21.14)	1.51
Sem±	-	0.11
CD (P=0.05)	-	0.34
CV%	-	10.95

*Disease severity scale (Narayanappa and Chandra *et al.*, 1984);

**Gall index (Taylor and Sasser *et al.*, 1978); Figures in parenthesis are *arcsine* transformed values

29. SM/92-338 (IC648625; INGR23063), a potato germplasm with highly resistant to bacterial wilt (*Ralstonia solanacearum*)

Vinay Bhardwaj¹, Vinay Sagar¹, Salej Sood^{1*}, Sanjeev Sharma¹, Vikas Mangal¹, Sundaresha S¹, Vinod Kumar¹, TA Joseph², SK Kaushik³, BP Singh¹, Mehi Lal⁴ and Devendra Kumar¹

¹ICAR-Central Potato Research Institute, Shimla, Himachal Pradesh, India

²ICAR-CPRI Regional Station, Ooty, Muthurai-643004, Tamil Nadu, India

³ICAR-National Bureau of Plant Genetic Resources, Pusa Campus-110012, New Delhi, India

⁴ICAR-CPRI Regional Station, Modipuram campus, Meerut-250 110 Uttar Pradesh, India

*Email: salej1plp@gmail.com

Potato (*Solanum tuberosum* L.) is the third most important food crop globally and a major staple in India (Sood *et al.*, 2022). Being a clonal crop, it attracts many pathogens that affect potato production. Although late blight, viruses, and potato cyst nematode (PCN) affect potato production globally to a major extent (Bhardwaj *et al.*, 2019), bacterial wilt or brown rot (*Ralstonia solanacearum*) has become a major problem in tropical, subtropical and warm temperate zones of the world (Sagar *et al.*, 2014). The pathogen infects more than 400 plant species and is considered a quarantine pest in seed potato production. Despite extensive screening of thousands of potato accessions and related germplasm, highly effective sources of resistance have remained scarce, restricting the development of resistant cultivars.

SM/92-338 is a promising potato germplasm possessing a high level of resistance to bacterial wilt. The clone was developed through a biparental cross between HB/82-372, a breeding line with resistance to late blight and good combining ability, and Kufri

Pukhraj, a widely adapted and popular Indian potato variety. The clone was selected in 1993 from a segregating population based on superior agronomic performance and disease resistance. Subsequent evaluations at Kufri and Modipuram demonstrated stable adaptation and superior yield performance.

The clone was evaluated for bacterial wilt resistance using a rapid in vitro screening protocol and subsequent pot and field assays. During in vitro screening, SM/92-338 remained healthy up to six days after inoculation, whereas the susceptible cultivar Kufri Jyoti collapsed within four days. Similar responses were observed under pot conditions, where SM/92-338 maintained resistance up to 28 days after inoculation, while the susceptible check wilted completely by 13 days. Replicated field testing in bacterial wilt-sick plots further confirmed its high level of resistance under natural disease pressure.

In addition to bacterial wilt resistance, SM/92-338 has shown agronomic superiority and attractive ovoid tubers. The combination of disease resistance, yield

superiority and desirable tuber characteristics makes this germplasm an important genetic resource for potato improvement programmes targeting bacterial wilt-prone regions of India.

References

Bhardwaj V, S Sood, A Kumar, G Vanishree, S Sharma, S Sundaresha, B Raigond, R Kumar, A Bairwa, M Lal and SK Chakrabarti (2019) Efficiency and reliability of marker-

assisted selection for resistance to major biotic stresses in potato. *Potato J.* 46(1): 56-66.

Sagar V, A Jeevalatha and S Mian (2014) Potato bacterial wilt in India caused by strains of phylotype I, II and IV of *Ralstonia solanacearum*. *Eur. J. Plant Pathol.* 138: 51-61.

Sood S, V Bhardwaj, V Kumar, R Das, VK Gupta, V Mangal, SK Luthra, R Kumar, S Kumar, RK Singh, M Kumar, A Rathore, NK Pandey and AK Singh (2022) Genetic progress in 50 years of potato breeding in India: where do we stand? *Euro. J. Agron.* 141: 126620.

30. ADJVN 01/AAD (IC648626; INGR23064), an arecanut germplasm with noticeably short internodes, dark green leaves, shorter inflorescences and highly fragrant flowers

B Augustine Jerard^{1*}, V Damodaran¹, I Jaisankar¹, Niral V² and NR Nagaraj³

¹ICAR-Central Island Agricultural Research Institute, Port Blair-744105, Andaman & Nicobar Islands, India

²ICAR-Central Plantation Crops Research Institute, Kasaragod-671124, Kerala, India

³ICAR-CPCRI Regional station, Dakshina Kannada District -574243, Karnataka, India

*Email: augustinejerard.b@icar.org.in

Arecanut (*Areca catechu* L.) palm is the predominantly grown crop in the Andaman and Nicobar Islands, next to the Coconut palm, which gives much remuneration to farmers. Characterization of a dwarf arecanut, which was collected and conserved at ICAR-CIARI for palm morphology, revealed that the dwarf type has noticeably short internodes, dense crown with closely arranged leaves and short inflorescences producing bolder nuts. Observations on seedling parameters revealed that more than 80% of the progenies of these dwarf type palms are found to be uniform for seedling traits such as germination, seedling height, number of leaves and collar girth of seedlings. The unique arecanut dwarf accession has immense potential for utilization in breeding for dwarfness in arecanut and has a great opportunity to use for ornamental planting owing to the beautiful appearance of foliage and bunches at an incredibly low height. The palms are observed to produce bunches within 60 cm of height with closely arranged leaf scars. Other unique features, such as compact canopy with shorter, dark green leaves, shorter inflorescences, and highly fragrant flowers, make this dwarf type distinct from the common tall cultivars.

The estimated chali yield of the dwarf arecanut palms ranged from 1.8 kg to 2.8 kg/palm/year. Comparison of 30-year-old tall and this dwarf type palms revealed that the dwarf palms reached a height of 3 to 5.3 m, whereas the tall type palms reached over 15 to 20 m height. A compact block of this dwarf arecanut is established at the Sipighat farm of ICAR-

CIARI. The seed nuts of this unique type were deposited with the National Active Germplasm Site for arecanut, ICAR-CPCRI, Kasaragod, for conservation at the National Gene Bank.

Characterization of a dwarf arecanut AAD-1 was continued, which was collected and conserved at ICAR-CIARI. Observations on seedling parameters revealed that more than 80% of the progenies of these dwarf type palms were uniform for seedling traits such as germination, seedling height, number of leaves and collar girth. The progressive growth rate of AAD-1 seedlings was compared with Samrudhi and Mangala arecanut seedlings. Significantly lower height increment was noticed in AAD-1 in 11 months, whereas number of leaves produced was nonsignificant, indicating the dwarfness of AAD-1 at early seedling stage itself. The AAD-1 seedlings showed 50% lesser height in the initial stages of growth up to 3 to 4-leaf stage and also recorded higher collar girth and produced dark green leaves than Samrudhi and Mangala. Preliminary observations on growth parameters recorded in the compact block of this dwarf arecanut, along with the hybrids using Dwarf with local and the local tall have revealed marked differences in the leaf length, petiole length, leaf lamina and palm height. The hybrid seedlings have exhibited vigorous growth over the dwarf parent. About 200 seed nuts were sown in poly bags nursery for seedling production to establish a new block of dwarf arecanut at the Institute.

31. IIRR-BIO-SB-9 (IC646828; INGR23065), a rice germplasm with tolerance to yellow stem borer

MS Madhav^{1*}, AP Padma Kumari¹, RM Sundaram¹, GS Laha¹, LV Subba Rao¹, P Senguttuvel¹, HK Patel², Ramesh V Sonti², B Vishalakshi¹, B Suneel¹, B Umakanth¹ and P Gopi¹

¹ICAR-Indian Institute of Rice Research, Hyderabad-500030, Telangana, India

²CSIR-Center for Cellular and Molecular Biology, Hyderabad-500007, Telangana, India

*Email: sheshu24@gmail.com

Yellow stem borer (YSB) is a major pest of rice (*Oryza sativa* L.) and the host plant resistance cannot be exploited due to the unavailability of resistance sources in the rice gene pool. To obtain novel resistant/tolerant sources for YSB, a 10,500 EMS-induced mutant population was developed in the genetic background of Samba Mahsuri (BPT 5204) in collaboration with CSIR-CCMB. After screening of 5000 M₂ families, a mutant line (IIRR-BIO-SB-9) was found to have tolerance to YSB. The selection of the IIRR-BIO-SB-9 was done in M₂ generation after screening every season, selected single tolerant plant in each generation and advancing to M₈ through panicle to row method.

For the estimation of genomic similarity with wild type, a total of 120 SSR markers (selected based on ~1 marker per 5Mb) which were spread uniformly across the twelve linkage groups were used. The results indicated that IIRR-BIO-SB-9 has 95% genomic similarity with wild type (BPT 5204 (Padmakumari *et al.*, 2017). It was evaluated for the agro-morphological variations at the Indian Institute of Rice Research Farm (ICAR-IIRR) and ICRISAT.

The mutagenized population (5000 M₂ lines) were screened for yellow stem borer tolerance in a phased manner at both vegetative and reproductive phases by augmenting the natural pest infestation through artificial releases. Damage was assessed at both dead hearts (symptoms at vegetative phase of rice plant) and white heads (symptoms at reproductive stage) at each

of the crop growth stages on the Standard Evaluation System (IRRI, 2002) SES scale. Evaluation of the M₂ lines, advancement of the promising lines and reconfirmation of the promising lines were carried out simultaneously till M₈ generation and the most promising line IIRR- BIO-SB-9. Further, IIRR-BIO-SB-9 was screened in a stem borer screening trial, a multi-location trial under AICRIP- Entomology Programme for three years (2015-2017). The results indicated that IIRR-BIO-SB-9 exhibited tolerance to YSB.

To identify mutated loci for conferring resistance to YSB, MutMap (NGS-based approach) analysis was performed in resistant and susceptible bulks of the F₂ population derived from SM-92 X BPT 5204. The identification and characterization of mutated loci responsible for YSB tolerance indicated three possible genomic regions (chromosomes -1, 10 & 12) of mutations. The identified mutant line can serve as the best genetic resource in the rice breeding programme, as there were no reports on the existence of high-tolerance-level genetic material in rice. Thus, IIRR- BIO-SB-9 can be used as a donor in rice improvement programmes for YSB tolerance.

Reference

Padmakumari AP, M Sheshu Madhav, B Umakanth, B Sunil, P Gopi, J Karteek, GS Laha, RM Sundaram, LV Subbarao, HK Patel and RV Sonti (2017) EMS mutants as novel source of tolerance to rice yellow stem borer. IIRR Newsletter. April- September. Vol. 15, No. 2&3.

32. BHS 480 (IC646837; INGR23066), a barley germplasm with resistant to all pathotypes of leaf and stem rust at the seedling stage (except for race 11)

Madhu Patial^{1*}, Dharam Pal¹, SC Bhardwaj², KK Pramanick¹, OP Gangwar¹ and SK Bishnoi³

¹ICAR-IARI Regional Station, Shimla-171004, Himachal Pradesh, India

²ICAR-IIWBR, Regional Station, Flowerdale-171002, Himachal Pradesh, India

³IIWBR Seed and Research Farm, C/o CIRB, Hisar-125001, Haryana, India

*Email: mcaquarian@gmail.com

BHS 480 (BBM 803) is a barley (*Hordeum vulgare* L.) line resistant to leaf and stem rust. It has shown seedling resistance against all the prevailing pathotypes of leaf

rust and stem rust (except race 11). BHS 480 (BBM 803) also showed adult plant resistance to stem rust with ACI less than 15, i.e. 5MS (Table 1). This line is developed

following the pedigree method of breeding involving crosses between barley local germplasm BLG 132 and registered barley genetic stock BHS 369 at ICAR-IARI, Regional Station, CHC, Amartara Cottage, Tutikandi Centre, Shimla (H.P.)

BHS 480 (BBM 803) has a semi-spreading growth habit with medium maturity (166 days) under Northern

Hill conditions. The average yield is 3.28 t/ha under rainfed conditions of the Northern Hill Zone of All India Co-ordinated trials 2019-20. The distinguishing features of BHS 480 (BBM 803) are six-rowed, hulled; semi-spreading growth habit; erect flag leaf attitude; green leaves and thousand grain weight of 43 g.

Table 1. Meteorological data from different locations used in the study

Diseases	Condition	Year	Response of Proposed Genetic stock BHS 480 (BBM 803)	Page No.
Leaf Rust (Resistant to brown rust)	NBDSN (Seedling)	2019-20	R (Resistant to all races)	3.28 of Reference (i)
Stem Rust (Resistant to black rust)	NBDSN (APR) NBDSN (Seedling)	2019-20 2019-20	HS=5MS R (Resistant) response to all races (Except for race 11)	3.14 of Reference (i) 3.28 of Reference (i)
Stripe rust (Yellow rust)	IBDSB	2018-19	Yellow rust ACI = 2.7 (Highest Score 5S)	3.7 of reference (ii)

ACI, Average Coefficient of incidence; HS, Highest Score; IBDSN, Initial Barley Disease Screening Nursery; NBDSN, National Barley Disease Screening nursery.

33. CV.WP-14 (IC251385; INGR23067), a blackgram germplasm with high resistance against *Callosobruchus chinensis* L.

Revanasidda A^{1*}, Aditya Pratap¹, Mohd. Akram¹, Basavaraja T¹, Sanjeev Gupta¹ and Latha M²

¹ICAR-Indian Institute of Pulses Research, Kanpur-208024, Uttar Pradesh, India

²ICAR-NBPGR Regional Station, Thrissur-680656, Kerala, India

*Email: rshlb1114@gmail.com

Bruchids (Coleoptera: Bruchinae) cause severe economic losses to stored pulses, and their management has remained challenging as conventional methods often prove ineffective across regions and time. In this context, host plant resistance offers an effective and environmentally friendly solution. Identifying resistant sources in traditional cultivars, landraces, or crop wild relatives (CWRs) and introgressing alien genes into cultivated legumes can diversify and strengthen the genetic base. Several wild legumes, particularly in the *Vigna* subgenus *Ceratotropis*, have been explored for resistance to biotic and abiotic stresses. Bruchid resistance was first identified in wild mungbean accession TC1966 against *Callosobruchus chinensis* and later in other cultivated and wild *Vigna* germplasms. Crosses between resistant wild accessions and cultivated types were attempted to transfer resistance genes (Mariyammal *et al.*, 2019). The susceptible mungbean cultivar VRM (Gg) 1 with the resistant rice bean cultivar TNAU Red, identified several QTLs linked to bruchid resistance. Despite these

advances, studies on diverse genetic resources of wild *Vigna* species and their transfer to cultivated forms remain limited, warranting further research.

India, being the center of diversity for *Vigna*, harbours a rich repository of wild, weedy, and cultivated forms. Gene banks across the country maintain large collections of wild species belonging to the subgenus *Ceratotropis*. However, these remain underexplored for bruchid resistance. To address this gap, the present study screened diverse endemic *Vigna* accessions to identify resistant donors against three prevalent bruchid species in India.

Between 2019–21, 42 wild *Vigna* accessions representing 13 species were evaluated against *C. chinensis*, *C. maculatus*, and *C. analis* at the Storage Entomology Laboratory, Division of Crop Protection, ICAR-Indian Institute of Pulses Research (IIPR), Kanpur. Seeds were sourced from the Germplasm Bank, Division of Crop Improvement, ICAR-IIPR. Qualitative and quantitative seed traits were recorded following NBPGR descriptors to assess their role in resistance. Screening

involved both 'Free-choice' and 'No-choice' tests (Eker *et al.*, 2018), with resistant accessions validated under 'No-choice' conditions. Reactions were categorized based on per cent seed damage (Eker *et al.*, 2018).

Among the accessions, IC251385 of *V. mungo* exhibited complete resistance to *C. chinensis*, recording 0% seed damage in both initial and confirmatory screenings, despite egg deposition averaging 1.7, 0.8, and 1.38 per seed. Correlation studies revealed that seed morphological traits did not significantly influence resistance, suggesting biochemical components in the seed coat and cotyledons play a major role. This accession thus represents an invaluable genetic resource for introgression breeding in mungbean and urdbean. Identification of QTLs associated with bruchid resistance in IC251385 further

strengthens its utility. Deploying these resistance genes through introgression breeding could lead to the development of superior bruchid-resistant cultivars, ensuring sustainable pulse storage and reduced economic losses.

References

- Eker T, F Erler, A Adak, B Imrek, G Guven, HS Tosun, D Sari, H Sari, HD Upadhyaya, C Toker and C Ikten (2018) Screening of chickpea accessions for resistance against the pulse beetle, *Callosobruchus chinensis* L. (Coleoptera: Bruchidae). *J. Stored Prod. Res.* 76: 51-57.
- Mariyammal I, D Seram, SM Samyuktha, A Karthikeyan, M Dhasarathan, J Murukarthick, JS Kennedy, D Malarvizhi, T Yang, M Pandiyan and N Senthil (2019) QTL mapping in *Vigna radiata* × *Vigna umbellata* population uncovers major genomic regions associated with bruchid resistance. *Mol. Breed.* 39: 110.

AUTHOR GUIDELINES

The Indian Journal of Plant Genetic Resources (IJPGR) is the official publication of the Indian Society of Plant Genetic Resources (ISPGR). For publication in the journal, the authors must be a member of the Society. UPGR publishes full-length papers or short communications of original scientific research in the field of plant genetic resources (PGR). Review articles (with prior consent or invitation only) summarizing the existing state of knowledge in topics related to PGR will also be published.

Contributions should be as concise as possible. The maximum length of the review article, full-length papers and short communications is usually restricted to 12, 6, 3 printed pages including illustrations and tables, respectively.

SCOPE

Basic research on biosystematics, genetics and genomics related to PGR

Applied research on field and laboratory evaluation of PGR

Supportive research in conservation and quarantine of PGR

Policy research on access and benefit sharing; IPRS

Status reports on ecology, conservation, traditional knowledge and use of PGR

Title Page: The title page of the manuscript should be the first page and should include the title, names and addresses of the authors, abstract and keywords.

Title: Keep the title brief, specific and informative and amendable to indexing. It should be typed in running text with first letter of word as capital and Latin names in italics.

Name and Address: The name of authors and the address of the institution where the work was carried out should be mentioned below the title. Present address of correspondence, if different, should be given as footnote indicating by asterisk (*) the author to whom the correspondence and reprint requests are to be made. E-mail addresses should also be indicated.

Abstract: The abstract should clearly state the rationale, objectives, methods, and important conclusions of the study. It should not exceed 150 words.

Key Words: The abstract should be followed by not more than five key words (following alphabetic order) indicating the contents of the paper and useful for abstracting purposes.

Main Text: The main text of the paper should start from the second page which should contain the title of the paper followed by the text divided into following main headings which are to be typed in running text and flushed with the margin: Introduction, Materials and Methods, Results, Discussion, Acknowledgements, References. Only relevant references should be included in reference section. Wherever appropriate, results and discussion can be combined and acknowledgements be omitted.

Introduction should be brief and limited to the statement of the problem and aim of the experiment.

Materials and Methods should include relevant details on the nature of material, experimental design, the techniques/methodology employed and the statistical method used. For well-known methods, citation of reference will suffice.

Results and Discussion should be clear to readers in different disciplines. Units of measurement should be SI.

Tables should be typed on separate sheets, each with a heading stating its contents clearly and concisely. Numerical data and calculations should be thoroughly checked.

Figures of only good quality that are essential to a clear understanding of the paper shall be accepted. Legends to the illustrations should be typed on separate paper. Information in the legend should not be repeated in the text and similarly, the same data should not be represented in both graph and table form. All figures, whether photographs, maps, graphs or line drawings should be numbered consecutively. Illustration number and title of the article with authors' name may be given at the back of the plates in soft pencil.

Line diagrams of high quality, preferable in the desired final size would be accepted. The inscriptions should be clearly legible.

Photographs for publication should be of high contrast, black and white/coloured, glossy print, trimmed at right angles. Magnification should be indicated with a bar scale on the photo. Authors need to indicate colour reproduction of photographs. Color photographs will be published free of cost however, cost of colour printing will be borne by the authors, if hard copy is required by author(s).

Acknowledgements should mention only guidance or assistance received in real terms, and financial grant provided by a funding agency. Acknowledgements for inspiration, typing etc., need not be mentioned.

References in text should be cited by author, year of publication (e.g. Joshi, 1995; Joshi and Rao, 1995) and multiple citations should be in chronological order (Withers and Englemann, 1998; Rao et al., 2001). References should be listed in alphabetical order under the first authors' name. The names of journals should be abbreviated according to the latest edition of the World List of Scientific Periodicals (eds P Brown and GB Stratton), Butterworths, London.

The following examples may be used for citations:

- Chao CCT and RR Krueger (2007) The date palm (*Phoenix dactylifera*); an overview of biology, uses and cultivation. *Hort. Sci.* 42(5): 1077-82.
- Muralidharan CM, SBS Tikka and P Verma (2008) Date palm cultivation in Kachchh, Technical Bulletin-2, Date palm Research Station, Sardarkrushinagar Dantiwada Agricultural University (SDAU), Mundra, Gujrat, India, p36.
- Wither LA and F Englemann (1998) In vitro conservation of plant genetic resources. In: A Altman (ed) *Agricultural Biotechnology*. Marcell Dekker Inc., New York, pp 57-58.
- WOI (1985) *Wealth of India - Raw Materials. A Dictionary of Indian Material and Industry Products - Raw Material Vol 1: A (Revised)*. Publication and Information Directorate, Council of Scientific and Industrial Research, New Delhi, 513 p.
- Engels JMM and V Ramanatha Rao (eds) (1998) *Regeneration of seed crops and their wild relatives*. Proceedings of a Consultation Meeting, 4-7 December 1995, ICRISAT, Hyderabad and IPGRI, Rome, Italy, 167 p.

For Journal title and Abbreviations, in general, authors are advised to follow Elsevier System (https://legacyfileshare.elsevier.com/promis_misc/BMCL_Abbreviations.pdf).

Short Communications

The style and format as mentioned for full-length papers should be followed for Short Communications. However, the abstract should be restricted to not more than 50 words and the remainder text should be continuous (without headings). Illustrative material should be kept at minimum, usually not more than one table or figure and only few references should be included (not more than 10). Authors can also submit meeting reports after consulting Editor-in-Chief, IJPGR.

Submission of the Manuscript Submission of an article will be held to imply that it has not been previously published or submitted for publication elsewhere. A cover letter including a statement to this effect should be submitted with the manuscript. Article are submitted through online submission system:

https://ispgr.in/index.php/ijpgr/about/submissions=manuscript_submission

Authors are requested to provide details (Name, designation, affiliation, specialization, email ID, phone number) of at least 4 experts/reviewers in the areas of your studies, in a separate MS Words file.

- Experts in the subject will review all the submitted manuscripts and the final decision about the acceptance of the manuscript rests with the Editorial Board.
- Publication of a paper in the Journal does not imply the responsibility for an agreement with the statements or view written therein, and rests entirely on the authors thereof.
- The authors will receive page proofs, which should be corrected and returned maximum within 3 days. Corrections must be kept to the minimum, and the proof stage should not be regarded as an opportunity for further editing and additions. Although, every effort is made by the editors to correct proofs of all the papers they assume no responsibility for errors that may remain in the final print.

ETHICS STATEMENT

IJPGR approves the spirit of the guidelines for journal editors developed by the Committee on Publication Ethics (COPE).

- Manuscripts submitted to IJPGR are evaluated entirely on the basis of their scientific content and relevance to PGR research community.
- IJPGR does not levy any publication charges to members of ISPGR.
- ISPGR, the publishers of the Journal, take all possible measures to uphold the highest standards of publication ethics and to prevent any malpractices.
- Authors who submit manuscripts to IJPGR declare that the submissions are original and unpublished and are not under consideration for publication elsewhere.
- Authors also declare that the manuscript is based on their own original work devoid of any nature and amount of plagiarism.
- IJPGR makes full efforts to eliminate any kind of conflict of interest of authors, editors or reviewers, resulting from competitive, collaborative or other relationship with any person, institution or agency connected to the submissions.

Ethics expected to be followed by IJPGR Editors:

- IJPGR Editorial Team is fully responsible for the decision on acceptance or rejection of a manuscript submitted to the Journal. An established procedure is followed to ensure unbiased decision making. Submissions are handled by different Editors based on their specialization. Editors obtain the reports of three referees with expertise in the topic dealt in the submission. Each Editor recommends rejection or acceptance (direct or after revisions) to the Editor-in-Chief who takes the final decision.
- The evaluation of manuscripts is made on the basis of their scholarly content in terms of relevance, novelty and scope as well as their presentation in terms of language and syntax. IJPGR does not consider any other factors like gender, race, religious belief, ethnic origin, citizenship, or political philosophy of the authors.
- IJPGR strictly follows transparency in the evaluation process and confidentiality of the submission. Reviewers, as on today, remain anonymous to authors.

Ethics expected to be followed by IJPGR Reviewers

- Peer review is the backbone of IJPGR manuscript flow. IJPGR depends upon the services of reviewers for maintaining ethical integrity and scholarly quality.
- IJPGR does not follow double blind review and, therefore, the reviewers are expected to maintain absolute confidentiality with regard to the contents of manuscripts.
- The reviews are conducted objectively with every comment/decision supported by clear reasons. Reviewers follow a non-ranking format provided by the IJPGR to facilitate decision making by Editor.

Ethics expected to be followed by Authors

- Authorship should be restricted to only those persons who have made a significant contribution to the conception, design, execution, or interpretation of the reported study. In recent times, symbiotic authorship has become a serious ethical issue which should be avoided.
- The authors must ensure that the experiment and manuscript are both entirely original works written in good English without any ambiguity. Every instance of use of work and/or words of other authors must be appropriately cited or quoted. Any kind of plagiarism detected at review stage results in rejection of the manuscript. If the plagiarism is detected post-publication, the article will be summarily retracted. If any fundamental errors are found in published papers, authors will inform IJPGR which will take corrective measures including retraction.
- IJPGR persuades authors to provide the basic data related to manuscripts such as description of germplasm and its availability to ensure greater utilization of diverse germplasm. Authors are also encouraged to provide relevant raw data for editorial review. Authors must prepare such data in a format to provide public access. Specific requests of data confidentiality shall be entertained by IJPGR on case-to-case basis.

- Authors must avoid any kind of duplication in their submissions i.e. attempting to publish same/significantly similar manuscripts in more than one journal. IJPGR does not consider the following as a prior publication: (i) Abstract in a conference/symposium; (ii) Academic thesis/dissertation; (iii) Invited talks and interviews on specific topics.
- Submission to IJPGR implies that authors have complied with ethical and technical standards of use of hazardous material, use of animal or human subjects, access and use of genetic resources, access and use of IP protected data. It is authors' duty to ensure no national or international legal requirement is compromised.
- Submission of a manuscript to IJPGR implies that the corresponding author has ensured that all co-authors have seen and approved the final version of the paper and have agreed to its submission for publication to IJPGR. All sources of financial support should also be disclosed.

PUBLISHER/EDITORIAL OFFICE ADDRESS

Office Block A, 2nd Floor, National Agricultural Science Complex, Dev Prakash Shastri (DPS) Marg, New Delhi-110012, India **E-mail:** ispgr2015@gmail.com

Publisher website: www.nbpgr.ernet.in/ispgr

Journal hosting and online submission of manuscript: <https://epubs.icar.org.in/index.php/IJPGR> Online Access to IJPGR

All the issues of Indian Journal of Plant Genetic Resources from 1988 [Vol 1(1)] till date [Vol 39 (2)] are accessible as pdf files of individual papers.


- The access is provided to life members only
- Access is through user ID and password
- Login credentials have been emailed to all life members by ispgr.in (if any life member has not received the user ID and password, please contact IJPGR office via E-mail: ispgr2015@gmail.com)
- Open the home page of IJPGR at <https://epubs.icar.org.in/index.php/IJPGR>
- You may also reach IJPGR home page via Google search of "Indian Journal of Plant Genetic Resources"
- Click on the LOGIN button and enter the user ID and password
- If you are a new member, then click on the REGISTER option, and register as AUTHOR.
- For a new submission, click on MAKE A SUBMISSION, or check our AUTHOR GUIDELINES.

Online Access to IJPGR

All the issues of Indian Journal of Plant Genetic Resources from 1988 [Vol 1 (1)] till date Vol 39 (2)] are accessible as pdf files of individual papers.



- IJPGR is Open access journal; Life Member of ISPGR can also access through user ID and password
- Login credentials have been emailed to all Life Members by ispgr.in (if any Life Member has not received the user ID and password, please contact IJPGR Office at ispgr2015@gmail.com)



- Open the home page of IJPGR at <https://epubs.icar.org.in/index.php/IJPGR>. You may also reach IJPGR home page via Google search of "Indian Journal of Plant Genetic Resources"
- Click on the LOGIN button and enter the user ID and password
- If you are a new member, then click on the REGISTER option, and register as AUTHOR
- For a new submission, click on MAKE A SUBMISSION, or check our AUTHOR GUIDELINES
- The login credentials are for the personal use of IJPGR life members. Please do not share with others

CONTENTS

Review Article

- 1 **Sustainability Benefits Following the Adoption of Genetically Modified Crops in Canada** 1
Stuart J Smyth

Research Articles

- 2 **Deciphering Genetic Variability and Stability for Yield Improvement in Mungbean (*Vigna radiata*)** 9
Chandra Mohan Singh, Mukul Kumar, Hitesh Kumar, Priyanka Kaundal, Chirag, Pradyum Maurya, Ruchi, Anurag and Aditya Pratap
- 3 **Aroma and Agro-morphological Diversity in Traditional Rice (*Oryza sativa*) Germplasm from Eastern India** 20
Pooja Belwal, Amrita Banerjee, Anima Mahato, Priyamedha, Sougata Bhattacharya, Nabaneeta Basak, Nimai Prasad Mandal and Somnath Roy
- 4 **Assessment of Molecular Diversity and Phylogenetic Analysis Among Chilli Genotypes Using Start Codon Targeted (SCoT) Markers** 29
Hanuman Ram, DK Samadia, Chet Ram and AK Verma
- 5 **Assessment of Antioxidant Activity and Quality Attributes in Indigenous Pran (Kashmiri Shallot) Genotypes** 38
Gazala Nazir, Suraya Ahsan, Mudasir Magray, Khursheed Hussain, Saima Tabasum, UH Masoodi, Syed Berges, Asima Amin, Abida Jabeen and AA Lone
- 6 **Morpho-biochemical Characterization and Multivariate Assessment of Genetic Diversity in Rajmah (*Phaseolus vulgaris* L.) Genotypes from Northeast India** 43
Bedashruti Saikia, Rupam Borgohain Gayatree Goswami, Jutika Das and Anubrat Borah
- 7 **Fatty Acid Composition of Seeds of Three Native *Garcinia* Species from Andaman and Nicobar Islands, India** 53
Pooja Bohra and Ajit Arun Waman
- 8 **Evaluation of Barley (*Hordeum vulgare* L.) Germplasm for Heat Tolerance Using Germination Indices and Seedling Traits** 58
Mohmad Amin Bhagat, Navjyot Kaur, Prinka Goyal and Simarjit Kaur
- 9 **Effect of Various Seed Priming Agents on Germination, Seedling Vigour and Grain Yield of Dry Direct-Seeded Rice in the Northwestern Indo Gangetic Plains of India** 67
Sukham Madaan, Mangat Ram, Lalita, Sumit Saini, Ashish Jain, Rakesh Kumar, Mahaveer Bochalya, Amit Kumar, Ashwani Kumar, Vishal Gandhi, Charan Singh, Parvesh Kumar, Sarita Devi and Anita Kumari
- 10 **Morphological Diversity Analysis in Brinjal (*Solanum melongena*) Germplasm** 74
TH Singh, Sushil Pandey, Chitra D Pandey and Suresh R Yerasu
- 11 **Preliminary Characterization of Wild *Actinidia Callosa* Var. *Strigillosa* as Potential Genetic Resources from North Sikkim, India** 83
Abhilash Padhan, Dinesh Singh Thakur, Vishal Singh Rana, RK Dogra, Narender Negi, Akriti Chauhan and Pooja Sharma
- 12 **Present status of *Pubistylus andamanensis* Thoth. in Andaman Islands and its IUCN Red List Assessment** 100
CS Purohit, Vivek CP, BC Dey and LJ Singh
- 13 **Correlation and Path Analysis for Seed Yield and Components Traits in Mothbean [*Vigna aconitifolia* (Jacq) Marechal] under Arid Environment** 105
Savita Meena, NK Sharma, Pinky Yadav, Rukoo Chawla and Ajit Kumar Meena
- 14 **Genetic Variation and Elite Clone Selection in *Gmelina arborea* for Seed Source Development in Northeast India** 114
Manish Kumar Singh
- 15 **Exploring Maize (*Zea mays* L.) Landraces: Characterization for Yield and Quality Traits to Foster Genetic Improvement** 125
Heena Sharma, Surinder Sandhu, Gagandeep Kaur and RK Bhavyasree

Book Review

- 16 **A Review of Code Paroda: A Life in Service of Agricultural Science** 135
A Amarender Reddy
- 17 **Plant Germplasm Registration Notice** 143
- 18 **Authors Guidelines** 169